

CAMPBELL REVIEW:

COGNITIVE BEHAVIOURAL THERAPY FOR MEN WHO PHYSICALLY ABUSE THEIR FEMALE PARTNER

Smedslund G, Dalsbø TK, Steiro AK, Winsvold A & Clench-Aas J

ISBN: 978-87-7487-880-3

2007 Nordic Campbell Center

This review is also published within the Cochrane Library

Nordic Campbell Center

Herluf Trolles Gade 11

1052 København K

Tel: +45 3369 7770

nc2@sfi.dk

www.nc2.net

Nordic Campbell Center's publications
should be cited with clear indication of source.

Please send a copy of references made to this
Campbell Review to Nordic Campbell Center.

Cover sheet

Title

Cognitive behavioural therapy for men who physically abuse their female partner

Reviewers

Smedslund G, Dalsbø TK, Steiro AK, Winsvold A, Clench-Aas J

Dates

Date edited: 13/07/2007
Date of last substantive update: 24/04/2007
Date of last minor update: 22/05/2007
Date next stage expected: 22/05/2009
Protocol first published: Issue 2, 2006
Review first published: Issue 3, 2007

Contact reviewer:

Geir Smedslund
Norwegian Knowledge Centre for Health Services
PB 7004 St. Olavs plass
Oslo
N-0130 NORWAY
Telephone 1: 47 2325 5155
Telephone 2: 47 9138 7076
Facsimile: 47 2325 5010
E-mail: ges@nokc.no, Geir.Smedslund@kunnskapssenteret.no

Internal sources of support

Norwegian Knowledge Centre for the Health Services, NORWAY

External sources of support

Nordic Campbell Centre, DENMARK

Contribution of reviewers

Dalsbo and Smedslund wrote the protocol.

All reviewers independently screened literature, reviewed potential trials, and extracted data. Smedslund analysed the data. Smedslund wrote the text of the completed systematic review. Steiro, Winsvold and Clench-Aas contributed by giving comments, assessing studies and acting as mediators if necessary. Responsibility for updating the review is jointly shared between Dalsbo and Smedslund.

Acknowledgements

Many thanks to Torill Johme for contributing to the planning of the study, for literature screening, hand-searching, and for performing the initial searches in 2003.

Thanks to Sigrun Espelien Aasen for carrying out updated searches in November-December 2006.

We also want to acknowledge the Norwegian Directorate for Health and Social Affairs, which

housed us and funded us during the initial phase of the development of this review. We are also grateful for the support by Andy Oxman during the start-up. Last, but not least, we want to thank Jane Dennis, the review group coordinator of the Cochrane Developmental, Psychosocial and Learning Problems Group for her continuous support and help in dealing with a number of different review authors and with different drafts leading up to the final review.

This review is co-registered within the Cochrane Developmental, Psychosocial and Learning Problems Group (Cochrane Collaboration).

Potential conflict of interest

None known.

What's new

Dates

Protocol first published:	Issue 2, 2006
Review first published:	Issue 3, 2007
Date of last substantive update:	24/04/2007
Date of last minor update:	22/05/2007
Date review re-formatted:	/ /
Date new studies sought but none found:	/ /
Date new studies found but not yet included/excluded:	/ /
Date new studies found and included or excluded:	/ /
Date reviewers' conclusions section amended:	/ /
Date comment/criticism added:	/ /
Date response to comment/criticism added:	/ /

Synopsis

Cognitive behavioural therapy for men who physically abuse their female partner

Domestic violence is common and serious. Many convicted men are court-ordered to receive cognitive behavioural therapy (CBT) in order to end their violence. Most men do not voluntarily seek such therapy but participate either after being pressured by their partners or after being legally required to do so. The review sought all randomised controlled evaluations about the effects of CBT on physical violence worldwide, but found only six small trials with 2343 participants met the inclusion criteria. The evidence from the included studies is insufficient to draw any conclusions.

Abstract

Background

In national surveys between 10 % and 34 % of women have reported being physically assaulted by an intimate male partner. Cognitive behavioural therapy (CBT) or programmes with elements of CBT are frequently used treatments for physically abusive men. Participants either enrol voluntarily or are obliged to participate in CBT by means of a court order. CBT not only seeks to change behaviour using established behavioural strategies, but also targets the thinking patterns and beliefs that are thought to contribute to violence.

Objectives

To measure the effects of CBT and similar interventions on men's physical abuse of their female partners.

Search strategy

We searched the Cochrane Controlled Trials Register (CENTRAL), C2-SPECTR, MEDLINE, EMBASE, CINAHL, PsycINFO, ERIC, Care Data/Social Care Online, Sociological Abstracts, Criminal Justice Abstracts, Bibliography of Nordic Criminology (all to late 2006), and SIGLE to 2003. *Santé mentale au Québec* was handsearched from 1976 to 2003 and reference lists of articles. We also contacted field experts and the authors of included studies.

Selection criteria

Randomised controlled trials (including cluster-randomised and quasi-randomised trials) of cognitive behavioural therapy with men who physically abuse their partners and reporting effects on continued violence.

Data collection & analysis

Two review authors independently assessed trial quality and extracted data. We contacted study authors for additional information.

Main results

Six trials, all from the USA, involving 2343 people, were included. A meta-analysis of four trials comparing CBT with a no-intervention control with 1771 participants, reported that the relative risk of violence was 0.86 (favouring the intervention group) with a 95% confidence interval (95% CI) of 0.54 to 1.38. This is a small effect size, and the confidence interval is so wide that there is no clear evidence for an effect. One study (Wisconsin Study) compared CBT with process-psychodynamic group treatment and found a relative risk of new violence of 1.07 (95% CI 0.68 to 1.68). Even though the process-psychodynamic treatment did marginally better than CBT, this result is also equivocal. Finally, one small study (N = 64) compared a combined treatment for substance abuse and domestic violence (SADV) with a Twelve-Step Facilitation (TSF) group. An analysis involving 58 participants investigated the effect on reduction in frequency of physical violence episodes. The effect size was 0.30 (favouring TSF) with 95% confidence interval from -0.22 to 0.81.

Reviewers' conclusions

There are still too few randomised controlled effect evaluations to conclude about the effects of cognitive behaviour therapy on domestic violence.

Background

Violent behaviour constitutes a serious problem in societies worldwide. Intimate partner abuse is especially problematic because it takes place in the private family sphere, making it a difficult arena for intervention and help. The physical abuse of women by their male partner is a serious concern because "it affects a distressingly high percentage of the population and it results in physical, psychological, social, and economic consequences" (CDCP 2003). The World Health Organisation (WHO) reported that "the overwhelming health burden of partner violence is borne by women at the hands of men" (WHO 2002). The WHO also provided evidence about the extent of the problem: in national surveys between 10 % and 34 % of the women reported being physically assaulted by an intimate male partner (WHO 2002).

Domestic violence occurs in the family and takes many different forms, including sexual, psychological, emotional and physical abuse. In this review the focus is solely on physical abuse. Domestic violence can occur between spouses/partners and between adults and children in the family. This review focuses only on partnership abuse, and specifically on men who physically abuse their female partner or ex-partner. The term domestic violence is therefore too broad to give meaning for this review, and more useful terms are physical abuse, battering, and intimate partner abuse. Another important limitation is that this review does not focus on the causes of violent behaviour. Several biological, psychological and sociological studies have attempted to find the one answer to what causes men to commit violent actions. In general there is now more focus on the correlation of different behavioural variables leading to violence. Therefore this review sets out to include more outcome variables than strictly physical violence, for example self-esteem, substance use and emotional problems.

US batterer intervention programmes were initially introduced as an alternative to incarceration for men arrested for domestic violence. This was a controversial strategy because many policy makers/advocates felt that this diversion sent the message that abuse was a mental health problem rather than a crime, and because funding these programmes may have diverted funds from programmes for abused women. The stated goals for the batterer intervention programmes differ widely, from statistically measurable reductions in violence, to holding men accountable to preparing men to take action against the woman battering culture.

One of the most frequently used treatment programmes for physically abusive men is a psychological intervention called cognitive behavioural therapy (CBT). CBT not only seeks to change behaviour using established behavioural strategies, but also targets the thinking patterns and beliefs that are thought to contribute to violence. CBT is "designed to help the patient test certain maladaptive cognitions and assumptions" (Beck 1979). CBT techniques aim to identify thoughts and beliefs that precede violent behaviour, challenging the patterns that violent men use to justify their violence after the event. The goal is to bring about changes in the way that physically abusive men think about violence and the circumstances which lead to violence, thereby interrupting the chain of events that lead to physical abuse. The CBT can be given in individual, couple or group format. A common intervention called the Duluth Model (Pence 1993) has many cognitive-behavioural components and is included in this review.

An American review of state and provincial programmes for intervening in spouse abuse cases reported simply "the jury remains out on the effectiveness of these programmes" (Arias 2002). When spouse abusers are sent on programmes, it is important to know the positive or negative effect. If a programme does not work or has adverse outcomes, we risk putting women in danger of future abuse.

The scope of this review is to determine the effectiveness of cognitive behavioural therapy

delivered to men engaged in physical abuse, against their female partner. A previous review of cognitive therapy (Butler 2000) for violent offenders did not include physically violent spouse abusers, but concluded that the therapy had a beneficial effect for those with problems such as marital distress and anger. To date, we know of no systematic review of the effects of CBT for men who are physically violent toward their partners that has employed a search strategy aiming to locate every randomised controlled study worldwide regarding this question. Davis and Taylor (Davis 1999) reviewed the literature but did not report a search strategy. A later review (Babcock 2004) searched only PsychINFO and used only four search terms. The results from the present review are of importance for perpetrators and victims of this form of violence, and those who seek this form of treatment for the problem, and also for therapists, researchers, the judiciary, and the general public.

Objectives

To measure the effects of cognitive behavioural therapy (CBT) and similar interventions on men's physical abuse of their female partners.

Criteria for considering studies for this review

Types of studies

Randomised controlled trials, including quasi- and cluster randomised controlled trials, were included in this review. The control group consisted of persons who received no intervention, other interventions, or were on a waiting list.

Types of participants

Men who physically abuse their female partner/spouse/wife. Primary studies where the focus is on women who abuse their partner/spouse were excluded from this review. In the event of trials having a mixed population of men who have been violent against women and those who have been violent against men, we requested data separately from the trial investigators. We recorded whether the perpetrators were living with their partners or whether they were ex-partners. Trials in which the participants attended the treatment programme voluntarily or were court-ordered to participate were included, and results were separately reported.

Types of interventions

Interventions stated by the authors to be cognitive behavioural or recognisably so (for example, Duluth Model) from the description provided. Programmes may be individual, couple or group based and delivered in any setting.

Types of outcome measures

The primary outcome measure was physically violent behaviour. We also included other violent behaviour, like verbal aggression and hostile attitudes. Other, secondary outcome measures were: improved self-esteem, reduced substance abuse and anger management. Regarding self-esteem, substance abuse, and emotional distress, these were recorded for both perpetrators and victims wherever possible. Measures of outcome data were grouped in to short follow-up time (0 to 6 months), intermediate 7 to 18 months, and long-term (19 months and beyond). Any formats for measuring the outcome were included but were separately reported (e.g. self reports, victim reports, judicial and police reports). Both standardised and unstandardised measures were included.

Search strategy for identification of studies

We searched the Cochrane Controlled Trials Register (CENTRAL) (The Cochrane Library Issue

4, 2006), MEDLINE (1966 to September week 3 2006), EMBASE (1980 to 2006 week 39), CINAHL (1982 to September 2006), PsycINFO (1806 to October 2006), ERIC (1966 to September 2006), C2-SPECTR (searched December 2006), Care Data/Social Care Online (searched September 2006), Sociological Abstracts (1963 to September 2006), Criminal Justice Abstracts (1968 to September 2003), Bibliography of Nordic Criminology (1999 to December 2006), and SIGLE (1980 to September 2003). Criminal Justice Abstracts and SIGLE were only searched until 2003 because we did not have access in 2006.

We handsearched *Santé mentale au Québec*, an online scientific journal, from 1976 to 2003. The review authors contacted field experts and the authors of retrieved studies in order to find additional studies. Conference papers were searched also, in order to minimise the threat of publication bias. Reference lists in included studies were searched for relevant literature. Studies were included regardless of language and country of origin.

Below is the search strategy used to search CENTRAL. The search strategies for the other electronic databases were modified to suit each database. They are included in [Table 02](#), [Table 03](#), [Table 04](#), [Table 05](#), [Table 06](#), [Table 07](#), [Table 08](#) and [Table 09](#).

Search strategy used in CENTRAL, The Cochrane Library Issue 4, 2006 (Sigrun Espelien Aasen)

```
#1MeSH descriptor Battered Women, this term only
#2MeSH descriptor Domestic Violence, this term only
#3MeSH descriptor Spouse Abuse, this term only
#4((family or domestic or conjugal or partner*) near/3 violence):ti,ab,kw
#5((abus* or batter* or beat* or assault*) near/3 (wom*n or partner* or spouse* or female* or wife or wives or domestic* or fiance or cohabitant* or live-in)):ti,ab,kw
#6((male* or men or man or partner* or spouse* or husband or fiance or cohabitant* or live-in) near/3 (batter* or perpetrator* or abus* or violen* or beat* or assault)):ti,ab,kw
#7(#1 OR #2 OR #3 OR #4 OR #5 OR #6)
#8MeSH descriptor Behavior Therapy, this term only
#9MeSH descriptor Cognitive Therapy, this term only
#10MeSH descriptor Psychotherapy, Rational-Emotive, this term only
#11(cognitive* near/3 (therap* or train* or techni* or question* or approach* or assessment*)):ti,ab,kw
#12((behavior* or behaviour*) near/3 (therap* or train* or modif* or experiment*)):ti,ab,kw
#13(rational* near/3 emotive*):ti,ab,kw
#14(cbt):ti,ab,kw
#15(schemas or schematas):ti,ab,kw
#16MeSH descriptor Imagery (Psychotherapy), this term only
#17(imager*):ti,ab,kw
#18((cognitive* or mental*) near/3 (map* or model*)):ti,ab,kw
#19(Socratic* near/3 (question* or method* or dialogue* or strateg* or sequence*)):ti,ab,kw
#20(dysfunctional near/2 (thought* or assumption* or rule* or appraisal* or belief* or attitude* or scheme*)):ti,ab,kw
#21(automatic near/3 (thought* or process*)):ti,ab,kw
#22(nat or nats):ti,ab,kw
#23(reattribution*):ti,ab,kw
#24((key or core) near/2 belief*):ti,ab,kw
#25(#8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24)
#26(#7 AND #25)
```

Methods of the review

Selection of studies

Selection of primary studies was based on the inclusion criteria described above. The Reference Manager database was transferred to SRS (software for electronic screening and data abstraction) (SRS 2005). At the first screening level, approval from a single review author took a citation to the second level, then two review authors working independently had to approve of a citation for it to be forwarded to Level 3 (and ordered in full text). Data from all relevant trials at Level 3 were extracted and presented in the 'Characteristics of included studies' table. All review authors contributed in this process. If two authors disagreed, a third author mediated, and the decision whether to include or exclude was reached through consensus.

Data extraction and management

Two review authors independently extracted data from the included studies using an online data extraction form. Any disagreement between review authors generated a conflict in SRS which had to be solved through a discussion. If disagreement persisted, a third review author was consulted. If outcome or other vital information was missing from the original reports, we contacted the author(s) by e-mail in an attempt to retrieve the necessary data for the analysis. The following data from the included studies were extracted:

Study characteristics: Country where the study was conducted, year of publication, publication type (e. g. journal article, report, book chapter).

Participants: age, socio-economic status, ethnicity, previous history of violent behaviour and treatment for it, current substance abuse, additional problems/disorders, and marital status.

Intervention: content, duration/time, profession of person delivering the programme (or intervention), gender and number of therapist(s)/group leader(s), support for women, the degree of mandatory delivery, attrition, adherence, type of comparison group (no intervention, other intervention).

Type of outcome measure: physical violence, aggression, self-esteem, substance abuse and managing anger.

Source of outcome data: official statistics; self-reports, partner report, or other forms for gathering outcome data.

Length of follow-up time: months and years.

Effect measures: standardised or unstandardised measures or raw data or both.

Quality assessments of included studies

Since "variation in validity can explain variation in the results of the studies included in a systematic review" (Higgins 2005) we assessed the internal validity of included studies. Two review authors independently assessed each selected study against quality categories described below. Uncertainty or disagreement was solved by discussion with a third reviewer. The review authors were not blinded to the authors or other information about the publication when assessing study validity. Whenever information about study quality, or other information about the study, was missing, we contacted the author(s) of the study, to minimise the danger of measuring the quality of the reporting, rather than of the study. Our aim was to get an overall assessment of internal validity based on a summary of the following six methodological criteria.

Generation of allocation sequence

MET = Resulting sequences are unpredictable (explicitly stated use of either computer-generated random numbers, table of random numbers, drawing lots or envelopes, coin tossing, shuffling cards, or throwing dice).

UNCLEAR = Statement that the study was randomised but not describing the generation of the allocation sequence.

NOT MET = Explicit description of inadequate generation of sequence, e. g. (e.g., using case record numbers, alternation, date of admission, date of birth).

Concealment of allocation sequence

MET = Neither participants nor investigators can foresee assignment (e.g. central randomisation performed at a site remote from trial location; or, use of sequentially numbered, sealed, opaque envelopes).

UNCLEAR = Statement that the study was randomised but not describing the concealment of allocation.

NOT MET = Explicit statement that allocation was not concealed OR statement indicating that participants and investigators can foresee upcoming assignment (e. g., open allocation schedule, unsealed or non-opaque envelopes).

Prevention of performance bias

MET = Interventions other than cognitive behavioural programmes avoided or controlled for across comparison groups.

UNCLEAR = Use of interventions other than cognitive behavioural programmes not reported and cannot be verified by contacting the investigators.

NOT MET = Dissimilar use of interventions other than cognitive behavioural programmes across comparison groups, i.e. differences in the care provided to the participants in the comparison groups other than the intervention under investigation.

Prevention of detection bias

MET = Assessor unaware of the assigned treatment when collecting outcome measures

UNCLEAR = "Blinding" of assessor not reported and cannot be verified by contacting investigators.

NOT MET = Assessor aware of the assigned treatment when collecting outcome measures.

Prevention of attrition bias

MET = Losses to follow up less than 20% and equally distributed (as judged by two reviewers) between comparison groups (e.g. 18% and 16%).

UNCLEAR = Losses to follow up not reported.

NOT MET = Losses to follow up 20% or greater, or not equally distributed (as judged by two reviewers) between comparison groups.

Intention-to-treat

MET = Intention to treat analysis performed or possible with data provided.

UNCLEAR = Intention to treat not reported, and cannot be verified by contacting the investigators.

NOT MET = Intention to treat analyses not done and not possible for reviewers to calculate independently.

Data analysis and presentation

We expressed binary outcome measures (for example, violent/not violent) as risk ratios (relative risks). Continuous measures were calculated as mean differences or (when different scales were used) standardised mean differences. We reported the 95% confidence intervals for all of the above.

If the primary studies were sufficiently homogenous, we performed a fixed-effect meta-analysis. Homogeneity was tested with the Q-test (Chi-square, P-value) and we measured degree of heterogeneity with I^2 (I-squared, Higgins 2002). But a decision did not rest solely on the outcome of these procedures. In addition to the formal procedures, we also took into account common sense, the nature of the measures, etc. If there was statistically significant heterogeneity among studies' effect sizes, a random-effects model was used. Effect sizes were pooled across studies using the DerSimonian and Laird method for random-effect models. For fixed-effect models we used the Mantel-Haenszel method for dichotomous data (except for data analysed using Peto's method) or the inverse variance method for continuous or generic inverse variance data. We also checked forest plots for detecting heterogeneity.

Cluster-randomised trials

No such trials were identified in this version of the review. See [Table 10](#) for plans for future updates.

Dealing with dependent outcomes

In some primary studies, several different outcomes are measured on the same participants. Sometimes the same outcome is measured at multiple points in time. As these data are from the same sample of participants, and, therefore, are not independent estimates of treatment effect, we aimed to analyse the data in such a way that any analysis contained a single, most recent, outcome (one measure from a single point in time) from each sample. The rationale for using the most recent outcome was that we opted for the longest possible follow-up time. In cases with several treatment arms, we compared only one of the treatment arms with the control group. The decisions and rationale for this are reported separately for each study in the Results section.

Heterogeneity

When there was statistically significant heterogeneity among primary outcome studies, the following factors were considered as possible explanations: voluntary or mandatory participation, intensity or length/period of the intervention, and differences in participant characteristics. If the primary studies were too heterogeneous to be grouped according to these characteristics, a meta-analysis was not performed. If there were many primary studies, we classified them according to these variables in order to identify possible sources of heterogeneity. We considered performing moderator analyses (stratification on subgroups, meta-analysis analogue to ANOVA, meta-regression) to explore how observed variables were related to heterogeneity.

Assessment of publication bias

We used a funnel plot ([Fig 01 - Funnel plot](#)) to explore the likelihood of publication bias. Asymmetry of the funnel plot may indicate possible publication bias in this review, but may also indicate other methodological or sample size issues within the trials. If asymmetry of the funnel plot was found, the clinical diversity of the studies was examined ([Egger 1997](#)).

Sensitivity analyses

Sensitivity analysis was inappropriate for this version of the review. See [Table 10](#) for plans for future updates.

Description of studies

Results of the search. From the initial search, 1969 records were identified, of which 1724 records were excluded as clearly irrelevant. Of the remaining 245 possibly relevant records, 75 were identified for further examination.

Included studies. After reading the full text reports, 12 records were included. They reported results from six randomised trials conducted in the USA. None of the studies were cluster-randomised. There were two types of comparisons; four studies compared CBT with a non-intervention control, while two studies compared CBT with another active treatment (See [Fig 02 - Flowchart](#)).

CBT versus Control.

In the [Bronx Exp. 2005 \(Bronx Exp. 2005\)](#), 420 convicted male domestic violence offenders were randomised into four experimental conditions:(1) batterer programme plus monthly judicial monitoring, (2) batterer programme plus graduated monitoring, (3) monthly monitoring only, and (4) graduated monitoring only. The model assumes that battering is rooted in societal norms that support male abuse of women. In order to measure the effect of the batterer programme we chose to combine the two groups receiving the batterer programme with the two other groups not receiving the batterer programme. The men were tracked for 12 months after court-ordering, and for up to eighteen months for most of them. The programme was in a group format and lasted for 26 weeks.

In the [Brooklyn Exp. 2000 \(Brooklyn Exp. 2000\)](#), 376 court-mandated offenders were randomly assigned to batterer treatment or to a treatment irrelevant to the violence problem (40 hours of community service). The length of the batterer treatment was 39 hours, but some men were assigned to complete the treatment in 26 weeks and others in 8 weeks. Interviews with victims and offenders were attempted at six and twelve months after the court-order date. Records of criminal justice agencies were also checked to determine if new crime reports or arrests had occurred involving the same defendant and victim.

A total of 404 men convicted of misdemeanour in Broward County, Florida ([Broward Exp. 2000](#)) were randomly assigned to a batterer programme using the Duluth Model or to a control group involving one year probation. The Duluth Model includes a feminist, cognitive psychoeducational curriculum provided in a group session. Its intent is to help domestic violence offenders develop an understanding of how battering is part of a range of male behaviours that seek to control women. The researchers hypothesised that men with a high stake in conformity would have a lower likelihood of recidivating than those with a low stake in conformity. The offenders were interviewed at time of adjudication and six and twelve months post-adjudication. Probation records and computer checks with the local police for all new arrests were used to track the defendants for one-year post-adjudication.

The largest study in this review was conducted in the military ([San Diego Navy 2000](#)). Participants were servicemen in the navy who had been substantiated as having physically assaulted their wives. The 861 couples in the study were randomly assigned to four groups: a men's group (N = 218), a conjoint group (the men participating with their wives) (N = 216), a rigorously monitored group (N = 213), and a control group (N = 214). The men's group met weekly for six months and then monthly for another six months and included both didactic and process activities. In the didactic part of the sessions, group leaders addressed perpetrator attitudes and values regarding women and violence toward women and taught the men a variety of skills thought to be important to the successful elimination of the continued abuse of women (e. g. cognitive restructuring, empathy enhancement, communication skills, anger modification, and jealousy). The process part of the sessions involved dealing with issue raised in the didactic segments of the session as well as with other issues that emerged. The conjoint group was similar to the men's group except for the presence of wives. The rigorously monitored group was formed based on a "stake in conformity" strategy to determine if male perpetrators held accountable for their abusiveness toward their wives, using systematized and official monitoring procedures, would stop the continued abuse. The interventions were cognitive-behavioural and outcome data

were collected both from the male servicemen and their female partners at roughly six-month intervals over the approximate 18-month experimental period. The interventions lasted for 12 months. We chose to compare the men's group with the control group because we thought that this comparison would be the most similar to the other included studies.

CBT versus Other therapy

In one study ([Wisconsin Study 1996](#)), 218 men were randomly assigned to receive either feminist-cognitive-behavioural group treatment (FCBT) or process-psychodynamic group treatment (PPT). FCBT focused on progressive relaxation, using coping thoughts, and becoming aware of feelings. PPT focused on childhood losses and rejections, childhood experience with violence and emotional safety in the group. FCBT also used leader role-play, lectures and giving advice, while PPT emphasised self-disclosure. Treatment integrity was verified through audio-taped recordings of each session. The partners of 79 % of the 136 treatment completers gave reports of the men's behaviour an average of two years post-treatment. There were also arrest records and self-reports by the men.

One study ([Yale Study 2007](#)), which was conducted in the state of Connecticut, randomly assigned substance dependent offenders with reported intimate partner violence to one of the following interventions: A 12-week substance abuse & domestic violence group (grounded in CBT) (N = 32) or a 12-week twelve-step facilitation group (N = 32). Data were collected using the SCID (Structured Clinical Interview for DSM-IV-TR), Addiction Severity Index, Substance Abuse Calendar, Conflict Tactics Scale Revised, breath samples, urine toxicology screens, and collateral reports from female partners at baseline, monthly, and post-treatment periods.

Methodological quality of included studies

All included trials were reported as randomised and the method of randomisation was adequate in four out of the six studies. Concealment of allocation was judged as adequate in three studies ([Bronx Exp. 2005](#); [San Diego Navy 2000](#), [Yale Study 2007](#)). In four studies it was clear that there were no systematic differences in treatment between groups aside from the experimental intervention (low risk of performance bias). Detection bias was unclear in all studies except for [Yale Study 2007](#), that is, it was not clear whether the persons collecting the results were blind to treatment group allocation. It should be noted, however, that this may not be important whenever the outcomes are written criminal justice records. There were high or unclear attrition in five out of six studies. Only the Navy ([San Diego Navy 2000](#)) managed to keep most of the participants in the intervention. Intention-to-treat was met in all studies. The randomisation was frequently compromised by judges overriding the allocation because they did not want certain of the defendants to be in the control group. In sum, all studies had a high risk of bias except for the [San Diego Navy 2000](#) Experiment ([San Diego Navy 2000](#)) which we judged to have a moderate risk of bias.

Results

CBT versus Control

Proportion new violence

The relative risk of 1.96 in the [Bronx Exp. 2005](#) ([Bronx Exp. 2005](#)) indicated that CBT had a harmful effect on new violence, but the 95% confidence interval from 0.96 to 3.99 was wide and not statistically significant. The CBT in the [Brooklyn Exp. 2000](#) ([Brooklyn Exp. 2000](#)) had a clearly positive effect on new violence (relative risk: 0.39, 95% CI 0.23 to 0.67). This means that the intervention on average reduced violence by 61% relative to the control group. The relative risk in the [Broward Exp. 2000](#) ([Broward Exp. 2000](#)) was 1.01 (no effect) with 95% confidence interval from 0.71 to 1.42. The results of the [San Diego Navy 2000](#) Study ([San Diego Navy 2000](#)) showed a positive effect of the intervention compared to the control group (relative risk 0.82), but

the effect was not statistically significant (95% CI 0.63 to 1.09).

In summary, four of the six included studies compared CBT with a control group looking at new violence. The effects were heterogeneous, with large effects in both directions. Only one study ([Brooklyn Exp. 2000](#)) showed a statistically significant effect, and it was positive. A meta-analysis involving 1771 men showed a risk ratio of 0.86, but the 95% confidence interval included zero difference (from 0.54 to 1.38).

CBT versus Other therapy

[Wisconsin Study 1996](#) (FCBT versus PPT)

Proportion new violence

The risk ratio (1.07) of the [Wisconsin Study 1996](#) ([Wisconsin Study 1996](#)) was not statistically significant (95 percent confidence interval from 0.68 to 1.68).

[Yale Study 2007](#) (SADV versus twelve step facilitation)

Frequency of new violence

In the Yale study ([Yale Study 2007](#)), the mean number of reported violent episodes per month was higher at post-treatment in the SADV group (0.95) than in the TSF group (0.73). This corresponds to a standardised mean difference of 0.30 (95% CI: -0.22 to 0.81).

Presence of new violence

There were reports of violence for three of the men in the SADV group and for two men in the TSF group. This corresponds to a risk ratio of 1.50 (95% CI: 0.27 to 8.32).

Discussion

There have been few randomised evaluations of cognitive behavioural therapy for men who physically abuse their female partner. All such studies have so far been conducted in the USA. The studies also have relatively small sample sizes, the largest study having 861 participants. This review has employed an extensive search strategy with no restrictions on publication language or geographical region. Because there were so few studies, we could not model sources of heterogeneity using e. g. meta-regression or stratified analyses. Only one meta-analysis was performed. The results were inconsistent and heterogeneous, but there was no clear evidence for publication bias (Figure 01). The number of studies in this funnel plot (N = 4) is, however, so small that great caution should be taken in its interpretation. The methodological quality of the included studies was generally low ([Table 01](#)). The concealment of the random allocation sequence was mostly unclear or not met. Furthermore, it was mostly unclear whether the assessors of the results were aware of group allocation. Finally, the attrition was either unclear or high. Unclear concealment of allocation and high attrition both contribute to high risk of selection bias, but blinding of the assessors is probably not of serious concern because criminal justice records are in written form.

Two anonymous peer-reviewers pointed to a number of issues pertaining to contemporary US debates which should be mentioned here. The partners of the male perpetrators were offered some support. This support might have made the partners more able to report new violence, independently of the effect of the interventions on the men. This review focuses exclusively on effects on physical violence. But even where interventions may have an impact on physical violence, perpetrators may continue or possibly increase the use of emotionally controlling behaviours ([Dobash 2000](#)). Reliance on criminal justice records is also problematic because much violence is never recorded by the police. Some authors have been worried about whether assignment to a batterer intervention programme actually sends an implicit message that

contradicts its official aim, namely that battering will not be taken seriously. This raises another debate: although some form of monitoring for compliance was present in all of the studies reviewed, sanctions for noncompliance are very rare. Moreover, the issue of attendance versus drop-out raises the important issue of selection bias, namely whether men who complete programmes are more motivated to stop their violence than those who never attend or drop out. While randomised controls should eliminate selection bias, the poor ways in which compliance is monitored suggest this may not be so. This review focuses on behaviour change, but many programmes actually view behavioural change as unlikely. They focus more on message sending and re-education or education.

CBT is a broad category that encompasses many different approaches. The most widely used model in the US, the Duluth Model, is included in this review because it has components of CBT. Not to include this model would eliminate most batterer intervention programmes in the US. But the fact that it has such components does not mean that everyone recognises it as a form of CBT. The Duluth Model includes strong elements of anti-sexist messaging, for instance, whereas traditional CBT approaches do not.

Care should be taken when generalising the results of this review to other parts of the world. First, there may be different baseline risk of violence across populations. Second, the motivation to comply with the treatment might vary greatly across populations of violent men. Men have different reasons for being in therapy. Their wives might have threatened to leave them if they do not enter therapy, or the men might have been court-ordered to take part. The men differ in their degree of stake in conformity, and such variables (employment and age) have predicted both attendance at treatment and re-offending ([Broward Exp. 2000](#)).

Reviewers' conclusions

Implications for practice

The research evidence is insufficient to draw conclusions about the effectiveness of cognitive behavioural interventions for spouse abusers in reducing or eliminating male violence against female partners. Note that this does not mean that there is evidence for no effect. We simply do not know whether the interventions help, whether they have no effect, or whether they are harmful.

Implications for research

In the USA, and in other countries, there is a need for more, and larger-scale, randomised interventions. This has been possible in other fields, such as welfare-to-work ([Smedslund 2006](#)), where the weight of evidence involves randomised evaluations with more than 400,000 participants. Each primary study has randomised several thousands of participants. In the rest of the world there has been a complete lack of randomised interventions. Such interventions are highly needed in order to estimate the effects of CBT on domestic violence, even though this might not be as easy to accomplish with domestic violence as it was with welfare-to-work.

Characteristics of included studies

Study ID	Methods	Participants	Interventions	Outcomes	Notes	Allocation concealment
Bronx Exp. 2005	Randomised controlled trial. 2x2 factorial design. Register data. Interviews.	420 offenders arraigned on a domestic violence misdemeanour, convicted of a violation, and sentenced to conditional discharge with a one-year protection order in favor of the victim. The mean age was 30.8 years, and there were 40% Blacks, 42% Hispanics, and 18% White or other ethnic group.	Four different interventions; (1) batterer program + monthly monitoring, (2) batterer program + graduated monitoring, (3) only monthly monitoring, (4) only graduated monitoring. The batterer program lasted for 26 weeks with classes meeting weekly for 75 minutes.	Official re-arrests, victim reports of re-abuse, victim satisfaction		A
Brooklyn Exp. 2000	Randomised controlled trial. Interviews with batterers and victims at 6 and 12 months after the sentence date. In addition, records of criminal justice agencies were checked for new crime reports or arrests.	376 court-mandated batterers.	39 hours of class time. Some were assigned to complete the treatment in 26 weeks and others in eight weeks. Men assigned to the control condition were sentenced to 40 hours of community service. The intervention included defining domestic violence, understanding historical and cultural aspects of domestic abuse and reviewing criminal/legal issues. Batterers were encouraged to take responsibility for their anger, actions, and reactions.	Proportion violent		C
Broward Exp. 2000	Randomised controlled trial. Outcomes were collected by means of interviews with	Men (N=404) convicted of misdemeanour domestic violence in Broward County	Duluth Model, which is a feminist, cognitive psychoeducational	Differences between the groups at time of adjudication (Time 1), at least 6-months	Predicted that having a stake in conformity would predict when an intervention will be	C

	batterers and victims, and police records of repeat violence.	during a 5-month period in 1997.	curriculum provided in 26-week group sessions. Men in the control group were sentenced to 1 year's probation.	postadjudication (Time 2), and changes between Time 1 and Time 2 were examined. Offender attitudes, beliefs, and self-reported behaviours were collected from the offenders. The men answered the revised Conflict Tactics Scale.	effective in reducing the likelihood of subsequent violence. There was controversy in the community around randomizing men into a spouse abatement programme. This led to low victim response rates, high staff turnover, delays, and other problems.	
San Diego Navy 2000	Randomised controlled trial. Sample selection took 46 months. A computer did the randomisation to one of four groups: a men's group, a conjoint group, a rigorous assessment group, or a control group.	861 married U.S. Navy couples in which active-duty husbands were substantiated as having physically assaulted their wives. Mean age was 27 years. Mean length of marriage was 47 months. 83% had a mean of 1.7 children. Mean number of school years was 12.6.	The men's group, which used a cognitive-behavioural model of change, met weekly for 6 months and then monthly for another 6 months, for a total of 1-year treatment period. The conjoint group also had 26 weekly sessions that included both didactic and process activities followed by six monthly sessions. The main difference from the men's group was the presence of wives. The rigorous monitoring group attempted to hold perpetrators accountable for their abusiveness. Every six weeks a record search was completed to determine if perpetrators had been arrested or referred to court anywhere in San Diego County. Wives	Four types of outcome measures were used. A self-reported episodic measure assessed the number of incidents or episodes in which a victim or perpetrator reported being abused across three different levels of abuse. The second outcome measure, the Modified Conflict Tactics Scale focused on types of abusive behaviours as reported by respondents. The third outcome measure consisted of official police and court records for all respondents (both victims and perpetrators) living within the boundaries of San Diego County. The fourth outcome measure focused on the date of the first instance in which a repeat case of spouse		A

			were called monthly and asked about new instances of abuse. Men assigned to the control group did not receive any treatment but their wives received preliminary assistance called 'stabilization and safety planning'.	assault occurred as indicated by both official arrest records and victim reports of new physical injuries.	
Wisconsin Study 1996	Randomised controlled trial. Treatment integrity was verified through audio-taped codings of each session.	218 men who had been referred by the criminal justice system and accepted for treatment at a family counselling agency that was certified as an outpatient mental health clinic.	Feminist-cognitive behavioural (FCBT) or process-psychodynamic (PPT) group treatments. The FCBT condition followed a highly structured format. Agendas and homework assignments were included in each session. Each session included a didactic session on communication and cognitive skills, relaxation/desensitization training, consciousness raising about sex roles and violence against women, and behavioural or cognitive rehearsal. The PPT did not use agendas but focused on building trust and a sense of safety, uncovering childhood traumas and reconnecting with traumatic childhood events.	The partners of 79% of the 136 treatment completers gave reports of the men's behaviour an average of 2 years post-treatment. These reports were supplemented with arrest records and self-reports.	Hypothesised that the feminist-cognitive-behavioural model is best suited to men with antisocial traits and that the process-psychodynamic model is most suited for men with moderate to high level of dependency needs because they are much more likely to engage in group process and methods for enhancing self-awareness.
Yale Study 2007	Randomised controlled trial	Substance dependent offenders (N = 64) with a	A 12 week Substance Abuse & Domestic Violence group	Data were collected using the SCID, Addiction Severity	B

history of intimate partner violence. Mean age was 38 and they had the following racial composition: 49% Caucasian, 33% African American, 10% Hispanics, and 8% other.

(grounded in CBT (N = 32) or a 12 week Twelve Step Facilitation group (N=32).

Index, Substance Use Calendar, Conflict Tactics Scale Revised, breath samples, urine toxicology screens, and collateral reports from female partners at baseline.

Characteristics of excluded studies

Study ID	Reason for exclusion
Bern 1984	Not an intervention study
Currie 1983	Not RCT. Not CBT.
Currie 1985	Not RCT. Not CBT.
Deschner 1986	Not RCT
Dowden 1999	Not RCT
Dutton 1986	Not RCT
DVFCT Program 2004	Subjects were randomised to individual couple therapy or to multi-couple group therapy. Random assignment was not applied in creating the comparison group.
Easton 2006	Not an intervention study.
Echeburúa 1997	Not RCT
Eckhardt 2004	Not RCT
Edleson 1990	Not CBT
Edleson 1991	Review. Not CBT.
Elliott 2003	Not RCT
Fals-Stewart 2001	Wrong participants
Fals-Stewart 2002	Not RCT
Fals-Stewart 2005	Not an intervention study
Flournoy 1992	The study was not randomized.
Goldkamp 1996	Not CBT
Gondolf 1985	Review
Gondolf 2000	Not RCT
Gondolf 2001	Not RCT
Gondolf 2004	Not RCT
Gondolf 2005	Not RCT
Gray 2004	Not RCT. Not CBT
Hamberger 1989	Not RCT

Hanson 2003	Not RCT
Harrell 1991	Not RCT
Harris 1981	We were not able to obtain the report in full text.
Hendricks 2006	Not a randomized trial
Howard County 2003	The study was not randomized.
Johnston 1985	Not RCT
Jones 2001	Not an intervention study
Jones 2002	Not CBT
Kriner 1988	Not RCT
Lanza 2002	Wrong participants
Lawson 2006	Not a randomized trial
Leonard 2005	Editorial
Loza 1999	Wrong participants. Not RCT. Probably not CBT
McAllister 2003	Not a randomized trial
Moore 1994	Not RCT
Morrel 2000	Not RCT
Morrel 2003	Not RCT
New Zealand 2005	Not a randomized trial
Nosko 1988	Not RCT
O'Farrell 1995	Not RCT
O'Farrell 2004	Not RCT
Palmer 1992	Not CBT
Pressman 1983	Not RCT
Rosenbaum 1997	Not RCT. Probably not CBT
Satel 2001	Review with wrong participants
Schuenger 1988	Not RCT and not CBT
Sherman 1984	Not CBT
Sherman 1991	Not CBT
Shupe 1987	Review

South Florida 2002	We could only obtain an abstract.
Stacey 1984	Not RCT
Sullivan 1990	Not CBT. Wrong participants
Taft 2004	Not RCT
Taylor 1984	Not RCT. Not CBT
Tennant 1998	Wrong participants. Not RCT
Upshaw 2005	The outcomes were symptomatic distress, interpersonal relations, social role adjustments, and well-being, not whether the men ceased to beat their partners.
Wade 1985	Not RCT
Watt 1999	Wrong participants
Weidman 1986	Not an intervention study
Williams 1995	Not RCT. Probably not CBT

References to studies

Included studies

Bronx Exp. 2005

{ published data only }

Labriola M, Rempel M, Davis RC. Testing the Effectiveness of Batterer Programs and Judicial Monitoring: Results from a Randomized Trial at the Bronx Misdemeanor Domestic Violence Court. New York, NY: Center for Court Innovation, 2005.

Brooklyn Exp. 2000

{ published data only }

Davis RC, Taylor BG, Maxwell CD. Does Batterer Treatment Reduce Violence? A Randomized Experiment in Brooklyn -- Executive Summary Included (available online: www.ncjrs.gov/pdffiles1/nij/grants/180772.pdf , accessed January 2006). Washington, DC: US Dept. of Justice, 2000.

Maxwell CD, Davis RC, Taylor BG. Results From the Brooklyn Domestic Violence Treatment Experiment (NCJ 199728) (available online: www.ncjrs.gov/pdffiles1/nij/199728, accessed January 2006). Washington DC: National Institute of Justice, US Dept. of Justice, 2004.

* Taylor BG, Davis RC, Maxwell CD. The effects of a group batterer treatment program: a randomized experiment in Brooklyn. *Justice Quarterly* 2001;18(1):171-201.

Broward Exp. 2000

{ published data only }

Feder L, Forde DR. Test of the efficacy of court-mandated counseling for domestic violence offenders: The Broward Experiment (available online: www.ncjrs.gov/pdffiles1/nij/grants/184631.pdf : accessed January 2006). Washington DC: US Dept. of Justice, 2000.

* Feder L, Dugan L. A test of the efficacy of court-mandated counseling for domestic violence offenders: The Broward Experiment. *Justice Quarterly* 2002;19(2):343-75.

San Diego Navy 2000

{ published data only }

* Dunford FW. The San Diego Navy experiment: an assessment of interventions for men who assault their wives. *Journal of Consulting & Clinical Psychology* 2000;68(3):468-76.

Dunford, FW. Determining program success: the importance of employing experimental research designs. *Crime and Delinquency* 2000;46(3):425-34.

Wisconsin Study 1996

{ published data only }

* Saunders DG. Feminist-cognitive-behavioral and process-psychodynamic treatments for men who batter: interaction of abuser traits and treatment models. *Violence & Victims* 1996;11(4):393-414.

Yale Study 2007

{ unpublished data only }

Easton CJ, Mandel D, Babuscio T, Rounsaville BJ, Carroll KM. Differences in treatment outcome between male alcohol dependent offenders of domestic violence with and without positive drug screens. *Addictive Behaviors* 2007;Feb 4; [Epub ahead of print]:doi: 10.1016/j.addict.2007.01.031.

* Easton CJ, Mandel DL, Hunkele KA, Nich C, Rounsaville BJ, Carroll KM. A cognitive behavioral therapy for alcohol-dependent domestic violence offenders: an integrated substance abuse-domestic violence treatment approach (SADV). *American Journal on Addictions* 2007;16(1):24-31.

Easton CJ. Treatment outcome among substance-dependent offenders of intimate partner violence: A randomized trial. CPDD-2005 Orlando Florida 2005.

Excluded studies

Bern 1984 {published data only}

Bern EH, Bern LL. A group program for men who commit violence towards their wives. *Social Work with Groups* 1984;7(1):63-77.

Currie 1983 {published data only}

Currie DW. A Toronto model. *Social Work with Groups* 1983;6(3-4):179-88.

Currie 1985 {published data only}

Currie D. Group model for men who assault their partners. In: D. Sinclair, editor(s). *Understanding Wife Assault: A Training Manual for Counsellors and Advocates*. Toronto: Ontario Government Bookstore, 1985:120-43.

Deschner 1986 {published data only}

Deschner J. *The Hitting Habit: Anger Control for Battering Couples*. New York: Free Press, 1986.

Dowden 1999 {published data only}

Dowden C, Blanchette K, Serin R. Anger management programming for federal male inmates: an effective intervention. Ottawa: Research Branch, Correctional Service Canada, 1999.

Dutton 1986 {published data only}

Dutton DG. The outcome of court-mandated treatment for wife-assault: a quasi-experimental evaluation. *Violence and Victims* 1986;1(3):163-175.

DVFCT Program 2004 {published data only}

Stith SM, Rosen KH; McCollum EE, Thomsen CJ. Treating intimate partner violence within intact couple relationships: Outcomes of multi-couple versus individual couple therapy. *Journal of Marital and Family Therapy* 2004;30(3):305-18.

Easton 2006 {published data only}

Easton CJ. The role of substance abuse in intimate partner violence. *Psychiatric Times* 2006;23:25-7.

Echeburúa 1997 {published data only}

Echeburúa E, Fernández-Montalvo J. Tratamiento cognitivo-conductual de hombres violentos en el hogar: Un estudio piloto [Cognitive-behavioral treatment of abusive men: a pilot study [Spanish]]. *Análisis y Modificación de Conducta* 1997;23(89):355-384.

Eckhardt 2004 {published data only}

Eckhardt CI. Partner assaultive men and the stages and processes of change. *Journal of Family Violence* 2004;19(2):81-93.

Edleson 1990 {published data only}

Edleson JL, Syers M. Relative effectiveness of group treatments for men who batter. *Social Work Research and Abstracts* 1990;26(2):10-8.

Edleson 1991 {published data only}

Edleson JL, Syers M. The effects of group treatment for men who batter: An 18-month follow-up study. *Research on Social Work Practice* 1991;1(3):227-43.

Elliott 2003 {published data only}

Elliott JD. Responsibility acceptance and aggression changes in group therapy for relationship abuse perpetrators (Phd). Baltimore, MD: University of Maryland, 2002.

Fals-Stewart 2001 {published data only}

Fals-Stewart W, O'Farrell TJ, Birchler GR. Behavioral couples therapy for male methadone maintenance patients: Effects on drug-using behavior and relationship adjustment. *Behavior Therapy* 2001;32(2):391-411.

Fals-Stewart 2002 {published data only}

Fals-Stewart W, Kashdan TB, O'Farrell TJ, Birchler GR. Behavioral couples therapy for drug-abusing patients: effects on partner violence. *Journal of Substance Abuse Treatment* 2002;22(2):87-96.

Fals-Stewart 2005 {published data only}

Fals-Stewart W, Kennedy C. Addressing intimate partner violence in substance-abuse treatment. *Journal of Substance Abuse Treatment* 2005;29:5-17.

Flournoy 1992 {published data only}

Flournoy PS. A comparison of groups for men who batter (Phd). Pullman WA: Washington State University, 1992.

Goldkamp 1996 {published data only}

Goldkamp JS, Weiland D, Collins M, White M. The role of drug and alcohol abuse in domestic violence and its treatment: Dade County's domestic violence court experiment. Final report. Philadelphia, PA: Crime and Justice Research Institute, 1996.

Gondolf 1985 {published data only}

Gondolf, E. Men who batter. An integrated approach for stopping wife abuse. Holmes Beach, FL: Learning Publications, 1985.

Gondolf 2000 {published data only}

Gondolf EW. A 30-month follow-up of court-referred batterers in four cities. *International Journal of Offender Therapy and Comparative Criminology* 2000;44(1):111-28.

Gondolf 2001 {published data only}

Gondolf E, Snow JA. The program effect of batterer programs in three cities. *Violence & Victims* 2001;16(6):693-704.

Gondolf 2004 {published data only}

Gondolf EW. Regional and cultural utility of conventional batterer counseling. *Violence against Women* 2004;10(8):880-900.

Gondolf 2005 {unpublished data only}

Gondolf E. Culturally-focused batterer counseling for African-American men. Final report. Washington, DC: National Institute of Justice, 2005.

Gray 2004 {published data only}

Gray YL. e-Health for all: designing nurses agenda for the future. In: 8th International Conference in Nursing informatics. Knowledge Representation: mapping violence and spousal abuse using the international Classification for Nursing Practice. E-papers Servicos Editoriais Ltda. Rio de Janeiro, Brazil, 2003.

Hamberger 1989 {published data only}

Hamberger LK, Hastings JE. Counseling male spouse abusers: characteristics of treatment completers and dropouts. *Violence & Victims* 1989;4(4):275-86.

Hanson 2003 {published data only}

Hanson RK, Wallace-Capretta S. A multi-site study of treatment for abusive men. Public Works and Government Services Canada 2003.

Harrell 1991 {published data only}

Harrell A. Evaluation of court-ordered treatment for domestic violence offenders: final report.

Washington DC: Urban Institute, Washington DC, 1991.

Harris 1981 {published data only}

Harris S, Sinclair D. Domestic violence project. A comprehensive model for intervention into the issue of domestic violence. Toronto, CA: Family Services Association of Metropolitan Toronto, 1981.

Hendricks 2006 {published data only}

Hendricks B, Werner T, Shipway L, Turinetti GJ. Recidivism among spousal abusers. Predictions and program evaluation. *Journal of Interpersonal Violence* 2006;21(6 (June)):703-16.

Howard County 2003 {published data only}

Elliott JD. Responsibility acceptance and aggression changes in group therapy for relationship abuse perpetrators. Baltimore MD: University of Maryland, 2002.

Morrel TM. Changes in self-efficacy, self-esteem and aggression in male batterers: A comparison of cognitive-behavioral and supportive group therapies (dissertation). University of Maryland: College Park, MD, 2000.

* Morrel TM, Elliott JD, Murphy CM, Taft CT. Cognitive behavioral and supportive group treatments for partner-violent men. *Behavior Therapy* 2003;34(1):77-95.

Johnston 1985 {published data only}

Johnston JD. A social learning and cognitive-behavioural analysis of aggressive and violent behaviour (PhD). Toronto: University of Toronto, 1985.

Jones 2001 {published data only}

Jones NG. A study of the influence of protective factors as a resource to traditional batterers' interventions in a population of African American active-duty male batterers (DSW). Norfolk: Norfolk State University, 2001.

Jones 2002 {published data only}

Jones NG. A study of the influence of protective factors as a resource to African American males in traditional batterers' interventions. *Journal of Health & Social Policy* 2002;16(1-2):169-183.

Kriner 1988 {published data only}

Kriner L, Waldron B. Group counseling: A treatment modality for batterers. *Journal for Specialists in Group Work* 1988;13(3):110-16.

Lanza 2002 {published data only}

Lanza ML, Anderson J, Boisvert CM, LeBlanc A, Fardy M, Steel B. Assaultive behavior intervention in the Veterans Administration: psychodynamic group psychotherapy compared to

cognitive behavior therapy. *Perspectives in Psychiatric Care* 2002;38(3):89-97.

Lawson 2006 {published data only}

Lawson DM, Barnes AD, Madkins JP, Francois-Lamonte BM. Changes in male partner abuser attachment styles in group treatment. *Psychotherapy: Theory, Research, Practice, Training* 2006;43(2):232-37.

Leonard 2005 {published data only}

Leonard KE. Alcohol and intimate partner violence: when can we say that heavy drinking is a contributing cause of violence? *Addiction* 2005;100:422-25.

Loza 1999 {published data only}

Loza W, Loza FA. The fallacy of reducing rape and violent recidivism by treating anger. *International Journal of Offender Therapy and Comparative Criminology* 1999;43(4):492-502.

McAllister 2003 {unpublished data only}

McAllister J. Intervention with batterers: Reexamining the significance of gender-role beliefs. San Francisco, CA: Saybrook Graduate School and Research Center, 2003.

Moore 1994 {published data only}

Moore KJ. The outcome evaluation of a cognitive behavior treatment program for court-mandated batterers residing in Broward County, Florida. Miami, FL: Florida International University, 1994.

Morrel 2000 {published data only}

Morrel TM. Changes in self-efficacy, self-esteem and aggression in male batterers: A comparison of cognitive-behavioral and supportive group therapies (dissertation). College Park, MD: University of Maryland, 2000.

Morrel 2003 {published data only}

Morrel TM, Elliot JD, Murphy CM, Taft CT. Cognitive behavioral and supportive group treatments for partner-violent men. *Behavior Therapy* 2003;34(1):77-95.

New Zealand 2005 {published data only}

* Polaschek DLL, Wilson NJ, Townsend MR, Daly LR. Cognitive-behavioral rehabilitation for high-risk violent offenders. An outcome evaluation of the violence prevention unit. *Journal of Interpersonal Violence* 2005;20(12):1611-27.

Nosko 1988 {published data only}

Nosko A, Wallace B. Group with abusive men: A multi-dimensional model. *Social Work with Groups* 1988;11(3):33-52.

O'Farrell 1995 {published data only}

O'Farrell TJ, Murphy CM. Marital violence before and after alcoholism treatment. 1995 *Journal of Consulting & Clinical Psychology*;63(2):256-62.

O'Farrell 2004 {published data only}

O'Farrell TJ, Murphy CM, Stephan SH, Fals-Stewart W, Murphy M. Partner violence before and after couples-based alcoholism treatment for male alcoholic patients: the role of treatment involvement and abstinence. *Journal of Consulting & Clinical Psychology* 2004;72(2):202-17.

Palmer 1992 {published data only}

Palmer S, Brown R, Barrera M. Group treatment program for abusive husbands: Long-term evaluation. *American Journal of Orthopsychiatry* 1992;62(2):276-83.

Pressman 1983 {published data only}

Pressman BM. *Family violence: Origin and treatment*. San Francisco: Jossey-Bass, 1983.

Rosenbaum 1997 {published data only}

Rosenbaum A, Gearan P, Ondovic, C. Completion and recidivism among court- and self-referred batterers in a psychoeducational group treatment program: implications for intervention and public policy. *Journal of Aggression, Maltreatment & Trauma* 2001;5(2):199-220.

Satel 2001 {published data only}

Satel SL. Who needs trauma initiatives? *Psychiatric Services* 2001;52(6):815-815.

Schuerger 1988 {published data only}

Schuerger JM, Reigle N. Personality and biographic data that characterize men who abuse their wives. *Journal of Clinical Psychology* 1988;44(1):75-81.

Sherman 1984 {published data only}

Sherman LW, Berk RA. The deterrent effects of arrest for domestic assault. *American Sociological Review* 1984;49(2):261-72.

Sherman 1991 {unpublished data only}

Sherman LW, Schmidt JD, Gartin RR, Rogan D, Collins DJ, Bacich A, Cohn EG. Effects of on-scene arrest on subsequent domestic violence: A randomized field experiment. In: *Joint Conference on Evaluation of the Organization for Economic Cooperation and Development (OECD)*. Washington DC: U.S. Department of Education and U.S. Department of Health and Human Services, 1991.

Shupe 1987 {published data only}

Shupe A, Stacey WA, Hazlewood LR. *Violent men, violent couples: The dynamics of domestic violence*. Lanham, MD: Lexington Books, 1987.

South Florida 2002 {published data only}

Pilet GA. Efficacy of adult psychoeducation in an anger management portion of a domestic violence treatment program (PhD). Tampa FL: University of South Florida, 2002.

Stacey 1984 {published data only}

Stacey WA, Shupe A. Research Monograph No. 29. Arlington, TX: Center for Social Research, The University of Texas, 1984.

Sullivan 1990 {published data only}

Sullivan JP, Mosher DL. Acceptance of guided imagery of marital rape as a function of macho personality. *Violence & Victims* 1990;5(4):275-286.

Taft 2004 {published data only}

Taft CT, Murphy CM, Musser PH, Remington NA. Personality, interpersonal, and motivational predictors of the working alliance in group cognitive-behavioral therapy for partner violent men. *Journal of Consulting and Clinical Psychology* 2004;72(2):349-354.

Taylor 1984 {published data only}

Taylor J. Structured conjoint therapy for spouse abuse cases. *Social Casework* 1984;65:11-8.

Tennant 1998 {published data only}

Tennant A, Hughes G. Men talking about dysfunctional masculinity: an innovative approach to working with aggressive, personality disordered offender-patients. *Psychiatric Care* 1998;5(3):92-9.

Upshaw 2005 {unpublished data only}

Upshaw R. The efficacy of thought field therapy as an adjunct treatment modality for male domestic-violence perpetrators with domestic abuse in their family of origin. Minneapolis, MN: Walden University, 2005.

Wade 1985 {published data only}

Wade TF. A time-limited conjoint cognitive-behavioral treatment for abusive spouses (dissertation). Missoula, MT: University of Montana, 1985.

Watt 1999 {published data only}

Watt B, Howells K. Skills training for aggression control: evaluation of an anger management programme for violent offenders. *Legal and Criminological Psychology* 1999;4(2):285-300.

Weidman 1986 {published data only}

Weidman A. Family therapy with violent couples. *Social Casework* 1986;67:211-18.

Williams 1995

{published data only}

Williams OJ. Treatment for African American men who batter. CURA Reporter: Bulletin of the Center for Urban and Regional Affairs, University of Minnesota 1995;25(3):12-6.

Studies awaiting assessment

Rathus 2006

{published data only}

Rathus JH, Cavanaugh N, Passarelli V. Dialectical behavior therapy (DBT): A mindfulness-based treatment for intimate partner violence. In: Baer RA, editor(s). Clinician's guide to evidence base and applications. Burlington, MA, USA: Academic Press, 2006:424.

** indicates the primary reference for the study*

Other references

Additional references

Arias 2002

Arias I, Dankwort J, Douglas U, Dutton MA, Stein K. Violence against women: the state of batterer prevention programs. *The Journal of Law, Medicine, & Ethics* 2002;30(3):157-65.

Babcock 2004

Babcock JC, Green CE, Robie C. Does batterers' treatment Work? A meta-analytic review of domestic violence treatment outcome research. *Clinical Psychology Review* 2004;23:1023-1053.

Beck 1979

Beck AT, Rush AJ, Shaw BF, Emery G. *Cognitive therapy of depression*. New York: The Guilford Press, 1979.

Butler 2000

Butler AC, Beck JS. Cognitive therapy outcomes: a review of meta-analyses. *Journal of the Norwegian Psychological Association* 2000;37:1-9.

CDCP 2003

Center for Disease Control and Prevention (CDCP). Domestic and Intimate Partner Violence. <http://www.cdc.gov/communication/tips/domviol.htm> (accessed December 2004) 2003 (July 23).

Davis 1999

Davis RC, Taylor BG. Does batterer treatment reduce violence? A synthesis of the literature. *Women and Domestic Violence: An Interdisciplinary Approach* 1999;10:69-93.

Dobash 2000

Dobash RE, Dobash RP. Evaluating criminal justice interventions for domestic violence. *Crime & Delinquency* 2000;46:252-70.

Egger 1997

Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal* 1997;315:629-34.

Higgins 2002

Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* 2002;21:1539-58.

Higgins 2005

Higgins JPT, Green S. Cochrane Reviewers' Handbook 4.2.5 [updated May 2005]. In: The Cochrane Library. Chichester: UK: John Wiley & Sons, Ltd, 2005.

Pence 1993

Pence E, Paymar M. Education groups for men who batter: The Duluth model. New York: Springer, 1993.

Smedslund 2006

Smedslund G, Hagen KB, Steiro A, Johme T, Dalsbo TK, Rud MG. Work programmes for welfare recipients (<http://www.campbellcollaboration.org/frontend2.asp?ID=54>). Philadelphia, PA: Campbell Collaboration, 2006 (August).

SRS 2005

SRS. Version 3.0 [Computer program]. Ottawa, Ontario, Canada: Trialstat, 2005.

WHO 2002

World Health Organization. Intimate partner violence.
http://www.who.int/violence_injury_prevention/violence/global_campaign/en/ipvfacts.pdf
(accessed 22 December 2004) 2002.

Table of comparisons

01 CBT versus control

01 Proportion new violence

02 CBT versus other therapy

01 Proportion new violence

02 Frequency of violence

03 Any violence

Additional tables

01 Quality assessment

Study name	Allocation sequence	Allocation conceal	Performance bias	Detection bias	Attrition bias	Intention-to-treat
Bronx	Met	Met	Met	Unclear	Not met	Met
Brooklyn	Met	Not met	Unclear	Unclear	Not met	Met
Broward	Met	Not met	Met	Unclear	Unclear	Met
San Diego	Met	Met	Met	Unclear	Met	Met
Wisconsin	Unclear	Unclear	Met	Unclear	Unclear	Met
Yale	Not met	Met	Unclear	Met	Not met	Met

Additional tables

02 MEDLINE search strategy

MEDLINE

MEDLINE

Database: MEDLINE 1966 to September Week 3 2006

Date: 05.10.2006

Search by Sigrun Espelien Aasen

Number of hits: 26

Search strategy in OVID : CBT_Medline 041006

1.Battered Women/

2.domestic violence/ or spouse abuse/

3.((family or domestic or conjugal or partner) adj3 violence).tw.

4.((abus\$ or batter\$ or beat\$ or assault\$) adj3 (wom?n or partner\$ or spouse\$ or female\$ or wife or wives or domestic\$ or fiance or cohabitant\$ or live?in)).tw.

5.((male\$ or men or man or partner\$ or spouse\$ or husband or fiance or cohabitant\$ or live?in) adj3 (batter\$ or perpetrator\$ or abus\$ or violen\$ or beat\$ or assault)).tw.

6.or/1-5

7.behavior therapy/ or cognitive therapy/

8.psychotherapy, rational-emotive/

9.(cognitiv\$ adj3 (therap\$ or train\$ or techni\$ or question\$ or approach\$ or assessment\$)).tw.

10.(behavio?r\$ adj3 (therap\$ or train\$ or modif\$ or experiment\$)).tw.

11.(rational\$ adj3 emotive\$).tw.

12.cbt.tw.

13.(schemas or schematas).tw.

14."Imagery (Psychotherapy)"/

15.imager\$.tw.

16.((cognitive\$ or mental\$) adj3 (map\$ or model\$)).tw.

17.(socratic\$ adj3 (question\$ or method\$ or dialogue\$ or strateg\$ or sequence\$)).tw.

18.(dysfunctional adj2 (thought\$ or assumption\$ or rule\$ or appraisal\$ or belief\$ or attitude\$ or scheme\$)).tw.

19.(automatic adj3 (thought\$ or process\$)).tw.

20.(nat or nats).tw.

21.reattribution\$.tw.

22.((key or core) adj2 belief\$).tw.

23.or/7-22

24.6 and 23

Additional tables

03 EMBASE search strategy

EMBASE

EMBASE

Database: EMBASE 1980 to 2006 Week 39

Date: 05.10.2006

Search by Sigrun Espelien Aasen

Number of hits: 112

Search strategy for OVID : CBT_Embase 041006

- 1.exp domestic violence/ or battered woman/ or family violence/ or partner violence/
- 2.((family or domestic or conjugal or partner) adj3 violence).tw.
- 3.((abus\$ or batter\$ or beat\$ or assault\$) adj3 (wom?n or partner\$ or spouse\$ or female\$ or wife or wives or domestic\$ or fiance or cohabitant\$ or live?in)).tw.
- 4.((male\$ or men or man or partner\$ or spouse\$ or husband or fiance or cohabitant\$ or live?in) adj3 (batter\$ or perpetrator\$ or abus\$ or violen\$ or beat\$ or assault)).tw.
- 5.or/1-4
- 6.behavior therapy/ or cognitive therapy/
- 7.behavior modification/
- 8.(cognitiv\$ adj3 (therap\$ or train\$ or techni\$ or question\$ or approach\$ or assessment\$)).tw.
- 9.(behavio?r\$ adj3 (therap\$ or train\$ or modif\$ or experiment\$)).tw.
- 10.(rational\$ adj3 emotive\$).tw.
- 11.cbt.tw.
- 12.(schema\$ or schemata\$).tw.
- 13.imagery/
- 14.imager\$.tw.
- 15.((cognitiv\$ or mental\$) adj3 (map\$ or model)).tw.
- 16.(socratic\$ adj3 (question\$ or method\$ or dialogue\$ or strateg\$ or sequence\$)).tw.
- 17.(dysfunctional adj2 (thought\$ or assumption\$ or rule\$ or appraisal\$ or belief\$ or attitude\$ or scheme\$)).tw.
- 18.(automatic adj2 (thought\$ or process\$)).tw.
- 19.(nat or nats).tw.
- 20.reattribution\$.tw.

21.((key or core) adj2 belief\$.tw.

22.or/6-21

23.5 and 22

24.limit 23 to yr="2005 - 2006

Additional tables

04 CINAHL search strategy

CINAHL

CINAHL

Database: CINAHL - Cumulative Index to Nursing & Allied Health Literature
1982 to September Week 5 2006
Date: 05.10.2006
Search by Sigrun Espelien Aasen
Number of hits: 14

Search strategy in OVID: CBT_Cinah1 041006

- 1.Battered Women/
- 2.Domestic Violence/
- 3.partner abuse/ or spouse abuse/
- 4.((family or domestic or conjugal or partner) adj3 violence).tw.
- 5.((abus\$ or batter\$ or beat\$ or assault\$) adj3 (wom?n or partner\$ or spouse\$ or female\$ or wife or wives or domestic\$ or fiance or cohabitant\$ or live?in)).tw.
- 6.((male\$ or men or man or partner\$ or spouse\$ or husband or fiance or cohabitant\$ or live?in) adj3 (batter\$ or perpetrator\$ or abus\$ or violen\$ or beat\$ or assault)).tw.
- 7.or/1-6
- 8.behavior therapy/ or cognitive therapy/
- 9.Behavior Modification/
- 10.(cognitiv\$ adj3 (therap\$ or train\$ or techni\$ or question\$ or approach\$ or assessment\$)).tw.
- 11.(behavio?r\$ adj3 (therap\$ or train\$ or modif\$ or experiment\$)).tw.
- 12.(rational\$ adj3 emotive\$).tw.
- 13.cbt.tw.
- 14.(schema\$ or schemata\$).tw.
- 15.imagination/ or guided imagery/
- 16.imager\$.tw.
- 17.Concept Mapping/
- 18.((cognitiv\$ or mental\$) adj3 (map\$ or model)).tw.
- 19.(socratic\$ adj3 (question\$ or method\$ or dialogue\$ or strateg\$ or sequence\$)).tw.

20.(dysfunctional adj2 (thought\$ or assumption\$ or rule\$ or appraisal\$ or belief\$ or attitude\$ or scheme\$)).tw.

21.(automatic adj2 (thought\$ or process\$)).tw.

22.(nat or nats).tw.

23.reattribution\$.tw.

24.((key or core) adj2 belief\$).tw.

25.or/8-24

26.7 and 25

27.limit 26 to yr="2005 - 2006"

Additional tables

05 PsycINFO search strategy

PsycINFO

PsycINFO

Database: PsycINFO 1806 to October Week 1 2006

Date: 05.10.2006

Search by Sigrun Espelien Aasen

Number of hits: 55

Search strategy in OVID: "CBT_PsychInfo 041006"

- 1.partner abuse/
- 2.family violence/
- 3.((family or domestic or conjugal or partner) adj3 violence).tw.
- 4.battered females/
- 5.((abus\$ or batter\$ or beat\$ or assault\$) adj3 (wom?n or partner\$ or spouse\$ or female\$ or wife or wives or domestic\$ or fiance or cohabitant\$ or live?in)).tw.
- 6.((male\$ or men or man or partner\$ or spouse\$ or husband or fiance or cohabitant\$ or live?in) adj3 (batter\$ or perpetrator\$ or abus\$ or violen\$ or beat\$ or assault)).tw.
- 7.or/1-6
- 8.cognitive therapy/
- 9.cognitive behavior therapy/
- 10.rational emotive behavior therapy/
- 11.exp behavior therapy/
- 12.behavior modification/
- 13.cognitive assessment/
- 14.(cognitiv\$ adj3 (therap\$ or train\$ or techni\$ or question\$ or approach\$ or assessment\$)).tw.
- 15.(behavio?r\$ adj3 (therap\$ or train\$ or modif\$ or experiment\$)).tw.
- 16.(rational\$ adj3 emotive\$).tw.
- 17.cbt.tw.
- 18.schema/
- 19.(schema\$ or schemata\$).tw.
- 20.exp imagery/ or conceptual imagery/
- 21.imager\$.tw.

22.cognitive maps/
23.mental models/
24.((cognitiv\$ or mental\$) adj3 (map\$ or model)).tw.
25.(socratic\$ adj3 (question\$ or method\$ or dialogue\$ or strateg\$ or sequence\$)).tw.
26.(dysfunctional adj2 (thought\$ or assumption\$ or rule\$ or appraisal\$ or belief\$ or attitude\$ or scheme\$)).tw.
27.(automatic adj2 (thought\$ or process\$)).tw.
28.(nat or nats).tw.
29.reattribution\$.tw.
30.((key or core) adj2 belief).tw.
31.or/8-29
32.7 and 31
33.limit 32 to yr="2005 - 2006"

Additional tables

06 ERIC search strategy

ERIC

ERIC

Search by Sigrun Espelien Aasen

Database: ERIC, searched 1966 to September 2006.

Date: 29.11.2006

Number of hits: 1

CBT_Eric via Ovid

1.family violence/

2.battered women/

3.(abuse\$ adj3 (wom?n or partner\$ or spouse\$ or female\$ or wife or wives or domestic\$)).tw.

4.(batter\$ adj3 (wom?n or partner\$ or spouse\$ or female\$ or wife or wives)).tw.

5.(violen\$ adj3 (partner\$ or spous\$ or family or families or domestic\$ or conjugal\$)).tw.

6.or/1-5

7.cognitive restructuring/

8.exp behavior modification/

9.(cognitiv\$ adj3 (therap\$ or train\$)).tw.

10.(behavio?r\$ adj3 (therap\$ or train\$)).tw.

11.(behavio?r\$ adj3 modif\$).tw.

12.or/7-11

13.6 and 12

Search in ERIC via CSA Illumina 29.11.06

((DE="family violence") or (DE="battered women") or ((TI=abuse* or AB=abuse*) within 3 (TI=(wom*n or partner* or spouse* or female* or wife or wives or domestic*) or AB=(wom*n or partner* or spouse* or female* or wife or wives or domestic*))) or ((TI=batter* or AB=batter*) within 3 (TI=(wom*n or partner* or spouse* or female* or wife or wives) or AB=(wom*n or partner* or spouse* or female* or wife or wives))) or ((TI=violen* or AB=violen*) within 3 (TI=(partner or spous* or family or families or domestic* or conjugal*) or AB=(partner or spous* or family or families or domestic* or conjugal*))) and (((DE="cognitive restructuring") or (DE=("behavior modification" or "contingency management" or "desensitization"))) or

((TI=(cognitiv* or behavio*r) or AB=(cognitiv* or behavio*r)) within 3 (TI=(therap* or train*) or AB=(therap* or train*))) or (TI=(behavio*r* within 3 modif*) or AB=(behavio*r* within 3 modif*)) or (TI=cbt or AB=cbt)

1 result found in Multiple Databases +

346 results found in Community of Scholars: Social Science

2 results found in Web Resources Related to the Social Sciences/Humanities

Date Range: 2005 to 2007

Additional tables

07 Sociological Abstracts search strategy

Sociol. Abstracts

CBT_Sociological Abstract2 search via Ovid

Searched 1963 to September 2006

- 1.exp spouse abuse/
- 2.battered women/
- 3.family violence/
- 4.((family or domestic or conjugal or partner) adj3 violence).tw.
- 5.((abus\$ or batter\$ or beat\$ or assault\$) adj3 (wom?n or partner\$ or spouse\$ or female\$ or wife or wives or domestic\$ or fiance or cohabitant\$ or live?in)).tw.
- 6.((male\$ or men or man or partner\$ or spouse\$ or husband or fiance or cohabitant\$ or live?in) adj3 (batter\$ or perpetrator\$ or abus\$ or violen\$ or beat\$ or assault)).tw.
- 7.or/1-6
- 8.behavior modification/
- 9.treatment programs/
- 10.treatment methods/
- 11.(cognitiv\$ adj3 (therap\$ or train\$ or techni\$ or question\$ or approach\$ or assessment\$)).tw.
- 12.(behavio?r\$ adj3 (therap\$ or train\$ or modif\$ or experiment\$)).tw.
- 13.(rational\$ adj3 emotive\$).tw.
- 14.cbt.tw.
- 15.(schema\$ or schemata\$).tw.
- 16.exp images/
- 17.imager\$.tw.
- 18.cognitive mapping/
- 19.((cognitiv\$ or mental\$) adj3 (map\$ or model)).tw.
- 20.(socratic\$ adj3 (question\$ or method\$ or dialogue\$ or strateg\$ or sequence\$)).tw.
- 21.(dysfunctional adj1 (thought\$ or assumption\$ or rule\$ or appraisal\$ or belief\$ or attitude\$ or scheme\$)).tw.
- 22.(automatic adj1 (thought\$ or process\$)).tw.
- 23.(nat or nats).tw.
- 24.reattribution\$.tw.
- 25.((key or core) adj1 belief\$).tw.
- 26.or/8-25

27.7 and 26

Additional tables

08 Bibliography of Nordic Criminology search strategy

Bib. Nordic Crim.

Bibliography of Nordic criminology
Searched 1999 to December 2006

Bibliography of Nordic criminology (<http://www.nsfk.org/>) was searched on December 11 2006 by GS, using the textword 'violence' and limiting the search to 2003-2006. The database was searched from 1999 up to 2003 by Torill Johme. There were 152 hits, but none were judged to be relevant.

Additional tables

09 C2-SPECTR search strategy

C2-SPECTR

C2-SPECTR

C2-SPECTR was searched on December 12, 2006 by GS. All indexed fields or all non-indexed fields were searched for the term 'violence'. Of 49 hits, none were judged as relevant.


Additional tables

10 Additional methods for future updates

Issue	Method
Cluster randomised controlled trials	In cluster-randomised trials, the elements are groups of individuals (e.g. courts, jurisdictions, prisons, geographical areas), rather than individuals themselves. In such studies, care should be taken to avoid unit-of-analysis errors. If there for instance are a total of 100 offenders with 25 offenders in each of four jurisdictions, and two jurisdictions are randomised to receive the intervention and the other two are randomised to receive the control, the correct N to use in the analysis is not 100 but smaller. The effective sample size of a single intervention group in a cluster-randomised trial is its original sample size divided by a quantity called the design effect. A common design effect is usually assumed across intervention groups. The design effect is $1+(m-1)r$, where m is the average cluster size and r is the intracluster correlation coefficient. If we include any cluster randomised controlled trials in this review, we try to measure the intra-cluster correlation. The total variance in the outcome can be partitioned into variance between groups (VBG) and variance within groups (VWG). The intracluster correlation (ICC) is calculated as $VBG/(VBG+VWG)$. But the ICC is seldom reported in the primary studies. The number of participants can be used in the analyses if the ICC is used as a correcting factor. For dichotomous data both the number of participants and the number experiencing the event can be divided by the same design effect (Higgins 2005).
Number needed to treat	For statistically significant meta-analyses, we plan to compute the number needed to treat (NNT).
Sensitivity analyses	If the number of included studies is sufficient, we will assess the impact of differing methodological quality by sensitivity analyses.


Additional figures

Fig 01 - Funnel plot



Additional figures

Fig 02 - Flowchart



Notes

Unpublished CRG notes

Exported from Review Manager 4.3

Published notes

This review is co-registered within the Cochrane Developmental, Psychosocial and Learning Problems Group (Cochrane Collaboration).

Amended sections

Cover sheet

Synopsis

Abstract

Background

Objectives

Criteria for considering studies for this review

Search strategy for identification of studies

Methods of the review

Description of studies

Methodological quality of included studies

Results

Discussion

Reviewers' conclusions

Acknowledgements

Potential conflict of interest

References to studies

Other references

Characteristics of included studies

Characteristics of excluded studies

Comparisons, data or analyses

Additional tables and figures

Contact details for co-reviewers

Asbjørn Steiro
The Norwegian Health Services Research Centre
PB 7004 St. Olavs plass
Oslo
NORWAY
N-0130
E-mail: ast@kunnskapssenteret.no

Aina Winsvold
Norwegian Knowledge Centre for Health Services
PB7004
Oslo
NORWAY
N-0130
E-mail: awi@nokc.no

Jocelyne Clench-Aas
Norwegian Institute of Public Health
PO BOX 4404 Nydalen
Oslo
NORWAY
N-0403
Telephone 1: +47 23408361
E-mail: joel@fhi.no


Therese K Dalsbø
Almeveien 36
Oslo
NORWAY
0855
Telephone 1: +47 48271234
E-mail: t.dalsbo-alumni@lse.ac.uk

Review: Cognitive behavioural therapy for men who physically abuse their female partner (Version 04)


Total number of included studies: 6

Comparison or outcome	Studies	Participants	Statistical method	Effect size
01 CBT versus control				
01 Proportion new violence	4	1771	RR (random), 95% CI	0.86 [0.54, 1.38]
02 CBT versus other therapy				
01 Proportion new violence			RR (random), 95% CI	No total
02 Frequency of violence	1	58	SMD (random), 95% CI	0.30 [-0.22, 0.81]
03 Any violence	1	58	RR (random), 95% CI	1.50 [0.27, 8.32]


Review: Cognitive behavioural therapy for men who physically abuse their female partner (For publication)
 Comparison: 01 CBT versus control
 Outcome: 01 Proportion new violence



Review: Cognitive behavioural therapy for men who physically abuse their female partner (For publication)
 Comparison: 02 CBT versus other therapy
 Outcome: 01 Proportion new violence



Review: Cognitive behavioural therapy for men who physically abuse their female partner (For publication)
 Comparison: 02 CBT versus other therapy
 Outcome: 02 Frequency of violence



Review: Cognitive behavioural therapy for men who physically abuse their female partner (For publication)
 Comparison: 02 CBT versus other therapy
 Outcome: 03 Any violence

