

Supplementary Material

Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, Alaska, 2002–2014

Suggested Citation: Alejandro Azofeifa, Margaret E. Mattison, and Rob Lyster. Supplementary Material. State Level Data: Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, Alaska, 2002–2014. Center for Behavioral Health Statistics and Quality. (2016). Substance Abuse and Mental Health Services Administration, Rockville, MD.

Notes: A) Two years of data were combined to allow for reliable state-level prevalence estimates. Some state-level prevalence estimates for combined years have been suppressed (indicated by an asterisk (*)) symbol in each cell based on the following National Survey on Drug Use and Health (NSDUH) suppression criteria. Prevalence estimates are suppressed if any of the following occurred: 1) the prevalence estimate is $<0.005\%$ or $>99.995\%$; 2) the relative standard error (RSE) of the negative natural logarithm of the estimated proportion p (where p is the prevalence divided by 100) is >0.175 if the prevalence is $\leq 50\%$; 3) the RSE of the negative natural logarithm of $(1-p)$ is >0.175 if the prevalence is $>50\%$; 4) the actual sample size is <100 ; or 5) the effective sample size (defined as the actual sample size divided by the design effect) is <68 . B) The estimates provided in this supplementary material for each state and the District of Columbia are from design-based direct estimation procedures (i.e., they are based directly on the respondent sample and the accompanying analytical weights with variances derived through a Taylor-linearization approach using SUDAAN software) as used for the national level estimates (available at: Azofeifa A, Mattison ME, Schauer G, McAfee T, Grant A, Lyster R. National Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, United States, 2002–2014. MMWR Surveill Summ 2016;65(Nb, SS-11):1–25. DOI: <http://dx.doi.org/10.15585/mmwr.ss6511a1>). Caution is advised when comparing data from other NSDUH reports that are based on different design-based estimation procedures. C) Interpretation of state-level trend data should be done with caution as some prevalence estimates are suppressed based on the above criteria and survey cycle differences may not be statistically significant.

BILAG 1

Table 1. Percentage of past month marijuana use among all persons aged ≥ 12, by selected demographic characteristics — National Survey on Drug Use and Health, Alaska, 2002–2014

Characteristics	2002–2003	2003–2004	2004–2005	2005–2006	2006–2007	2007–2008	2008–2009	2009–2010	2010–2011	2011–2012	2012–2013	2013–2014
Age group (yrs)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	9.9 (8.3–11.8)	10.6 (9.0–12.4)	11.5 (9.3–14.0)	9.3 (7.5–11.4)	8.4 (6.7–10.4)	9.8 (7.9–12.1)	12.2 (10.0–14.8)	12.8 (10.0–16.1)	12.9 (10.4–15.9)	14.1 (11.7–16.8)	13.0 (10.8–15.7)	12.4 (10.2–15.0)
12–17	10.7 (8.2–13.8)	13.0 (10.1–16.7)	11.5 (8.5–15.4)	9.1 (6.7–12.2)	7.8 (5.5–10.9)	7.6 (5.4–10.7)	9.3 (6.8–12.6)	8.2 (6.0–11.3)	7.7 (5.3–11.1)	7.3 (5.2–10.3)	5.9 (4.3–8.2)	7.9 (5.8–10.6)
18–25	25.4 (21.0–30.4)	23.2 (19.0–27.9)	21.1 (17.4–25.5)	20.3 (16.7–24.3)	19.7 (15.4–25.0)	20.5 (15.9–26.1)	26.1 (22.0–30.6)	25.3 (20.3–31.0)	25.1 (20.0–30.9)	25.6 (21.2–30.7)	21.6 (17.5–26.3)	22.0 (17.4–27.3)
≥26	7.2 (5.3–9.6)	7.8 (6.0–10.1)	9.5 (7.1–12.7)	7.2 (5.0–10.2)	6.4 (4.2–9.4)	8.2 (6.0–11.1)	10.0 (7.6–13.0)	10.9 (7.7–15.3)	11.2 (8.5–14.7)	12.8 (10.0–16.3)	12.4 (9.6–15.8)	11.2 (8.5–14.6)
26–34	15.9 (11.7–21.2)	12.8 (8.3–19.2)	13.2 (8.6–19.5)	9.8 (6.1–15.3)	10.2 (5.5–18.3)	15.8 (10.3–23.4)	16.2 (11.0–23.2)	13.2 (8.6–19.7)	16.8 (11.5–24.0)	22.6 (16.2–30.5)	15.7 (10.9–22.1)	14.9 (10.9–20.1)
35–44	8.4 (5.3–13.0)	12.7 (8.2–19.2)	15.6 (10.3–23.0)	10.3 (6.2–16.7)	7.5 (4.3–12.7)	8.5 (4.6–15.2)	9.1 (5.2–15.6)	13.3 (7.3–23.1)	13.2 (7.3–22.7)	12.5 (7.1–21.1)	15.3 (9.4–24.0)	12.6 (8.0–19.3)
45–54	6.6 (3.1–13.7)	5.9 (2.6–12.8)	6.8 (3.5–12.7)	5.6 (2.7–11.2)	5.3 (2.8–9.7)	7.0 (3.5–13.5)	12.4 (7.6–19.5)	14.2 (8.6–22.7)	14.2 (8.6–22.7)	10.4 (5.5–18.9)	10.1 (4.9–19.7)	11.8 (7.0–19.2)
55–64	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
≥65	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Sex												
Male	12.2 (9.6–15.4)	12.9 (10.7–15.6)	13.7 (10.7–17.4)	11.4 (8.8–14.8)	9.9 (7.3–13.4)	12.2 (9.2–16.1)	15.7 (12.1–20.1)	16.2 (11.8–21.7)	17.5 (13.9–21.7)	18.2 (14.4–22.7)	14.9 (12.0–18.4)	13.4 (10.8–16.6)
Female	7.6 (5.6–10.3)	8.2 (6.3–10.7)	9.2 (7.1–11.9)	7.1 (5.4–9.4)	6.8 (4.9–9.5)	7.4 (5.2–10.4)	8.6 (6.5–11.4)	9.3 (7.0–12.3)	8.1 (5.5–11.8)	9.8 (7.6–12.5)	11.1 (8.1–15.0)	11.3 (8.1–15.5)
Race/ethnicity												
Non-Hispanic white	9.4 (7.4–11.7)	11.2 (9.1–13.8)	12.8 (10.1–16.1)	8.4 (6.4–11.1)	7.2 (5.1–9.9)	10.4 (7.9–13.5)	12.5 (9.7–15.9)	12.6 (9.3–16.8)	12.0 (9.4–15.2)	13.0 (10.3–16.2)	12.6 (10.0–15.8)	11.2 (8.9–13.9)
Non-Hispanic black	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic American Indian/Alaska Native	13.6 (8.7–20.7)	15.0 (10.1–21.7)	12.9 (8.3–19.4)	13.3 (9.4–18.4)	12.9 (9.2–17.8)	11.3 (7.7–16.3)	21.4 (15.0–29.8)	22.0 (14.3–32.5)	22.2 (14.4–32.5)	(*)	(*)	(*)
Non-Hispanic Hawaiian/Other Pacific Islander	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic Asian	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic two or more races	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Hispanic	(*)	(*)	(*)	4.9 (2.3–9.9)	(*)	(*)	7.5 (3.6–15.2)	(*)	(*)	9.7 (5.2–17.5)	9.4 (4.6–16.3)	7.9 (3.6–16.5)
Current employment status												

Full-time	9.9 (7.5-13.0)	10.6 (8.1-13.8)	11.2 (8.8-14.1)	7.6 (5.7-10.0)	7.7 (5.4-10.8)	10.1 (7.4-13.6)	11.0 (8.1-14.8)	12.6 (9.0-17.5)	13.0 (9.8-16.9)	14.0 (10.6-18.3)	13.7 (10.2-18.3)	12.3 (9.2-16.3)
Part-time	7.5 (4.5-12.3)	7.9 (4.0-15.0)	10.9 (6.7-17.2)	11.1 (6.8-17.4)	11.2 (5.8-20.4)	9.1 (4.8-16.6)	14.4 (8.9-22.7)	15.4 (9.0-25.1)	13.5 (7.2-24.2)	20.2 (13.2-29.5)	(*)	14.4 (8.4-23.7)
Unemployed	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Other ²	9.6 (6.2-14.6)	9.1 (5.6-14.4)	11.5 (6.8-18.7)	11.3 (6.8-18.2)	7.5 (4.6-12.2)	8.3 (5.4-12.5)	12.9 (8.9-18.2)	12.5 (7.8-19.4)	11.9 (7.1-19.2)	11.5 (7.4-17.5)	10.6 (7.1-15.4)	13.0 (8.2-20.2)
Highest level of education completed												
< High school	16.9 (11.0-25.2)	20.5 (13.5-30.0)	22.4 (15.9-30.5)	14.2 (9.8-20.1)	14.2 (9.0-21.5)	15.9 (10.4-23.5)	18.2 (10.8-29.0)	(*)	29.6 (21.1-39.9)	(*)	(*)	20.0 (12.4-30.7)
High school graduate	10.6 (7.5-14.9)	10.5 (6.8-15.7)	12.1 (8.8-16.4)	10.8 (8.3-13.8)	9.1 (6.3-12.9)	9.0 (6.3-12.6)	14.7 (10.9-19.4)	14.9 (11.1-19.6)	15.4 (11.2-20.7)	18.7 (14.2-24.3)	17.1 (12.1-23.6)	16.4 (11.5-22.9)
Some college	11.3 (7.3-17.1)	10.4 (7.4-14.4)	10.2 (7.0-14.8)	8.6 (5.3-13.7)	8.4 (5.1-13.6)	13.7 (8.7-20.8)	16.0 (11.0-22.7)	14.0 (9.0-21.3)	10.2 (6.1-16.6)	11.3 (7.8-16.1)	12.2 (8.5-17.2)	10.8 (8.1-14.4)
College graduate	4.3 (2.4-7.8)	4.5 (2.3-8.8)	6.5 (3.0-13.2)	5.6 (2.8-11.1)	5.1 (2.5-10.0)	5.4 (2.9-10.1)	4.5 (2.4-8.4)	5.3 (2.7-10.1)	6.6 (3.6-11.6)	7.4 (4.2-12.6)	8.8 (5.2-14.8)	8.6 (5.3-13.7)
Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.												
Abbreviation: CI = confidence interval.												
* Past month use of marijuana is defined as those who reported use of marijuana within 30 days preceding the date of interview.												
† Current employment and highest level of education completed are only defined for 18 or older.												
‡ *Other' employment category includes students, persons keeping house or caring for children full time, retired or disabled persons or other persons not in the labor force.												

Table 2. Percentage of past year marijuana use among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Alaska, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	16.2 (14.3-18.2)	16.2 (14.3-18.3)	17.6 (14.8-20.9)	15.6 (12.9-18.7)	14.8 (12.3-17.7)	15.9 (13.4-18.6)	17.2 (14.5-20.3)	18.8 (15.3-22.8)	18.6 (15.9-21.6)	20.6 (18.1-23.2)	20.8 (18.1-23.8)	20.0 (17.3-23.0)
12-17	18.3 (15.1-22.1)	17.6 (14.1-21.9)	16.8 (13.3-21.0)	16.2 (12.9-20.2)	15.6 (12.5-19.4)	15.6 (12.3-19.6)	16.1 (12.8-20.1)	16.5 (13.1-20.7)	17.0 (13.7-20.9)	14.6 (11.7-18.1)	13.9 (11.2-17.1)	17.2 (13.8-21.3)
18-25	40.1 (34.9-45.6)	37.4 (32.7-42.3)	35.1 (30.2-40.3)	33.7 (28.8-39.0)	33.4 (27.5-39.9)	32.3 (26.5-38.8)	34.5 (29.7-39.6)	35.8 (30.3-41.8)	38.4 (32.7-44.4)	39.8 (34.2-45.6)	36.1 (31.2-41.2)	36.6 (31.1-42.6)
≥26	11.7 (9.3-14.5)	11.9 (9.8-14.5)	14.4 (11.0-18.5)	11.9 (8.9-15.9)	11.2 (8.1-15.2)	12.8 (10.0-16.2)	14.1 (10.9-18.1)	15.8 (11.6-21.1)	14.9 (12.1-18.3)	17.8 (15.0-20.9)	18.9 (15.5-22.8)	17.2 (13.9-21.1)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 Past year use of marijuana is defined as those who reported use of marijuana within 12 months preceding the date of interview; — these also include those who reported use of marijuana within 30 days preceding the date of interview.

Table 3. Percentage of daily or almost daily marijuana use in the past year and past month among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Alaska, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Daily or almost daily marijuana use in the past year¹												
Total	2.1 (1.3-3.3)	3.0 (1.9-4.7)	3.7 (2.4-5.7)	2.5 (1.6-3.9)	1.9 (1.2-3.0)	2.3 (1.6-3.3)	2.5 (1.7-3.6)	2.8 (1.9-4.1)	3.6 (2.3-5.7)	4.2 (2.6-6.3)	3.8 (2.7-5.3)	3.5 (2.6-4.8)
12-17	1.6 (0.8-3.2)	2.4 (1.2-4.7)	2.5 (1.4-4.4)	1.7 (0.9-3.2)	1.7 (0.9-3.4)	2.1 (1.1-4.1)	1.1 (0.5-2.6)	0.6 (0.3-1.2)	1.7 (0.9-3.2)	2.0 (1.0-4.0)	1.5 (0.7-3.0)	1.3 (0.6-2.8)
18-25	5.0 (3.1-7.8)	6.0 (4.1-10.0)	6.5 (4.1-10.0)	5.1 (3.5-7.3)	4.0 (2.6-6.1)	5.9 (4.0-8.5)	7.7 (5.5-10.8)	7.2 (5.1-10.0)	8.2 (5.5-12.2)	10.8 (7.5-15.4)	9.8 (6.8-13.8)	7.7 (5.2-11.2)
≥26	1.7 (0.9-3.5)	2.5 (1.2-5.0)	3.4 (1.8-6.1)	2.2 (1.1-4.1)	1.5 (0.8-3.0)	1.7 (1.0-3.0)	1.7 (0.9-3.1)	2.2 (1.2-4.0)	3.0 (1.6-5.5)	3.3 (2.0-5.6)	3.0 (1.9-4.8)	3.0 (2.0-4.6)
Daily or almost daily marijuana use in the past month²												
Total	3.4 (2.4-4.7)	3.8 (2.5-5.7)	4.7 (3.2-6.8)	3.6 (2.5-5.1)	3.4 (2.4-4.7)	4.1 (3.0-5.6)	4.4 (3.4-5.7)	5.2 (3.7-7.2)	5.9 (4.2-8.3)	6.0 (4.4-8.1)	4.9 (3.7-5.6)	5.5 (4.1-7.4)
12-17	2.1 (1.3-3.5)	2.1 (1.1-3.6)	3.1 (1.8-5.3)	2.7 (1.5-4.8)	1.8 (1.0-3.3)	3.6 (2.3-5.6)	3.4 (1.9-5.9)	1.7 (0.8-3.7)	2.1 (1.1-4.1)	2.7 (1.5-4.6)	2.1 (1.3-3.6)	2.0 (1.1-3.9)
18-25	8.3 (5.7-12.0)	9.2 (6.2-13.6)	10.1 (7.1-14.1)	8.9 (6.7-11.6)	7.9 (5.6-11.0)	9.5 (6.5-13.6)	11.8 (8.6-16.0)	10.4 (8.0-13.5)	11.2 (7.8-15.8)	14.1 (10.4-19.0)	12.0 (8.8-16.1)	12.3 (8.7-17.1)
≥26	2.8 (1.7-4.5)	3.1 (1.6-6.0)	3.9 (2.2-6.7)	2.7 (1.5-4.7)	2.8 (1.6-4.7)	3.2 (1.9-5.3)	3.2 (2.0-5.0)	4.7 (2.8-7.7)	5.4 (3.4-8.5)	5.0 (3.3-7.5)	4.0 (2.6-6.0)	4.7 (3.1-7.1)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Past year daily or almost daily use is defined as having used marijuana on 300 or more days in the past year.

² Past month daily or almost daily use is defined as having used marijuana on 20 or more days in the past month.

Table 4. Percentage of daily or almost daily marijuana use in the past year and past month among marijuana users aged ≥12, by age group — National Survey on Drug Use and Health, Alaska, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Daily or almost daily marijuana use in the past year												
Total	13.1 (8.6-19.3)	18.4 (12.3-26.7)	21.0 (14.5-29.4)	16.2 (11.0-23.2)	12.9 (8.6-18.9)	14.6 (10.7-19.7)	14.5 (10.3-20.2)	14.8 (10.1-21.2)	19.6 (13.0-28.5)	20.6 (14.4-28.7)	18.3 (13.2-24.8)	17.6 (13.2-23.2)
12-17	8.7 (4.3-17.0)	13.7 (7.2-24.3)	15.0 (9.1-23.8)	(*) (*)	(*) (*)	13.7 (7.6-23.3)	6.9 (3.0-14.9)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)
18-25	12.4 (8.1-18.6)	16.1 (10.1-24.7)	18.5 (12.2-27.0)	15.2 (10.5-21.5)	11.9 (7.8-17.6)	18.2 (12.5-25.7)	22.3 (16.3-29.9)	20.0 (14.7-26.6)	21.5 (15.6-28.8)	27.2 (20.1-35.7)	27.1 (19.3-36.6)	21.0 (14.5-29.3)
≥26	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	12.3 (6.8-21.1)	14.1 (7.8-24.3)	18.7 (11.3-29.5)	18.7 (11.3-29.5)	15.9 (10.1-24.4)	17.6 (11.6-25.9)
Daily or almost daily marijuana use in the past month												
Total	34.0 (26.0-42.9)	36.2 (26.0-47.8)	40.9 (31.1-51.6)	38.4 (29.6-48.2)	40.2 (30.9-50.3)	42.1 (32.2-52.8)	36.3 (28.8-44.6)	40.8 (32.1-50.1)	45.9 (35.7-56.5)	42.7 (33.0-53.0)	37.9 (27.9-49.2)	44.7 (34.7-55.1)
12-17	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)
18-25	32.8 (23.9-43.0)	39.8 (29.4-51.3)	47.7 (36.9-58.7)	43.7 (34.2-53.7)	40.1 (30.9-49.9)	46.2 (35.8-56.9)	45.4 (35.0-56.2)	41.3 (32.4-50.8)	44.8 (35.7-54.3)	55.1 (45.7-64.3)	55.6 (44.4-66.2)	(*) (*)
≥26	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Analysis limited to marijuana users (i.e., estimate of daily or almost daily use in the past year is among past year marijuana users; estimates of daily or almost daily use in the past month is among past month marijuana users).
 † Past year daily or almost daily use is defined as having used marijuana on 300 or more days in the past year.
 ‡ Past month daily or almost daily use is defined as having used marijuana on 20 or more days in the past month.

Table 5. Percentage of past year marijuana initiation among persons aged ≥12 years at risk for initiation, estimated number at risk for initiation, and mean age at first use of marijuana among past year marijuana initiates, by age group — National Survey on Drug Use and Health, Alaska, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Past Year marijuana initiation among persons at risk for initiation¹												
Total	1.8 (1.3-2.7)	1.8 (1.1-2.8)	1.7 (1.1-2.5)	1.6 (1.1-2.4)	1.6 (1.1-2.5)	2.2 (1.0-4.6)	2.3 (1.1-4.6)	2.6 (1.6-4.4)	2.8 (1.9-4.3)	2.2 (1.7-3.0)	2.9 (1.9-4.3)	3.6 (2.1-6.0)
12-17	5.5 (3.8-7.9)	5.1 (3.1-8.3)	5.3 (3.2-8.5)	4.3 (2.6-6.9)	3.4 (2.1-5.6)	4.1 (2.7-6.2)	6.5 (4.6-9.2)	8.3 (5.9-11.4)	7.8 (5.7-10.7)	6.5 (4.8-8.6)	5.5 (3.6-8.2)	6.7 (4.6-9.6)
18-25	6.7 (3.7-11.7)	6.1 (3.2-11.4)	4.6 (2.2-9.3)	4.5 (2.1-9.1)	6.0 (3.3-10.7)	4.4 (2.1-8.8)	1.7 (0.7-3.9)	3.2 (1.6-6.6)	5.4 (3.4-8.5)	6.0 (3.5-10.3)	6.7 (4.2-10.5)	8.7 (4.8-15.0)
≥26	(*)	(*)	(*)	0.2 (0.0-1.4)	0.2 (0.0-1.4)	(*)	(*)	(*)	(*)	(*)	(*)	1.6 (0.4-6.3)
Estimated no. (in 1,000s) of past year marijuana initiation among those at risk for initiation¹												
Total	5 (3-6)	4 (3-7)	4 (3-6)	4 (3-6)	4 (3-6)	6 (3-12)	6 (3-12)	7 (4-11)	7 (5-11)	5 (4-7)	7 (4-13)	9 (5-15)
12-17	3 (2-4)	3 (2-4)	3 (2-5)	2 (1-4)	2 (1-3)	2 (1-3)	3 (2-5)	4 (3-6)	4 (3-5)	3 (3-5)	3 (2-4)	3 (2-5)
18-25	2 (1-3)	2 (1-3)	1 (1-3)	1 (1-3)	2 (1-4)	2 (1-3)	1 (0-1)	1 (1-2)	2 (1-3)	2 (1-4)	3 (2-4)	3 (2-5)
≥26	(*)	(*)	(*)	0 (0-2)	0 (0-2)	(*)	(*)	(*)	(*)	(*)	(*)	3 (1-11)
Mean age at first use among past year marijuana initiates												
Total	15.9 (15.2-16.7)	16.1 (14.8-17.3)	15.8 (14.4-17.1)	18.0 (13.4-22.6)	19.2 (15.4-23.0)	37.0 (11.2-52.8)	35.2 (9.1-61.3)	24.7 (10.7-38.7)	24.5 (11.0-38.0)	16.6 (15.9-17.4)	20.0 (15.5-24.5)	22.1 (17.0-27.2)
12-17	14.3 (13.8-14.8)	14.0 (13.5-14.5)	13.9 (13.5-14.4)	14.2 (13.4-15.0)	14.3 (13.5-15.1)	14.5 (14.0-15.0)	14.8 (14.1-15.4)	14.9 (14.3-15.5)	15.1 (14.6-15.7)	15.3 (14.6-16.0)	15.3 (14.9-15.8)	14.7 (14.3-15.2)
18-25	19.1 (18.1-20.1)	19.4 (17.7-21.0)	19.8 (17.8-21.7)	18.6 (17.6-19.6)	19.8 (18.3-21.3)	(*)	(*)	19.6 (18.1-21.2)	19.1 (18.2-20.1)	18.9 (18.0-19.8)	19.7 (18.2-21.1)	19.9 (18.9-20.9)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Past year marijuana initiation (initiates) among persons at risk for initiation is defined as those having used marijuana within 12 month period before date of the interview.

Table 6. Percentage of perceived great risk and no risk from smoking marijuana once a month and once or twice a week among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Alaska, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived great risk from smoking marijuana once a month												
Total	26.0 (22.6-29.8)	28.0 (22.2-30.2)	24.9 (21.5-28.7)	27.2 (23.9-30.9)	28.4 (24.1-33.1)	29.5 (25.3-34.1)	26.9 (23.6-30.5)	22.5 (18.8-26.7)	19.7 (16.8-22.9)	18.6 (16.1-21.4)	18.7 (15.9-21.9)	17.4 (14.9-20.1)
12-17	26.0 (22.8-29.5)	28.1 (23.9-32.6)	29.1 (24.1-34.5)	27.3 (22.8-32.2)	27.7 (23.3-32.6)	28.9 (25.0-33.1)	27.5 (23.4-31.9)	27.6 (23.4-32.2)	27.8 (23.4-32.7)	24.0 (20.6-27.7)	19.1 (16.3-22.2)	16.7 (13.6-20.2)
18-25	17.0 (13.4-21.5)	20.1 (16.9-23.7)	18.9 (15.2-23.2)	18.4 (15.1-22.2)	19.3 (16.2-22.9)	17.8 (14.3-21.9)	17.1 (13.9-21.0)	12.9 (9.4-17.4)	10.4 (7.4-14.3)	12.3 (8.5-17.3)	13.9 (10.2-18.8)	11.2 (8.1-15.1)
≥26	27.6 (22.9-32.7)	26.7 (21.9-32.2)	25.4 (21.0-30.3)	28.9 (24.6-33.7)	30.2 (24.7-36.4)	31.9 (26.5-37.7)	28.7 (24.4-33.4)	23.7 (19.1-28.9)	20.3 (16.6-24.6)	19.0 (15.6-22.9)	19.6 (16.0-23.8)	18.6 (15.7-22.0)
Perceived no risk from smoking marijuana once a month												
Total	14.6 (11.8-18.0)	14.0 (11.1-17.6)	15.7 (12.9-19.0)	14.1 (11.4-17.3)	16.0 (13.3-19.1)	16.9 (13.8-20.5)	17.0 (14.4-19.9)	18.8 (15.7-22.3)	21.5 (18.0-25.5)	23.8 (20.4-27.8)	25.1 (22.0-28.4)	27.6 (24.7-30.7)
12-17	11.8 (9.2-15.0)	9.6 (7.1-12.9)	10.5 (7.5-14.4)	10.4 (7.6-14.1)	9.3 (6.7-12.6)	10.6 (7.6-14.4)	12.9 (10.0-16.4)	13.5 (10.0-18.1)	13.8 (10.8-17.6)	14.3 (11.2-18.0)	16.9 (13.7-20.8)	23.3 (19.0-28.1)
18-25	21.6 (18.0-25.6)	24.2 (20.0-29.0)	25.0 (21.2-29.3)	21.5 (17.6-26.0)	22.4 (18.9-26.2)	25.0 (21.0-29.3)	24.7 (20.8-29.0)	27.8 (23.1-32.9)	35.8 (31.4-40.4)	38.4 (34.3-42.7)	37.7 (33.3-42.3)	39.3 (34.5-44.3)
≥26	14.0 (10.5-18.3)	12.9 (9.3-17.7)	14.8 (11.3-19.1)	13.3 (9.9-17.5)	15.9 (12.5-20.0)	16.3 (12.6-20.8)	16.1 (12.9-20.0)	17.7 (13.9-22.3)	19.8 (15.6-24.9)	22.4 (18.4-26.9)	23.8 (20.1-27.9)	25.9 (22.3-29.8)
Perceived great risk from smoking marijuana once or twice a week												
Total	38.9 (35.8-44.2)	39.5 (34.9-44.2)	35.2 (31.7-38.9)	38.2 (33.8-42.9)	41.3 (37.7-45.0)	41.9 (38.2-45.7)	38.1 (34.4-42.0)	30.0 (26.3-33.9)	27.8 (24.5-31.4)	27.4 (24.3-30.7)	24.2 (21.6-27.0)	23.2 (20.8-25.8)
12-17	44.5 (40.5-48.6)	49.8 (44.9-54.8)	49.2 (43.5-54.9)	46.9 (42.3-51.6)	47.1 (42.4-51.7)	48.3 (43.8-53.0)	48.5 (43.3-53.7)	44.8 (40.4-49.3)	41.7 (36.5-47.1)	40.8 (35.5-46.3)	36.1 (31.1-41.3)	29.7 (25.1-34.8)
18-25	28.2 (24.1-32.7)	31.1 (26.9-35.7)	28.7 (24.3-33.6)	28.2 (23.8-33.0)	31.1 (26.3-36.3)	28.4 (23.5-33.9)	25.5 (20.9-30.7)	21.6 (17.0-27.2)	17.5 (13.7-22.0)	18.0 (14.1-22.7)	17.6 (14.1-21.7)	13.8 (10.5-17.9)
≥26	41.1 (35.8-46.6)	39.1 (33.4-45.2)	34.0 (29.7-38.5)	38.7 (33.2-44.5)	42.3 (37.5-47.2)	43.5 (38.7-48.5)	39.0 (34.5-43.7)	29.5 (25.0-34.5)	27.9 (23.5-32.7)	27.3 (23.4-31.5)	23.8 (20.8-27.2)	24.2 (21.3-27.3)
Perceived no risk from smoking marijuana once or twice a week												
Total	8.8 (6.9-11.2)	7.8 (5.9-10.3)	9.8 (7.6-12.5)	9.4 (7.1-12.3)	10.0 (7.5-13.3)	10.6 (8.4-13.4)	12.1 (9.7-14.9)	14.4 (11.3-18.2)	16.7 (13.5-20.4)	17.8 (14.5-21.6)	18.5 (15.3-22.2)	20.6 (17.9-23.6)
12-17	6.5 (4.9-8.6)	4.6 (3.3-6.5)	5.5 (3.6-8.3)	6.2 (4.1-9.3)	5.3 (3.4-8.1)	7.1 (4.8-10.5)	10.2 (7.8-13.1)	9.6 (6.9-13.2)	8.9 (6.6-12.0)	8.4 (6.4-11.1)	12.4 (8.9-17.0)	18.3 (14.2-23.3)
18-25	15.6 (12.2-19.7)	12.8 (9.4-17.2)	13.0 (9.8-17.0)	14.6 (10.9-19.2)	16.7 (13.3-20.9)	20.4 (16.7-24.6)	20.3 (16.6-24.5)	20.7 (16.6-25.5)	28.8 (24.6-33.3)	31.7 (27.3-36.4)	30.4 (25.6-35.6)	31.9 (27.1-37.1)
≥26	8.1 (5.7-11.4)	7.4 (5.1-10.8)	9.9 (7.2-13.5)	8.9 (6.2-12.6)	9.5 (6.3-14.2)	9.3 (6.7-12.9)	10.8 (7.9-14.5)	13.8 (10.0-18.8)	15.4 (11.7-20.2)	16.5 (12.6-21.2)	17.1 (13.2-21.8)	18.7 (15.4-22.6)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
Abbreviation: CI = confidence interval.
Perceived great risk and no risk from smoking marijuana is defined as those who reported that smoking marijuana once a month and once or twice a week might cause harm [great risk] and might not cause harm [no risk].

Table 7. Percentage of perceived parental disapproval of trying marijuana once or twice and using marijuana once a month or more among persons aged 12 to 17 — National Survey on Drug Use and Health, Alaska, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	%	%	%	%	%	%	%	%	%	%	%	%
	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)
Perceived parental disapproval of trying marijuana once or twice ¹												
12–17 years	93.8 (90.4-96.1)	94.2 (91.7-96.0)	94.8 (91.8-96.7)	95.3 (92.6-97.1)	96.3 (94.1-97.7)	94.9 (92.3-96.7)	93.9 (91.0-95.9)	94.9 (91.5-97.0)	94.4 (91.3-96.5)	94.2 (92.0-96.9)	93.1 (90.3-95.2)	92.6 (89.7-94.7)
Perceived parental disapproval of using marijuana once a month or more ²												
12–17 years	95.1 (92.2-97.0)	94.5 (91.8-96.3)	95.1 (92.3-96.9)	96.6 (94.4-97.9)	97.7 (96.0-98.7)	95.8 (93.4-97.3)	94.4 (91.2-96.4)	94.9 (91.5-97.0)	94.3 (91.3-96.2)	95.1 (93.1-96.5)	94.3 (91.4-96.2)	93.7 (90.8-95.7)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Perceived parental disapproval of trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of trying marijuana once or twice.

² Perceived parental disapproval of using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of using marijuana once a month or more.

Table 8. Percentage of perceived parental disapproval of trying marijuana once or twice and using marijuana once a month or more among past month marijuana users aged 12 to 17 — National Survey on Drug Use and Health, Alaska, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	%	%	%	%	%	%	%	%	%	%	%	%
	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)
Perceived parental disapproval of trying marijuana once or twice ¹												
12–17 years	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Perceived parental disapproval of using marijuana once a month or more ²												
12–17 years	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Perceived parental disapproval of trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of trying marijuana once or twice.

² Perceived parental disapproval of using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of using marijuana once a month or more. Indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.

Table 9. Percentage of perceived attitudes towards peers trying marijuana once or twice and using marijuana once a month or more among persons aged 12 to 17 – National Survey on Drug Use and Health, Alaska, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived attitudes towards peers trying marijuana once or twice ¹												
12–17 years	74.2 (70.6-77.5)	75.7 (70.9-79.9)	76.5 (70.9-81.2)	79.2 (75.0-82.8)	80.9 (77.1-84.2)	79.4 (75.1-83.1)	77.0 (72.2-81.2)	77.2 (72.4-81.4)	78.5 (74.3-82.2)	81.3 (77.5-84.5)	78.9 (74.6-82.7)	74.6 (69.3-79.2)
Perceived attitudes towards peers using marijuana once a month or more ²												
12–17 years	75.8 (72.2-79.0)	79.0 (74.7-82.6)	79.0 (74.3-83.0)	79.5 (75.5-83.0)	81.2 (77.2-84.6)	80.6 (76.3-84.3)	78.5 (73.8-82.5)	78.4 (73.8-82.5)	79.3 (74.8-83.3)	80.5 (77.0-83.5)	78.3 (73.5-82.4)	74.7 (69.7-79.2)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
¹Disapproving attitudes toward peers trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers trying marijuana once or twice.
²Disapproving attitudes toward peers using marijuana once or a month or more is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers using marijuana once a month or more.

Table 10. Percentage of perceived attitudes towards peers trying marijuana once or twice and using marijuana once a month or more among past month marijuana users aged 12 to 17 – National Survey on Drug Use and Health, Alaska, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived attitudes towards peers trying marijuana once or twice ¹												
12–17 years	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)
Perceived attitudes towards peers using marijuana once a month or more ²												
12–17 years	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
¹Disapproving attitudes toward peers trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers trying marijuana once or twice. Indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.
²Disapproving attitudes toward peers using marijuana once or a month or more is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers using marijuana once a month or more. Indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.

Table 11. Percentage of perceived availability among all persons aged ≥12 and mode of acquisition of marijuana among past year marijuana users aged ≥12 by age group — National Survey on Drug Use and Health, Alaska, 2002–2014

Age group (yrs)	2002–2003		2003–2004		2004–2005		2005–2006		2006–2007		2007–2008		2008–2009		2009–2010		2010–2011		2011–2012		2012–2013		2013–2014	
	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Perceived availability																								
Total	65.5	(61.3–69.6)	68.0	(63.8–72.0)	65.7	(61.5–69.8)	63.9	(60.0–67.6)	60.6	(56.3–64.9)	59.8	(54.8–62.6)	58.7	(54.5–62.8)	59.1	(54.7–63.4)	61.7	(56.3–66.9)	63.6	(59.8–67.2)	62.9	(59.4–66.4)	64.6	(61.3–67.8)
12–17	58.8	(54.0–63.4)	56.7	(51.7–61.6)	57.2	(51.6–62.7)	53.4	(48.1–58.6)	50.3	(44.9–55.7)	51.0	(45.8–56.2)	49.1	(43.8–54.5)	47.3	(42.4–52.3)	46.8	(42.2–51.5)	43.6	(39.7–47.6)	42.2	(38.6–48.0)	47.6	(41.4–53.9)
18–25	81.2	(78.0–84.0)	79.4	(73.7–84.1)	75.7	(70.8–80.0)	75.7	(71.4–79.4)	73.6	(67.6–78.9)	70.4	(63.8–76.2)	69.1	(63.4–74.3)	71.3	(65.8–76.3)	74.0	(68.2–79.0)	73.2	(67.4–78.3)	70.2	(65.8–74.3)	72.0	(67.3–76.3)
≥26	64.1	(59.2–69.5)	68.0	(62.3–73.2)	65.2	(59.5–70.6)	63.5	(58.3–68.3)	59.9	(54.0–65.4)	57.8	(52.5–62.9)	58.2	(52.9–63.3)	58.4	(52.8–63.8)	61.4	(54.4–68.0)	64.5	(59.8–69.1)	64.4	(60.1–68.5)	65.6	(61.3–69.6)
Mode of acquisition of marijuana																								
Bought it																								
Total	45.2	(37.2–53.5)	48.0	(40.3–55.8)	44.5	(36.9–52.5)	38.4	(31.3–46.1)	38.9	(30.7–47.8)	38.6	(30.2–47.9)	38.6	(30.5–47.3)	46.2	(37.4–55.1)	46.2	(44.7–59.3)	46.2	(38.2–54.3)	45.4	(36.9–54.1)	43.3	(36.4–50.5)
12–17	42.9	(33.7–52.7)	42.1	(32.2–52.8)	34.4	(24.1–46.4)	(*)	(*)	(*)	(*)	(*)	(*)	27.8	(18.6–39.4)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
18–25	40.5	(33.3–48.1)	39.9	(32.3–48.0)	43.5	(35.7–51.6)	42.7	(34.4–51.5)	39.8	(31.5–48.7)	49.6	(38.9–60.4)	52.6	(44.0–61.1)	48.3	(40.4–56.3)	50.2	(43.0–57.5)	51.1	(42.8–59.4)	49.6	(42.0–57.1)	50.0	(40.1–59.9)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	
Traded something for it																								
Total	1.0	(0.3–3.7)	2.1	(0.8–5.5)	1.5	(0.5–4.0)	(*)	(*)	(*)	(*)	3.8	(1.3–10.2)	4.6	(1.8–11.2)	2.0	(0.6–6.3)	1.7	(0.5–5.5)	1.7	(0.5–5.0)	1.2	(0.4–3.7)	2.4	(1.1–5.0)
12–17	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	
18–25	0.8	(0.3–2.8)	2.1	(0.7–5.6)	1.9	(0.6–5.6)	1.6	(0.6–4.3)	1.2	(0.4–3.9)	0.9	(0.2–3.7)	0.8	(0.2–3.4)	(*)	(*)	0.2	(0.0–1.7)	(*)	(*)	2.6	(0.8–8.0)	2.0	(0.7–5.6)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	
Got it for free or shared with someone else																								
Total	52.7	(44.5–60.7)	49.3	(41.7–57.0)	51.0	(44.0–57.9)	55.4	(47.6–63.0)	55.0	(46.3–63.5)	54.8	(45.0–64.3)	55.0	(45.4–64.3)	51.0	(42.3–59.5)	45.2	(38.4–52.1)	50.2	(41.9–58.5)	51.8	(43.0–60.4)	52.0	(44.8–59.1)
12–17	56.5	(46.7–65.8)	57.2	(46.5–67.2)	62.8	(50.7–73.5)	(*)	(*)	(*)	(*)	(*)	(*)	70.8	(59.3–80.1)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
18–25	58.6	(50.9–65.9)	57.8	(48.8–66.3)	54.4	(45.5–63.0)	55.2	(46.6–63.5)	57.6	(48.5–66.2)	47.3	(37.2–57.7)	44.6	(36.4–53.0)	50.7	(42.8–58.6)	49.2	(42.0–56.6)	47.0	(38.4–55.7)	45.5	(37.1–54.1)	46.2	(36.1–56.6)
≥26	(*)	(*)	(*)	(*)	46.9	(36.6–57.5)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	52.8	(42.3–63.0)
Grew it yourself																								
Total	1.1	(0.5–2.3)	0.5	(0.2–1.1)	2.9	(1.5–5.1)	3.5	(2.1–5.6)	2.7	(1.6–4.5)	2.8	(1.7–4.5)	1.7	(0.9–3.0)	0.9	(0.4–1.8)	1.1	(0.5–2.1)	2.0	(1.1–3.5)	1.6	(0.8–2.8)	2.3	(1.2–4.1)

12-17	(0.3-4.6) (*)	(1.2-6.9) (*)	(1.5-7.7) (*)	(0.9-7.5) (*)	(1.0-7.7) (*)	(0.5-5.5) (*)	(0.2-3.6) (*)	(0.3-4.4) (*)	(0.8-4.9) (*)	(0.7-3.9) (*)	(1.1-5.0) (*)
18-25	0.2 (0.0-1.8) (*)	0.3 (0.0-1.8) (*)	1.4 (0.3-5.6) (*)	2.2 (0.7-6.7) (*)	2.0 (0.6-6.4) (*)	1.0 (0.2-4.5) (*)	0.3 (0.0-2.1) (*)	0.7 (0.2-3.0) (*)	2.4 (0.9-6.3) (*)	1.8 (0.6-5.3) (*)	2.4 (0.8-7.0) (*)
≥26	4.7 (1.9-11.0) (*)	4.7 (1.9-11.0) (*)	4.7 (1.9-11.0) (*)	4.7 (1.9-11.0) (*)	4.7 (1.9-11.0) (*)	4.7 (1.9-11.0) (*)	4.7 (1.9-11.0) (*)	4.7 (1.9-11.0) (*)	4.7 (1.9-11.0) (*)	4.7 (1.9-11.0) (*)	4.7 (1.9-11.0) (*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Abbreviation: CI = confidence interval.

* Perceived availability is defined as those who reported that it would be fairly easy or very easy for them to obtain marijuana if they wanted some.

† Mode of acquisition is defined as those who reported how they got the marijuana they used for the last time [i.e., bought it, traded something for it, got it for free or shared with someone else, or grew it yourself]; this question is only asked of those respondents indicating use of marijuana in the past 12 months.

Table 12. Percentage of past year marijuana dependence and abuse* among all persons aged ≥12 by age group — National Survey on Drug Use and Health, Alaska, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	2.6 (1.8-3.6)	3.2 (2.3-4.3)	3.2 (2.3-4.3)	2.8 (2.0-4.0)	2.4 (1.6-3.5)	2.2 (1.5-3.2)	2.0 (1.4-2.8)	1.5 (1.1-2.0)	1.6 (1.1-2.2)	1.8 (1.3-2.5)	1.9 (1.4-2.6)	2.4 (1.5-3.8)
12-17	5.5 (3.8-7.9)	5.4 (3.7-7.9)	5.0 (3.3-7.6)	5.4 (3.6-8.0)	4.3 (2.8-6.6)	4.2 (2.6-6.6)	3.2 (2.1-4.9)	3.7 (2.2-6.1)	5.6 (3.6-8.4)	4.0 (2.6-6.0)	1.9 (0.9-3.8)	2.9 (1.8-4.8)
18-25	7.1 (4.7-10.6)	7.7 (5.6-10.5)	6.7 (5.1-8.8)	6.3 (4.6-8.6)	6.4 (4.2-9.7)	5.1 (3.2-7.9)	5.3 (3.6-7.8)	4.9 (3.3-7.2)	5.1 (3.5-7.3)	6.1 (4.5-8.2)	6.9 (5.2-9.1)	7.1 (5.1-10.0)
≥26	1.2 (0.5-2.7)	1.9 (1.1-3.2)	2.1 (1.2-3.6)	1.7 (0.9-3.3)	1.3 (0.6-2.8)	1.3 (0.6-2.7)	1.1 (0.5-2.2)	0.5 (0.2-1.1)	0.3 (0.1-1.1)	0.7 (0.3-1.5)	0.9 (0.4-2.1)	1.4 (0.5-3.7)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Substance use disorder (also known as dependence or abuse) is defined as meeting criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association [APA], 1994) for either dependence or abuse for illicit drugs or alcohol.

Table 13. Percentage of past year marijuana dependence and abuse* among past year marijuana users aged ≥12, by age group — National Survey on Drug Use and Health, Alaska, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	15.8 (12.0-20.7)	19.5 (14.2-26.1)	18.0 (13.4-23.7)	18.2 (13.7-23.9)	16.1 (11.1-22.8)	13.8 (9.7-19.2)	11.4 (8.3-15.5)	7.8 (5.8-10.6)	8.5 (6.1-11.6)	8.6 (6.1-12.1)	9.0 (6.3-12.5)	11.9 (7.6-18.0)
12-17	30.0 (21.8-39.8)	30.8 (21.5-41.9)	30.0 (21.0-40.8)	(*)	(*)	26.7 (18.0-37.7)	20.1 (13.8-28.4)	(*)	(*)	(*)	(*)	(*)
18-25	17.6 (11.8-25.6)	20.5 (15.2-27.1)	19.2 (14.8-24.6)	18.8 (14.4-24.1)	19.1 (13.2-26.8)	15.7 (10.4-22.9)	15.4 (10.7-21.6)	13.6 (9.2-19.6)	13.3 (9.2-18.8)	15.2 (11.1-20.5)	19.1 (14.8-24.2)	19.5 (13.7-26.9)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	8.1 (4.3-14.5)	3.1 (1.4-6.6)	(*)	3.7 (1.5-8.8)	4.9 (2.0-11.1)	(*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Substance use disorder (also known as dependence or abuse) is defined as meeting criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association [APA], 1994) for either dependence or abuse for illicit drugs or alcohol.

* Analysis limited to past year marijuana users (defined as those who reported use of marijuana within 12 months preceding the date of interview, including those who reported use of marijuana within 30 days preceding the date of interview).

Table 14. Percentage of perception of maximum legal penalty in your state for first offense possession of an ounce or less of marijuana for your own use among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Alaska, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Fine												
Total	35.5 (30.1-41.3)	33.9 (28.6-39.8)	40.0 (35.3-44.9)	36.5 (32.7-40.4)	25.3 (21.3-29.6)	28.9 (24.7-33.5)	32.8 (28.9-37.0)	29.7 (24.9-35.0)	28.0 (23.5-33.0)	29.5 (26.2-33.0)	30.8 (27.4-34.5)	32.0 (28.6-35.7)
12-17	23.6 (19.4-28.2)	22.2 (17.9-27.1)	23.0 (18.1-28.9)	20.8 (15.9-26.8)	18.2 (13.9-23.7)	20.4 (16.2-25.2)	22.2 (19.0-25.7)	20.7 (16.1-26.1)	18.6 (14.4-23.7)	18.9 (15.5-22.8)	18.6 (15.0-22.7)	19.7 (16.1-23.9)
18-25	31.4 (26.5-36.7)	36.0 (30.2-42.3)	43.3 (37.8-48.9)	36.8 (31.4-42.6)	27.1 (21.5-33.7)	30.3 (24.6-36.7)	27.9 (22.1-34.4)	26.5 (21.9-31.8)	34.1 (27.8-40.9)	33.1 (28.8-37.7)	31.9 (27.6-36.4)	31.3 (26.5-36.6)
≥26	39.1 (31.7-46.9)	36.1 (28.6-44.3)	42.9 (36.3-49.7)	39.6 (34.3-45.2)	26.2 (20.7-32.6)	30.2 (24.7-36.3)	35.7 (30.6-41.1)	31.9 (25.8-38.7)	28.1 (22.1-35.0)	30.3 (26.1-34.9)	32.5 (27.5-37.9)	34.3 (29.3-39.6)
Probation												
Total	13.4 (11.2-16.1)	14.4 (11.7-17.5)	13.2 (10.2-16.9)	10.3 (7.9-13.2)	12.1 (9.3-15.6)	10.7 (7.8-14.6)	10.9 (8.3-14.2)	11.5 (8.8-14.9)	9.7 (7.6-12.3)	10.4 (8.1-13.3)	9.5 (7.3-12.3)	8.9 (6.8-11.6)
12-17	17.8 (14.4-21.9)	17.3 (12.9-22.8)	14.5 (9.8-21.0)	12.2 (9.0-16.3)	13.7 (10.7-17.3)	14.6 (10.9-19.3)	13.7 (10.2-18.4)	16.6 (12.4-21.8)	21.8 (17.3-27.0)	18.0 (13.7-23.3)	14.5 (10.5-19.5)	17.3 (13.7-21.7)
18-25	14.1 (10.6-18.6)	9.7 (7.0-13.3)	10.3 (7.6-13.9)	12.5 (9.3-16.5)	12.6 (9.0-17.3)	13.2 (9.1-18.7)	12.5 (9.2-16.8)	13.5 (9.6-18.7)	12.6 (9.0-17.3)	11.1 (8.5-14.4)	11.9 (9.6-14.8)	12.8 (9.8-16.6)
≥26 years	12.3 (9.4-16.0)	14.8 (11.1-19.4)	13.6 (9.5-19.0)	9.4 (6.3-13.7)	11.7 (8.1-16.7)	9.5 (6.0-14.6)	10.0 (6.7-14.8)	10.2 (7.0-14.7)	6.9 (4.4-10.5)	9.0 (6.1-13.1)	8.1 (5.3-12.3)	6.5 (4.0-10.6)
Community Service												
Total	9.5 (6.7-13.3)	11.2 (8.0-15.5)	10.3 (7.5-14.0)	8.4 (6.5-10.7)	7.8 (5.9-10.3)	6.9 (5.4-8.7)	6.1 (4.5-8.2)	5.8 (3.4-9.7)	5.4 (3.1-9.2)	6.6 (4.7-9.3)	6.7 (4.8-9.4)	6.9 (4.2-9.0)
12-17	18.5 (15.1-22.5)	17.5 (13.2-22.7)	16.2 (12.4-20.9)	15.6 (12.4-19.5)	12.1 (9.3-15.6)	12.5 (8.3-18.5)	14.0 (9.9-19.3)	10.0 (6.8-14.4)	8.8 (6.4-12.0)	10.4 (7.4-14.4)	11.7 (8.3-16.1)	13.8 (10.1-18.6)
18-25	8.3 (5.7-12.1)	6.2 (3.9-9.8)	9.2 (6.4-13.0)	9.2 (6.7-12.5)	7.9 (5.2-11.7)	6.4 (3.7-10.7)	4.6 (3.2-6.8)	4.6 (3.1-6.8)	3.8 (2.0-7.0)	8.7 (5.5-13.5)	9.6 (6.8-13.5)	7.2 (4.6-11.0)
≥26	7.7 (4.2-13.8)	11.0 (6.8-17.2)	9.3 (5.7-14.9)	6.7 (4.4-10.1)	7.0 (4.3-11.2)	5.9 (3.9-8.9)	5.1 (3.2-8.3)	5.4 (2.5-11.3)	5.1 (2.3-11.0)	5.5 (3.3-9.3)	5.3 (2.8-9.7)	4.2 (2.3-7.6)
Possible Prison Sentence												
Total	33.9 (29.0-39.3)	33.4 (28.5-38.8)	30.2 (26.4-34.4)	23.7 (20.5-27.3)	19.2 (15.7-23.2)	21.3 (18.0-24.9)	23.6 (20.2-27.3)	24.0 (20.5-27.9)	24.8 (21.1-28.8)	26.0 (22.0-30.5)	25.6 (21.3-30.4)	22.2 (18.9-26.0)

12-17	29.9 (24.9-35.4)	33.4 (27.3-40.0)	35.5 (29.2-42.4)	30.4 (25.3-36.1)	25.7 (21.1-30.9)	27.5 (23.0-32.6)	28.0 (22.9-33.6)	28.3 (22.6-35.0)	28.2 (22.8-34.3)	34.0 (28.4-40.0)	33.2 (26.5-40.6)	24.2 (20.1-28.9)
18-25	35.8 (31.4-40.5)	37.7 (32.1-43.5)	31.7 (26.2-37.8)	24.4 (20.9-28.3)	21.9 (17.8-26.6)	21.0 (16.4-26.4)	27.0 (21.4-33.4)	25.8 (20.6-31.7)	21.9 (18.3-26.0)	24.1 (20.1-28.8)	25.5 (20.4-31.5)	24.1 (19.2-29.9)
≥26	34.5 (27.8-41.8)	32.5 (25.9-39.9)	28.8 (23.4-34.8)	22.2 (17.6-27.5)	17.3 (12.6-23.3)	20.1 (16.1-24.9)	22.1 (17.6-27.2)	22.9 (18.4-28.1)	24.9 (20.2-30.3)	25.2 (19.8-31.4)	24.5 (18.9-31.1)	21.5 (17.1-26.5)
Mandatory Prison Sentence												
Total	7.6 (6.6-10.5)	7.0 (4.9-9.9)	6.3 (4.2-9.3)	6.1 (4.8-7.7)	5.7 (4.7-7.0)	5.2 (3.5-7.7)	5.3 (3.6-7.8)	5.6 (4.0-7.8)	5.7 (4.1-7.7)	4.6 (3.2-6.7)	3.5 (2.4-5.0)	3.3 (2.3-4.7)
12-17	10.2 (7.4-13.9)	9.8 (6.9-13.7)	10.7 (7.6-14.9)	14.1 (10.2-19.3)	17.3 (13.5-22.0)	14.1 (10.0-19.6)	10.5 (7.8-13.9)	12.6 (9.1-17.3)	12.4 (8.9-17.1)	7.8 (5.6-10.8)	7.5 (4.9-11.2)	7.0 (4.8-10.2)
18-25	10.3 (7.4-14.2)	10.4 (7.2-14.7)	5.5 (3.4-8.9)	4.7 (2.5-8.5)	7.5 (4.7-11.8)	7.2 (4.4-11.4)	6.4 (4.4-9.2)	6.8 (4.8-9.6)	5.3 (3.3-8.2)	6.2 (3.9-9.9)	5.5 (3.1-9.5)	5.2 (3.0-8.6)
≥26	6.5 (4.0-10.4)	5.7 (3.0-10.4)	5.5 (2.9-10.4)	4.7 (3.0-7.3)	3.1 (2.0-4.8)	3.1 (1.3-7.0)	4.2 (2.3-7.8)	4.2 (2.4-7.4)	4.6 (2.8-7.5)	3.8 (2.0-6.9)	2.3 (1.2-4.7)	2.2 (1.2-4.2)
No penalty												
Total	DNA	DNA	DNA	15.1 (12.0-18.8)	29.9 (25.2-35.0)	27.1 (22.4-32.2)	21.3 (17.4-25.8)	23.4 (19.0-28.5)	26.6 (22.3-31.3)	22.8 (19.1-27.1)	23.9 (20.0-28.3)	27.6 (24.2-31.4)
12-17	DNA	DNA	DNA	6.8 (4.2-11.0)	13.0 (9.4-17.6)	10.9 (8.1-14.4)	11.7 (8.7-15.6)	11.8 (8.5-16.3)	10.2 (7.6-13.6)	11.0 (7.9-15.1)	14.7 (11.2-18.9)	17.9 (13.8-22.9)
18-25	DNA	DNA	DNA	12.3 (9.5-15.9)	23.0 (19.0-27.6)	22.0 (17.0-28.1)	21.5 (17.2-26.8)	22.8 (18.3-28.1)	22.4 (17.9-27.5)	16.6 (13.2-20.7)	15.6 (12.0-19.9)	19.4 (14.8-25.0)
≥26	DNA	DNA	DNA	17.5 (13.1-22.8)	34.7 (28.3-41.7)	31.2 (25.0-38.2)	22.8 (17.6-29.1)	25.4 (19.3-32.7)	30.4 (24.2-37.4)	26.2 (21.3-31.8)	27.4 (22.3-33.2)	31.3 (26.4-36.5)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.
Abbreviations: CI = confidence interval; DNA = data not available.
Perception of legal penalty for marijuana possession is defined as those who reported their perception of the maximum legal penalty in their state of residence for first offense possession of an ounce or less of marijuana for their own use.

Supplementary Material

Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, Colorado, 2002–2014

Suggested Citation: Alejandro Azofeifa, Margaret E. Mattson, and Rob Lyerla. Supplementary Material. State Level Data: Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, Colorado, 2002–2014. Center for Behavioral Health Statistics and Quality. (2016). Substance Abuse and Mental Health Services Administration, Rockville, MD.

Notes: A) Two years of data were combined to allow for reliable state-level prevalence estimates. Some state-level prevalence estimates for combined years have been suppressed (indicated by an asterisk (*) symbol in each cell) based on the following National Survey on Drug Use and Health (NSDUH) suppression criteria. Prevalence estimates are suppressed if any of the following occurred: 1) the prevalence estimate is $<0.005^*$ or $>99.995^*$; 2) the relative standard error (RSE) of the negative natural logarithm of the estimated proportion p (where p is the prevalence divided by 100) is ≥ 0.175 if the prevalence is $\leq 50^*$; 3) the RSE of the negative natural logarithm of $(1-p)$ is ≥ 0.175 if the prevalence is $>50^*$; 4) the actual sample size is <100 ; or 5) the effective sample size (defined as the actual sample size divided by the design effect) is <68 . B) The estimates provided in this supplementary material for each state and the District of Columbia are from design-based direct estimation procedures (i.e., they are based directly on the respondent sample and the accompanying analytical weights with variances derived through a Taylor-linearization approach using SUDAAN software) as used for the national level estimates (available at: Azofeifa A, Mattson ME, Schauer G, McAfee T, Grant A, Lyerla R. National Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, United States, 2002–2014. MMWR Surveill Summ 2016;65(No. SS-11):1–25. DOI: <http://dx.doi.org/10.15585/mmwr.ss6511a1>). Caution is advised when comparing data from other NSDUH reports that are based on different design-based estimation procedures. C) Interpretation of state-level trend data should be done with caution as some prevalence estimates are suppressed based on the above criteria and survey cycle differences may not be statistically significant.

Table 1. Percentage of past month marijuana use among all persons aged ≥ 12, by selected demographic characteristics — National Survey on Drug Use and Health, Colorado, 2002–2014

Characteristics	2002–2003	2003–2004	2004–2005	2005–2006	2006–2007	2007–2008	2008–2009	2008–2010	2010–2011	2011–2012	2012–2013	2013–2014
Age group (yrs)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	8.8 (7.4–10.4)	7.0 (5.8–8.4)	6.9 (5.5–8.5)	7.5 (5.8–9.6)	8.8 (7.1–10.9)	10.3 (8.2–12.8)	10.1 (8.2–12.5)	12.1 (9.6–15.2)	11.1 (9.3–13.1)	10.0 (7.6–12.8)	13.1 (10.5–16.3)	15.0 (12.7–17.6)
12–17	10.6 (7.9–13.9)	11.6 (9.2–14.6)	10.4 (8.0–13.4)	7.0 (5.2–9.5)	8.3 (6.2–11.1)	9.9 (7.6–12.8)	11.1 (8.6–14.3)	9.9 (7.3–13.4)	10.8 (7.9–14.6)	10.9 (8.4–14.1)	12.3 (9.0–16.6)	14.9 (11.5–19.0)
18–25	21.9 (17.7–26.9)	18.5 (14.9–22.8)	21.5 (17.0–26.9)	22.9 (18.4–28.3)	22.9 (18.8–27.5)	24.4 (19.6–29.6)	24.8 (19.8–30.5)	26.2 (21.8–31.1)	28.5 (24.5–32.8)	28.5 (23.2–34.4)	29.9 (24.3–36.2)	31.5 (26.1–37.6)
≥26	6.2 (4.7–8.2)	4.3 (3.1–6.1)	3.8 (2.5–5.8)	4.9 (3.2–7.4)	6.5 (4.7–9.0)	7.9 (5.5–11.2)	7.6 (5.5–10.4)	10.0 (7.3–13.5)	8.1 (6.2–10.5)	6.8 (4.5–10.0)	10.4 (7.5–14.1)	12.2 (9.7–15.3)
26–34	11.4 (6.9–16.2)	7.9 (4.0–14.9)	8.3 (4.4–15.1)	9.7 (5.4–17.0)	8.7 (5.3–13.9)	10.1 (6.2–16.2)	14.5 (9.5–21.6)	18.4 (12.7–25.8)	12.0 (6.7–20.7)	8.8 (5.1–14.9)	15.2 (10.3–21.8)	21.9 (15.7–29.7)
35–44	10.2 (6.1–16.7)	7.5 (4.6–12.0)	4.9 (2.5–9.3)	4.2 (2.2–8.1)	6.2 (2.9–12.9)	10.9 (6.4–18.0)	9.8 (6.3–15.0)	8.8 (4.7–15.8)	7.9 (4.4–13.7)	10.7 (6.3–17.5)	16.3 (10.6–24.1)	16.1 (11.4–22.4)
45–54	4.2 (2.1–8.1)	2.1 (0.7–5.8)	2.2 (0.8–5.7)	6.2 (2.9–12.6)	10.6 (6.2–17.6)	9.0 (4.4–17.6)	2.6 (0.9–7.6)	10.0 (4.8–19.5)	10.9 (5.9–19.2)	8.7 (4.3–16.9)	11.6 (6.9–18.8)	10.7 (6.7–16.8)
55–64	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
≥65	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Sex												
Male	11.7 (9.3–14.6)	7.8 (6.0–10.0)	7.7 (5.8–10.2)	10.1 (7.3–13.9)	11.1 (8.3–14.7)	13.0 (9.7–17.3)	13.3 (10.2–17.2)	14.5 (10.7–19.4)	12.9 (9.9–16.6)	12.4 (9.0–16.9)	17.6 (13.3–23.0)	20.7 (17.3–24.6)
Female	5.9 (4.6–7.7)	6.2 (4.9–7.8)	6.0 (4.5–8.0)	4.9 (3.4–7.0)	6.6 (4.6–9.4)	7.5 (5.2–10.6)	7.0 (5.2–9.4)	9.8 (7.1–13.3)	9.3 (7.2–11.9)	7.7 (5.7–10.2)	8.8 (6.7–11.4)	9.4 (7.2–12.1)
Race/ethnicity												
Non-Hispanic white	8.7 (7.0–10.6)	7.1 (5.5–9.1)	7.0 (5.5–8.8)	7.0 (5.4–9.1)	8.7 (6.7–11.4)	10.9 (8.3–14.2)	11.2 (8.7–14.4)	13.3 (10.1–17.1)	11.3 (8.9–14.2)	10.3 (7.4–14.1)	14.5 (11.2–18.6)	15.6 (12.9–18.6)
Non-Hispanic black	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic American Indian/Alaska Native	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic Hawaiian/Other Pacific Islander	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Asian	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic two or more races	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Hispanic	7.5 (4.7–11.7)	5.7 (3.8–8.6)	7.8 (4.5–13.2)	9.1 (5.6–14.3)	7.8 (5.1–11.9)	8.5 (5.6–12.6)	7.0 (4.3–11.2)	8.7 (5.3–13.9)	11.1 (7.3–16.7)	10.3 (7.4–14.3)	11.2 (7.8–15.8)	13.6 (9.6–19.0)
Current employment status												

Full-time	9.4 (7.0-12.5)	6.8 (4.9-9.5)	5.6 (3.8-8.2)	7.2 (5.1-10.0)	10.0 (7.4-13.5)	11.3 (8.2-15.3)	11.4 (8.4-15.3)	13.0 (9.3-18.1)	9.7 (7.0-13.4)	9.0 (6.2-13.1)	14.2 (10.5-18.8)	15.3 (12.3-18.9)
Part-time	10.8 (7.0-16.1)	8.0 (4.9-12.5)	8.6 (5.7-12.8)	9.5 (6.1-14.3)	12.4 (7.1-20.9)	14.7 (9.2-25.2)	9.6 (5.2-17.3)	15.6 (9.8-23.9)	17.5 (11.8-25.1)	12.4 (8.2-18.5)	16.4 (10.9-23.7)	20.3 (13.9-28.7)
Unemployed	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Other ²	3.8 (2.2-6.7)	3.1 (1.7-5.4)	5.6 (3.1-9.9)	5.9 (3.0-11.5)	4.0 (1.8-8.7)	4.8 (2.5-8.8)	6.1 (3.3-11.1)	7.6 (4.3-13.2)	8.5 (4.9-14.5)	9.6 (6.1-14.6)	9.0 (5.8-13.6)	10.4 (7.2-14.8)
Highest level of education completed ¹												
< High school	11.2 (6.7-18.1)	8.2 (5.3-12.5)	12.5 (7.7-19.6)	12.7 (7.6-20.3)	12.6 (7.0-21.6)	12.2 (6.2-22.6)	9.4 (6.0-14.3)	13.7 (8.0-22.5)	14.3 (7.9-24.5)	17.4 (10.7-26.8)	21.5 (14.1-31.4)	18.5 (12.2-26.9)
High school graduate	11.1 (7.7-15.6)	8.3 (5.6-12.2)	7.2 (4.7-10.9)	8.4 (5.8-12.0)	10.1 (6.7-14.9)	9.2 (6.1-13.8)	8.3 (5.0-13.5)	14.3 (9.7-20.5)	13.7 (9.8-18.9)	10.6 (6.7-16.2)	12.5 (8.7-17.5)	17.6 (12.6-23.9)
Some college	8.2 (5.1-12.8)	6.2 (3.5-10.7)	6.5 (4.1-10.0)	6.6 (4.4-9.9)	7.7 (4.8-12.0)	14.2 (9.3-21.0)	14.2 (9.7-20.4)	14.7 (11.5-18.7)	13.4 (9.3-18.9)	10.4 (7.1-14.9)	11.9 (8.3-16.8)	13.5 (10.1-17.9)
College graduate	5.9 (3.4-10.0)	4.1 (1.9-8.5)	3.1 (1.7-5.7)	5.6 (2.7-11.2)	7.2 (4.5-11.2)	7.6 (4.6-12.3)	8.6 (5.6-13.1)	9.6 (5.9-15.2)	7.3 (4.8-11.1)	7.0 (4.0-11.9)	11.9 (7.2-18.9)	13.4 (9.7-18.2)
Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.												
Abbreviation: CI = confidence interval.												
* Past month use of marijuana is defined as those who reported use of marijuana within 30 days preceding the date of interview.												
¹ Current employment and highest level of education completed are only defined for 18 or older.												
² Other employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.												

Table 2. Percentage of past year marijuana use among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Colorado, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	15.7 (13.8-17.7)	14.1 (11.9-16.6)	12.0 (10.0-14.2)	12.5 (10.6-14.7)	13.7 (11.8-15.9)	14.3 (11.9-17.1)	15.7 (12.9-19.0)	18.5 (15.2-22.3)	16.7 (14.6-18.9)	15.9 (13.2-19.1)	19.2 (15.7-23.3)	20.7 (17.9-23.8)
12-17	19.1 (15.9-22.9)	18.8 (15.5-22.6)	17.3 (14.3-20.9)	16.8 (13.7-20.4)	17.7 (14.1-22.0)	17.4 (13.7-21.9)	20.0 (16.1-24.5)	20.5 (16.9-24.7)	20.0 (16.5-24.1)	17.1 (13.9-21.0)	18.4 (14.6-23.0)	22.8 (18.9-27.3)
18-25	35.9 (30.9-41.2)	30.2 (25.8-35.0)	32.2 (26.7-38.4)	36.8 (30.9-43.1)	35.3 (30.5-40.4)	35.4 (29.7-41.5)	37.4 (29.5-45.9)	39.9 (32.7-47.5)	42.7 (37.6-48.1)	41.6 (36.0-47.5)	44.2 (38.2-50.3)	45.2 (40.0-50.6)
≥26	11.6 (9.5-14.2)	10.6 (8.0-14.0)	7.5 (5.6-10.3)	7.7 (5.9-10.0)	9.5 (7.4-12.2)	10.4 (7.9-13.7)	11.6 (8.8-15.1)	14.5 (11.1-18.8)	11.7 (9.7-14.1)	11.4 (8.5-15.1)	15.1 (11.2-19.9)	16.3 (13.2-19.9)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

Past year use of marijuana is defined as those who reported use of marijuana within 12 months preceding the date of interview. — these also include those who reported use of marijuana within 30 days preceding the date of interview.

Table 3. Percentage of daily or almost daily marijuana use in the past year and past month among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Colorado, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Daily or almost daily marijuana use in the past year ¹												
Total	2.4 (1.6-3.7)	1.9 (1.3-2.9)	1.9 (1.2-2.8)	2.0 (1.3-3.1)	2.1 (1.3-3.3)	2.9 (1.8-4.4)	2.8 (1.9-4.3)	3.5 (2.3-5.1)	3.7 (2.6-5.2)	3.3 (2.2-5.0)	4.4 (3.1-6.3)	5.3 (4.1-6.9)
12-17	2.3 (1.3-4.2)	2.1 (1.2-3.7)	1.7 (0.8-3.6)	1.0 (0.4-2.4)	0.9 (0.4-2.0)	1.0 (0.5-2.1)	1.4 (0.7-2.8)	1.9 (0.9-3.9)	1.2 (0.6-2.5)	1.4 (0.6-3.3)	2.4 (1.2-4.5)	2.6 (1.4-4.7)
18-25	6.3 (4.5-8.6)	6.2 (4.4-8.7)	8.0 (5.6-11.4)	8.5 (5.7-12.4)	7.3 (4.9-10.6)	9.6 (6.7-13.6)	9.5 (7.0-12.7)	9.1 (6.6-12.5)	11.2 (8.1-15.4)	11.8 (8.3-16.5)	12.6 (8.9-17.5)	12.9 (9.3-17.4)
≥26	1.8 (0.9-3.6)	1.2 (0.5-2.5)	0.8 (0.3-2.0)	1.0 (0.4-2.4)	1.4 (0.6-3.3)	2.0 (0.9-4.2)	1.9 (0.9-3.9)	2.7 (1.4-4.9)	2.7 (1.6-4.7)	2.1 (1.1-4.2)	3.3 (1.9-5.6)	4.3 (3.0-6.2)
Daily or almost daily marijuana use in the past month ²												
Total	3.8 (2.9-5.1)	2.4 (1.6-3.6)	2.7 (1.9-3.7)	3.2 (2.3-4.4)	3.6 (2.5-5.1)	4.3 (2.9-6.2)	3.8 (2.7-5.5)	5.0 (3.6-6.9)	5.0 (3.8-6.5)	4.2 (2.8-6.1)	5.6 (4.1-7.5)	7.2 (5.7-9.1)
12-17	4.4 (2.8-6.8)	2.0 (1.1-3.7)	1.7 (0.9-3.3)	2.5 (1.5-4.2)	2.6 (1.6-4.3)	2.5 (1.5-4.3)	2.8 (1.6-5.1)	2.2 (0.9-5.4)	0.9 (0.3-2.6)	2.1 (1.1-3.9)	4.5 (2.8-7.0)	3.8 (2.4-6.0)
18-25	10.9 (8.2-14.3)	9.0 (6.6-12.1)	10.3 (7.6-14.0)	11.0 (8.0-14.9)	10.1 (6.9-14.5)	12.2 (8.9-16.4)	12.0 (9.1-15.7)	12.4 (9.3-16.3)	14.4 (11.1-18.6)	15.4 (11.6-20.1)	16.1 (11.8-21.5)	16.3 (12.1-21.5)
≥26	2.5 (1.5-4.1)	1.3 (0.6-3.0)	1.4 (0.7-2.7)	1.9 (1.1-3.4)	2.6 (1.4-4.6)	3.2 (1.7-5.9)	2.6 (1.4-4.7)	4.0 (2.6-6.2)	3.8 (2.6-5.6)	2.5 (1.3-5.0)	3.9 (2.5-6.3)	6.1 (4.5-8.2)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Past year daily or almost daily use is defined as having used marijuana on 300 or more days in the past year.

² Past month daily or almost daily use is defined as having used marijuana on 20 or more days in the past month.

Table 4. Percentage of daily or almost daily marijuana use in the past year and past month among marijuana users¹ aged ≥12, by age group — National Survey on Drug Use and Health, Colorado, 2002–2014

Age group (yrs)	2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014	
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Daily or almost daily marijuana use in the past year																								
Total	15.5 (10.1-23.1)	13.7 (9.2-20.1)	15.6 (10.5-22.6)	16.2 (11.0-23.3)	15.4 (10.0-22.9)	20.0 (13.1-29.2)	18.0 (12.8-24.7)	18.7 (13.2-25.9)	22.3 (16.4-29.6)	21.0 (15.1-28.5)	23.0 (17.2-30.1)	25.6 (20.0-31.1)												
12-17	12.0 (6.5-21.2)	11.4 (6.4-19.4)	5.9 (*)	5.9 (2.6-13.0)	5.0 (2.3-10.9)	5.8 (2.7-12.2)	6.9 (3.4-13.3)	9.1 (4.8-16.8)	6.2 (3.2-11.9)	(*)	(*)	11.4 (6.1-20.3)												
18-25	17.5 (12.7-23.6)	20.5 (15.3-26.8)	24.9 (18.9-32.0)	23.1 (17.0-30.6)	20.6 (15.0-27.7)	27.2 (19.2-37.0)	25.3 (19.2-32.5)	22.8 (17.1-29.8)	26.3 (19.2-34.9)	28.3 (20.7-37.3)	28.5 (20.9-37.5)	28.4 (21.2-37.0)												
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	21.8 (13.7-32.8)	26.7 (20.1-34.6)												
Daily or almost daily marijuana use in the past month																								
Total	(*)	(*)	38.7 (29.4-48.9)	42.4 (31.8-53.7)	40.5 (29.8-52.1)	41.7 (31.7-52.4)	37.8 (29.2-47.3)	41.0 (31.5-51.2)	44.7 (36.7-53.1)	41.9 (33.1-51.3)	42.6 (35.0-50.6)	48.1 (41.0-55.2)												
12-17	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)												
18-25	49.6 (40.6-58.6)	48.5 (38.1-58.9)	48.0 (39.0-57.1)	47.8 (40.2-55.4)	(*)	(*)	48.4 (38.1-58.8)	47.2 (37.8-56.8)	50.7 (42.0-59.3)	54.0 (43.1-64.6)	53.7 (43.0-64.1)	51.6 (41.1-61.9)												
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	49.9 (39.6-60.1)												

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Analysis limited to marijuana users (i.e., estimate of daily or almost daily use in the past year is among past year marijuana users, estimates of daily or almost daily use in the past month is among past month marijuana users).
 † Past year daily or almost daily use is defined as having used marijuana on 300 or more days in the past year.
 ‡ Past month daily or almost daily use is defined as having used marijuana on 20 or more days in the past month.

Table 5. Percentage of past year marijuana initiation among persons aged ≥12 years at risk for initiation, and mean age at first use of marijuana among past year marijuana initiates, by age group — National Survey on Drug Use and Health, Colorado, 2002–2014

Age group (yrs)	2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014	
	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Past Year marijuana initiation among persons at risk for initiation																								
Total	2.0	(1.3-2.9)	1.9	(1.3-2.7)	1.7	(1.1-2.6)	2.3	(1.7-3.2)	2.8	(2.1-3.8)	1.8	(1.4-2.4)	2.0	(1.4-2.9)	2.7	(2.0-3.5)	2.7	(2.1-3.5)	2.9	(1.9-4.5)	3.0	(1.8-4.9)	2.6	(1.8-3.6)
12-17	7.3	(5.1-10.5)	8.8	(6.5-11.9)	7.4	(5.1-10.5)	7.7	(5.3-11.1)	8.7	(6.1-12.3)	6.4	(4.0-10.1)	8.2	(5.5-12.2)	11.2	(8.4-14.7)	12.4	(9.4-16.1)	9.2	(6.8-12.5)	7.2	(4.8-10.6)	9.0	(6.3-12.9)
18-25	4.1	(2.1-7.7)	3.0	(1.2-7.4)	2.7	(1.0-6.9)	5.4	(3.0-9.7)	8.8	(5.9-12.9)	7.2	(3.8-13.0)	6.1	(3.2-11.4)	6.2	(3.5-11.0)	7.2	(4.2-11.8)	7.7	(4.7-12.3)	8.7	(5.3-13.9)	9.3	(5.7-14.9)
≥26	*	(*)	*	(*)	0.3	(0.0-1.8)	0.7	(0.2-2.4)	0.5	(0.1-2.3)	*	(*)	*	(*)	0.1	(0.0-0.6)	0.1	(0.0-0.6)	1.0	(0.2-4.9)	*	(*)	*	(*)
Estimated no. (in 1,000s) of past year marijuana initiation among those at risk for initiation																								
Total	35	(24-52)	34	(24-48)	33	(22-50)	48	(34-66)	54	(40-72)	35	(26-46)	40	(28-57)	52	(40-67)	57	(45-74)	65	(42-101)	59	(35-98)	50	(35-71)
12-17	23	(16-33)	28	(21-38)	24	(17-34)	26	(18-37)	28	(20-40)	21	(13-32)	26	(18-39)	36	(27-47)	41	(31-54)	32	(23-43)	24	(16-35)	30	(20-42)
18-25	8	(4-15)	6	(2-15)	5	(2-13)	11	(6-19)	18	(12-26)	14	(8-26)	14	(7-26)	14	(8-25)	15	(9-25)	16	(10-26)	18	(11-29)	20	(12-32)
≥26	*	(*)	*	(*)	4	(0-27)	11	(3-37)	8	(2-32)	*	(*)	*	(*)	1	(0-9)	1	(0-9)	17	(4-82)	*	(*)	*	(*)
Mean age at first use among past year marijuana initiates																								
Total	17.2	(14.3-20.0)	16.3	(14.5-18.1)	17.9	(13.6-22.1)	19.9	(16.1-23.7)	18.3	(15.7-20.8)	16.3	(15.3-17.4)	16.1	(15.2-17.0)	16.4	(15.5-17.2)	16.5	(15.8-17.2)	26.1	(12.0-40.3)	27.0	(11.8-42.2)	16.4	(15.6-17.3)
12-17	14.3	(13.6-15.0)	14.4	(13.9-14.8)	14.5	(14.1-14.9)	14.9	(14.4-15.3)	14.9	(14.3-15.5)	14.8	(14.1-15.5)	14.6	(14.2-15.1)	14.9	(14.6-15.3)	15.1	(14.6-15.5)	14.9	(14.5-15.4)	14.5	(13.8-15.2)	14.6	(14.0-15.1)
18-25	19.5	(18.4-20.5)	*	(*)	*	(*)	19.1	(17.8-20.4)	18.6	(17.6-19.7)	18.6	(17.7-19.5)	18.9	(18.4-19.5)	19.2	(18.4-20.0)	19.8	(18.6-20.9)	19.8	(18.7-20.8)	19.0	(17.8-20.3)	19.1	(18.0-20.1)
≥26	*	(*)	*	(*)	*	(*)	*	(*)	*	(*)	*	(*)	*	(*)	*	(*)	*	(*)	*	(*)	*	(*)	*	(*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Past year marijuana initiation (initiates) among persons at risk for initiation is defined as those having first used marijuana within the 12-month period before the date of the interview.

Table 6. Percentage of perceived great risk and no risk from smoking marijuana once a month and once or twice a week among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Colorado, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived great risk from smoking marijuana once a month												
Total	27.9 (23.4-32.8)	30.6 (26.9-34.6)	31.9 (27.8-36.3)	28.9 (24.8-33.2)	26.8 (22.2-30.5)	26.1 (21.5-30.3)	20.5 (17.0-24.4)	19.6 (16.4-23.2)	23.5 (20.0-27.5)	21.1 (17.8-24.8)	21.1 (17.8-24.8)	17.0 (14.7-19.6)
12-17	28.6 (25.2-32.2)	29.7 (25.6-34.1)	28.2 (23.7-33.1)	26.6 (23.2-30.2)	25.7 (21.6-30.3)	25.6 (21.5-30.3)	22.4 (18.3-27.0)	20.8 (16.8-25.3)	21.7 (18.5-25.3)	18.8 (15.5-22.7)	18.8 (15.5-22.7)	16.9 (13.5-20.9)
18-25	17.7 (13.6-22.8)	22.0 (18.5-25.9)	20.9 (16.6-26.0)	17.9 (13.4-23.5)	14.9 (11.9-18.6)	13.5 (10.3-17.5)	13.0 (9.5-17.7)	10.1 (7.6-13.4)	10.1 (7.7-13.2)	9.9 (7.2-13.6)	9.9 (7.2-13.6)	7.7 (5.7-10.3)
≥26	29.6 (24.0-35.8)	32.3 (27.6-37.3)	34.4 (29.4-39.8)	31.4 (26.2-37.0)	31.5 (26.7-36.7)	28.3 (23.4-33.7)	21.5 (17.3-26.5)	21.1 (17.1-25.8)	26.1 (21.7-31.0)	23.3 (19.4-27.9)	23.3 (19.4-27.9)	18.6 (15.6-22.0)
Perceived no risk from smoking marijuana once a month												
Total	12.4 (10.1-15.1)	11.4 (9.6-13.6)	11.4 (9.3-13.9)	14.9 (12.0-18.3)	16.0 (13.3-19.2)	15.7 (12.7-19.3)	20.1 (16.8-23.9)	22.0 (18.6-25.8)	20.2 (16.3-24.9)	23.0 (19.5-26.9)	23.0 (19.5-26.9)	24.6 (21.7-27.8)
12-17	9.2 (7.2-11.7)	7.8 (5.6-10.9)	9.1 (6.8-12.1)	8.6 (6.4-11.3)	9.4 (7.4-11.9)	11.9 (8.9-15.6)	12.6 (9.4-16.6)	16.3 (13.6-19.4)	18.2 (14.4-22.7)	19.9 (16.5-23.7)	19.9 (16.5-23.7)	21.2 (17.7-25.2)
18-25	23.3 (19.3-27.8)	21.4 (18.4-24.8)	23.8 (20.0-28.2)	22.1 (18.1-26.7)	29.5 (25.7-33.7)	31.7 (26.9-36.9)	34.3 (29.5-39.5)	39.2 (34.2-44.4)	40.9 (35.3-46.8)	42.5 (37.3-47.8)	42.5 (37.3-47.8)	43.8 (39.1-48.7)
≥26	10.9 (8.3-14.2)	10.1 (7.8-13.1)	9.6 (7.0-12.9)	14.2 (10.9-18.4)	14.5 (11.1-18.8)	13.6 (10.0-18.1)	18.6 (14.7-23.2)	19.7 (15.6-24.5)	16.9 (12.8-22.1)	20.0 (16.0-24.6)	20.0 (16.0-24.6)	21.7 (18.2-25.7)
Perceived great risk from smoking marijuana once or twice a week												
Total	44.6 (40.9-48.5)	44.8 (40.9-48.8)	45.0 (40.8-49.3)	42.8 (39.3-46.3)	42.5 (38.3-46.8)	38.5 (33.8-43.4)	30.2 (26.1-34.6)	29.8 (25.9-34.0)	33.6 (29.0-38.4)	30.6 (27.0-34.6)	30.6 (27.0-34.6)	24.5 (21.4-27.8)
12-17	51.2 (46.8-55.6)	51.6 (46.9-56.3)	50.8 (45.6-56.0)	52.2 (46.6-57.8)	47.7 (43.3-52.0)	44.3 (38.7-50.1)	37.9 (32.8-43.4)	36.4 (31.9-41.2)	39.5 (35.3-43.9)	35.1 (30.1-40.4)	35.1 (30.1-40.4)	29.1 (24.5-34.1)
18-25	28.1 (24.2-32.5)	28.9 (25.2-32.8)	28.9 (24.0-34.3)	26.1 (21.8-31.0)	24.6 (21.1-28.4)	22.8 (18.3-27.9)	19.8 (15.5-25.1)	16.1 (12.5-20.5)	14.4 (10.8-18.8)	13.2 (9.3-18.3)	13.2 (9.3-18.3)	11.6 (8.9-15.1)
≥26	46.7 (41.7-51.7)	46.6 (42.0-51.3)	47.0 (41.8-52.3)	44.4 (40.1-48.8)	44.9 (39.6-50.2)	40.4 (34.9-46.2)	31.1 (26.0-36.7)	31.4 (26.5-36.7)	36.1 (30.9-41.7)	33.1 (28.6-37.8)	33.1 (28.6-37.8)	26.1 (22.3-30.3)
Perceived no risk from smoking marijuana once or twice a week												
Total	7.1 (5.4-9.4)	6.1 (4.7-8.0)	5.9 (4.3-8.0)	7.1 (5.1-9.6)	9.2 (7.1-11.8)	9.6 (7.4-12.3)	12.6 (9.9-15.8)	14.3 (11.9-17.1)	13.3 (10.4-16.9)	15.6 (12.7-18.9)	15.6 (12.7-18.9)	18.0 (15.6-20.7)
12-17	5.5 (3.8-7.7)	4.3 (2.7-6.8)	4.7 (3.2-6.8)	3.6 (2.2-5.0)	3.7 (2.4-5.8)	7.5 (4.9-11.2)	9.4 (6.5-13.4)	10.8 (8.7-13.4)	11.6 (8.7-15.2)	13.6 (10.3-17.8)	13.6 (10.3-17.8)	16.3 (12.9-20.5)
18-25	14.4 (10.7-19.3)	12.1 (9.3-15.7)	13.4 (10.3-17.2)	13.1 (9.9-17.2)	20.6 (16.8-25.1)	21.1 (15.8-27.6)	22.0 (18.2-26.2)	26.9 (23.0-31.2)	30.7 (26.7-35.1)	32.8 (27.6-38.5)	32.8 (27.6-38.5)	34.3 (29.8-39.1)
≥26	6.1 (4.1-8.9)	5.3 (3.6-7.8)	4.7 (2.9-7.7)	6.3 (4.0-9.8)	7.9 (5.4-10.4)	7.9 (5.5-11.2)	11.3 (8.2-15.4)	12.5 (9.5-16.4)	10.5 (7.3-14.9)	12.8 (9.5-17.0)	12.8 (9.5-17.0)	15.4 (12.4-18.9)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
Abbreviation: CI = confidence interval.
Perceived great risk and no risk from smoking marijuana is defined as those who reported that smoking marijuana once a month and once or twice a week might cause harm (great risk) and might not cause harm (no risk).

Table 7. Percentage of perceived parental disapproval of trying marijuana once or twice and using marijuana once a month or more among persons aged 12 to 17 — National Survey on Drug Use and Health, Colorado, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived parental disapproval of trying marijuana once or twice ¹												
12–17 years	94.5 (91.9-96.4)	93.9 (91.3-95.8)	95.2 (92.9-96.9)	96.3 (93.9-97.7)	94.9 (92.0-96.8)	94.7 (92.1-96.5)	95.0 (92.1-96.9)	95.3 (92.0-97.3)	96.8 (94.7-98.1)	96.6 (94.5-98.0)	95.9 (93.8-97.4)	94.6 (92.3-96.3)
Perceived parental disapproval of using marijuana once a month or more ²												
12–17 years	94.9 (92.0-96.8)	94.9 (92.2-96.7)	96.7 (94.8-97.9)	97.2 (95.4-98.3)	95.5 (92.7-97.3)	95.4 (92.6-97.1)	95.9 (93.2-97.5)	96.3 (93.5-98.0)	96.5 (94.1-97.9)	96.5 (94.0-98.0)	96.1 (93.8-97.5)	94.5 (92.0-96.3)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Perceived parental disapproval of trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of trying marijuana once or twice.

² Perceived parental disapproval of using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of using marijuana once a month or more.

Table 8. Percentage of perceived parental disapproval of trying marijuana once or twice and using marijuana once a month or more among past month marijuana users aged 12 to 17 — National Survey on Drug Use and Health, Colorado, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived parental disapproval of trying marijuana once or twice ¹												
12–17 years	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Perceived parental disapproval of using marijuana once a month or more ²												
12–17 years	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Perceived parental disapproval of trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of trying marijuana once or twice.

² Perceived parental disapproval of using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of using marijuana once a month or more. Indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.

Table 9. Percentage of perceived attitudes towards peers trying marijuana once or twice and using marijuana once a month or more among persons aged 12 to 17— National Survey on Drug Use and Health, Colorado, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
Perceived attitudes towards peers trying marijuana once or twice ¹	77.1 (72.4-81.2) %	76.1 (71.8-80.0) %	75.8 (71.9-79.2) %	77.4 (73.4-81.1) %	75.8 (71.9-79.3) %	76.7 (72.1-80.6) %	76.0 (71.2-80.3) %	75.7 (71.0-79.9) %	75.3 (70.5-79.6) %	76.8 (72.6-80.6) %	75.3 (71.2-79.1) %	74.3 (69.3-78.7) %
Perceived attitudes towards peers using marijuana once a month or more ²	79.6 (75.2-83.5) %	78.2 (74.8-81.3) %	77.5 (74.4-80.3) %	81.3 (77.5-84.4) %	79.8 (75.9-83.2) %	78.0 (73.3-82.1) %	77.0 (72.0-81.3) %	77.2 (72.6-81.2) %	76.4 (72.0-80.3) %	77.8 (74.1-81.2) %	75.9 (71.2-80.0) %	74.6 (70.0-78.7) %

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
¹Disapproving attitudes toward peers trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers trying marijuana once or twice.
²Disapproving attitudes toward peers using marijuana once or a month or more is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers using marijuana once a month or more.

Table 10. Percentage of perceived attitudes towards peers trying marijuana once or twice and using marijuana once a month or more among past month marijuana users aged 12 to 17— National Survey on Drug Use and Health, Colorado, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
Perceived attitudes towards peers trying marijuana once or twice ¹	80.0 (75.0-85.0) %	78.0 (73.0-83.0) %	76.0 (71.0-81.0) %	74.0 (69.0-79.0) %	72.0 (67.0-77.0) %	70.0 (65.0-75.0) %	68.0 (63.0-73.0) %	66.0 (61.0-71.0) %	64.0 (59.0-69.0) %	62.0 (57.0-67.0) %	60.0 (55.0-65.0) %	58.0 (53.0-63.0) %
Perceived attitudes towards peers using marijuana once a month or more ²	82.0 (77.0-87.0) %	80.0 (75.0-85.0) %	78.0 (73.0-83.0) %	76.0 (71.0-81.0) %	74.0 (69.0-79.0) %	72.0 (67.0-77.0) %	70.0 (65.0-75.0) %	68.0 (63.0-73.0) %	66.0 (61.0-71.0) %	64.0 (59.0-69.0) %	62.0 (57.0-67.0) %	60.0 (55.0-65.0) %

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
¹Disapproving attitudes toward peers trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers trying marijuana once or twice, restricted to respondents aged 12–17 years who reported past month marijuana use.
²Disapproving attitudes toward peers using marijuana once or a month or more is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers using marijuana once a month or more; indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.

Table 11. Percentage of perceived availability among all persons aged ≥12 and mode of acquisition of marijuana among past year marijuana users aged ≥12 by age group — National Survey on Drug Use and Health, Colorado, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived availability												
Total	64.1 (59.3-68.6)	66.2 (62.1-70.1)	61.6 (57.5-65.6)	60.7 (56.7-64.5)	62.0 (57.9-66.0)	60.5 (56.2-64.6)	58.7 (54.3-62.9)	61.9 (57.8-65.8)	66.3 (62.0-70.3)	67.4 (63.1-71.5)	72.0 (68.1-75.7)	80.6 (77.4-83.5)
12-17	55.0 (50.1-59.9)	57.1 (52.5-61.6)	56.6 (51.9-61.0)	51.8 (46.6-57.1)	53.6 (48.4-59.7)	55.2 (50.4-59.9)	56.1 (51.5-60.6)	54.6 (49.4-59.7)	55.7 (50.8-60.4)	58.7 (53.0-64.2)	62.4 (57.2-67.3)	61.0 (55.1-66.6)
18-25	84.0 (80.3-87.1)	84.7 (81.1-87.6)	82.5 (78.2-86.2)	82.5 (78.3-86.1)	84.3 (80.8-87.3)	85.6 (83.3-87.6)	85.0 (81.2-88.2)	83.2 (78.8-86.8)	83.1 (79.3-86.2)	83.1 (79.2-86.5)	84.0 (80.0-87.4)	87.7 (84.1-90.7)
≥26	61.7 (55.7-67.4)	64.1 (59.1-68.9)	58.5 (53.3-63.5)	57.9 (52.9-62.8)	59.1 (54.1-64.0)	56.7 (51.6-61.7)	54.4 (49.2-59.5)	59.0 (54.2-63.7)	64.6 (59.4-69.4)	65.7 (60.6-70.5)	71.1 (65.9-75.8)	81.8 (77.7-85.2)
Mode of acquisition of marijuana												
Bought it												
Total	39.2 (32.7-46.1)	37.9 (31.5-44.7)	43.3 (36.7-51.2)	43.2 (35.4-51.5)	45.9 (38.2-53.9)	52.7 (42.1-63.0)	45.0 (37.4-52.9)	43.9 (36.4-51.7)	48.2 (41.5-55.0)	44.1 (36.1-52.5)	46.1 (40.1-52.4)	56.4 (50.6-62.0)
12-17	21.7 (13.9-32.3)	*	*	34.1 (23.7-46.2)	21.9 (14.5-31.8)	*	33.2 (24.1-43.7)	32.1 (24.4-41.0)	33.5 (24.0-44.6)	33.3 (23.2-45.2)	*	*
18-25	34.8 (27.9-42.4)	41.2 (32.3-50.7)	46.5 (39.0-54.2)	41.8 (35.3-48.6)	43.2 (36.4-50.2)	58.9 (52.2-65.3)	60.0 (52.7-66.8)	49.7 (42.0-57.5)	50.2 (42.5-57.9)	51.2 (44.9-57.5)	50.6 (42.3-59.0)	55.1 (45.9-64.0)
≥26	*	*	*	*	*	*	*	43.0 (32.1-54.7)	*	*	45.3 (35.7-55.2)	59.8 (50.7-68.2)
Traded something for it												
Total	0.4 (0.1-1.3)	0.8 (0.3-1.9)	1.1 (0.4-2.9)	0.6 (0.2-2.0)	0.3 (0.1-1.6)	0.4 (0.1-1.7)	0.3 (0.1-1.1)	0.3 (0.1-0.8)	0.4 (0.1-1.9)	1.3 (0.4-4.2)	3.1 (1.4-6.7)	3.3 (1.5-7.2)
12-17	1.9 (0.5-7.5)	*	*	*	0.8 (0.2-3.6)	0.2 (0.0-1.5)	0.2 (0.0-1.9)	*	*	*	*	*
18-25	*	2.2 (0.9-5.3)	3.1 (1.2-7.5)	1.4 (0.4-4.9)	*	*	0.6 (0.1-2.3)	0.5 (0.1-2.1)	*	1.7 (0.5-5.4)	3.0 (1.2-7.2)	2.1 (0.8-5.8)
≥26	*	*	*	*	*	*	*	*	*	*	3.3 (1.0-10.0)	4.2 (1.6-10.8)
Got it for free or shared with someone else												
Total	59.8 (52.9-66.2)	59.8 (52.7-66.5)	54.5 (46.6-62.2)	54.3 (46.4-62.0)	52.1 (45.6-58.5)	46.6 (36.3-57.1)	54.5 (46.6-62.2)	54.7 (46.8-62.3)	49.3 (42.0-56.6)	51.8 (43.0-60.5)	45.8 (38.9-52.8)	37.0 (31.6-42.7)
12-17	75.4 (65.0-83.5)	*	*	64.6 (52.5-75.0)	76.1 (67.6-82.9)	*	66.6 (56.1-75.7)	66.6 (57.6-74.6)	65.4 (54.2-75.1)	66.0 (53.7-76.4)	*	60.1 (48.2-71.0)
18-25	63.6 (55.8-70.6)	55.2 (46.2-64.0)	50.2 (42.2-58.3)	56.3 (48.8-63.5)	55.7 (48.8-62.4)	39.9 (33.2-46.9)	39.0 (31.9-46.6)	47.9 (40.1-55.8)	47.3 (40.0-54.8)	45.9 (39.4-52.4)	43.2 (36.3-50.4)	40.2 (32.1-48.8)
≥26	*	*	*	*	*	*	*	55.9 (44.2-66.9)	*	*	44.9 (34.1-56.1)	31.6 (23.8-41.1)
Grew it yourself												

Total	0.7 (0.2-1.8)	1.5 (0.4-5.1)	1.1 (0.2-4.8)	0.3 (0.1-1.4)	0.2 (0.0-1.1)	1.2 (0.4-3.8)	2.1 (0.7-6.0)	2.7 (0.9-7.5)	5.0 (2.4-9.8)	3.3 (1.6-6.7)
12-17	1.6 (0.4-6.5)	1.4 (0.3-5.6)	0.5 (0.1-2.3)	0.3 (0.1-1.4)	0.2 (0.0-1.1)	1.2 (0.4-3.8)	2.1 (0.7-6.0)	2.7 (0.9-7.5)	5.0 (2.4-9.8)	3.3 (1.6-6.7)
18-25	1.2 (0.3-5.0)	0.2 (0.0-1.3)	0.5 (0.1-2.3)	0.3 (0.1-1.4)	0.2 (0.0-1.1)	1.2 (0.4-3.8)	2.1 (0.7-6.0)	2.7 (0.9-7.5)	5.0 (2.4-9.8)	3.3 (1.6-6.7)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Abbreviation: CI = confidence interval.

* Perceived availability is defined as those who reported that it would be fairly easy or very easy for them to obtain marijuana if they wanted some

* Mode of acquisition is defined as those who reported how they got the marijuana they used for the last time (i.e., bought it, traded something for it, got it for free or shared with someone else, or grew it yourself); this question is only asked of those respondents indicating use of marijuana in the past 12 months.

Table 12. Percentage of past year marijuana dependence and abuse among all persons aged ≥12 by age group — National Survey on Drug Use and Health, Colorado, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	1.9 (1.4-2.5)	1.7 (1.2-2.4)	1.6 (1.1-2.1)	1.9 (1.4-2.6)	2.3 (1.3-3.9)	2.6 (1.6-4.2)	2.3 (1.6-3.3)	2.1 (1.5-2.9)	1.9 (1.4-2.6)	2.2 (1.5-3.2)	2.1 (1.4-3.1)	2.1 (1.4-2.9)
12-17	3.0 (1.8-4.9)	3.9 (2.5-5.9)	4.8 (3.2-7.1)	4.6 (3.0-7.1)	4.2 (2.8-6.3)	4.5 (3.2-6.2)	6.7 (4.9-9.2)	7.1 (4.9-10.0)	5.2 (3.7-7.4)	4.7 (3.2-6.9)	4.6 (2.9-7.3)	4.1 (2.6-6.4)
18-25	9.3 (6.7-12.8)	7.1 (4.9-10.0)	5.9 (3.9-8.8)	6.9 (4.8-9.9)	4.8 (3.2-7.2)	5.9 (3.7-9.5)	6.6 (4.2-10.5)	7.2 (5.1-10.2)	8.5 (6.2-11.5)	8.8 (6.2-12.3)	8.2 (5.4-12.1)	6.1 (3.7-9.9)
≥26	0.4 (0.1-1.3)	0.4 (0.1-1.3)	0.3 (0.1-1.0)	0.7 (0.3-1.6)	1.6 (0.6-4.3)	1.8 (0.8-4.4)	1.1 (0.5-2.2)	0.6 (0.2-1.6)	0.4 (0.1-1.2)	0.8 (0.3-2.1)	0.8 (0.3-2.0)	1.1 (0.6-2.1)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 † Substance use disorder (also known as dependence or abuse) is defined as meeting criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association [APA], 1994) for either dependence or abuse for illicit drugs or alcohol.

Table 13. Percentage of past year marijuana dependence and abuse among past year marijuana users aged ≥12, by age group — National Survey on Drug Use and Health, Colorado, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	11.9 (8.8-15.9)	11.9 (8.2-16.9)	13.0 (9.5-17.5)	15.5 (11.7-20.3)	16.5 (9.6-27.0)	18.4 (11.8-27.4)	14.7 (10.5-20.3)	11.2 (8.3-15.0)	11.6 (8.5-15.5)	13.7 (10.0-18.6)	11.0 (7.9-15.2)	10.0 (7.3-13.5)
12-17	15.7 (9.7-24.3)	20.7 (13.6-30.1)	27.5 (18.8-38.3)	27.4 (19.1-37.7)	23.8 (16.7-32.8)	25.6 (17.9-35.3)	33.7 (24.8-43.9)	34.5 (26.3-43.7)	26.1 (18.8-35.1)	27.3 (18.9-37.6)	*	18.0 (11.6-26.9)
18-25	25.9 (19.5-33.6)	23.3 (16.7-31.6)	18.3 (12.9-25.2)	18.9 (13.4-25.9)	13.7 (9.4-19.3)	16.8 (11.0-24.7)	17.8 (11.2-27.2)	18.1 (12.8-25.0)	19.9 (15.0-25.8)	21.2 (15.8-27.7)	18.5 (13.1-25.5)	13.5 (8.5-20.9)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	4.1 (1.5-10.3)	(*)	(*)	5.2 (2.2-12.1)	7.0 (4.0-12.1)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 † Substance use disorder (also known as dependence or abuse) is defined as meeting criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association [APA], 1994) for either dependence or abuse for illicit drugs or alcohol.
 * Analysis limited to past year marijuana users (defined as those who reported use of marijuana within 12 months preceding the date of interview, including those who reported use of marijuana within 30 days preceding the date of interview).

Table 14. Percentage of perception of maximum legal penalty in your state for first offense possession of an ounce or less of marijuana for your own use among all persons aged 212, by age group — National Survey on Drug Use and Health, Colorado, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Fine												
Total	32.8 (28.8-37.0)	32.9 (27.7-38.5)	38.8 (34.1-43.7)	42.9 (38.7-47.2)	46.3 (41.5-51.1)	49.6 (44.8-54.4)	47.8 (43.2-52.5)	46.8 (42.2-51.5)	46.5 (41.5-51.5)	47.0 (43.3-50.7)	39.8 (34.8-44.7)	26.3 (22.4-30.5)
12-17	17.1 (13.1-22.0)	11.8 (9.2-15.1)	11.1 (8.3-14.6)	20.0 (16.6-23.8)	27.5 (23.3-32.1)	27.2 (23.7-30.9)	26.4 (22.8-30.5)	26.1 (21.9-30.7)	28.0 (23.3-33.2)	28.7 (23.6-34.5)	25.4 (20.5-30.9)	26.0 (20.9-31.8)
18-25	30.4 (25.5-35.8)	26.5 (22.4-31.0)	27.4 (22.7-32.8)	35.0 (29.9-40.5)	42.4 (35.4-49.6)	46.0 (39.9-52.2)	44.9 (39.8-50.1)	44.3 (38.1-50.6)	47.1 (40.8-53.5)	49.6 (43.7-55.4)	47.4 (41.8-53.1)	37.5 (32.1-43.4)
≥26	35.9 (30.9-41.2)	37.9 (30.8-45.6)	46.1 (40.0-52.4)	48.1 (42.7-53.6)	49.9 (43.3-56.5)	53.7 (46.9-60.4)	51.5 (45.0-57.9)	50.1 (44.2-56.0)	48.7 (42.4-55.1)	48.9 (44.1-53.7)	39.8 (33.4-46.7)	24.1 (19.3-29.6)
Probation												
Total	16.6 (13.7-20.0)	13.5 (10.7-16.9)	10.5 (8.5-13.0)	10.1 (8.0-12.7)	10.1 (7.9-12.9)	9.5 (6.9-12.9)	8.6 (6.5-11.4)	7.6 (5.8-10.0)	11.6 (8.8-15.2)	12.9 (10.8-15.7)	10.0 (8.2-12.1)	7.7 (6.2-9.5)
12-17	15.0 (11.3-19.7)	14.3 (11.3-18.0)	16.3 (13.1-20.2)	16.2 (12.6-20.5)	17.8 (14.8-21.4)	19.3 (15.4-24.0)	15.2 (11.8-19.3)	14.0 (10.5-18.3)	19.2 (14.3-25.3)	21.9 (17.3-27.3)	21.3 (16.0-27.7)	19.5 (14.6-25.6)
18-25	16.8 (13.2-21.2)	16.9 (12.4-22.5)	16.2 (12.3-21.0)	14.4 (11.3-18.3)	14.1 (10.8-18.2)	14.2 (10.9-18.2)	13.6 (10.4-17.6)	12.8 (10.1-16.1)	13.6 (10.7-17.2)	16.0 (12.3-20.7)	13.6 (10.3-17.8)	9.6 (7.0-12.9)
≥26 years	16.9 (13.1-21.5)	12.6 (9.1-17.2)	8.2 (5.7-11.8)	8.2 (5.6-11.9)	8.1 (5.3-12.2)	7.0 (3.9-12.4)	6.6 (3.9-10.9)	5.7 (3.6-8.9)	10.2 (6.8-15.0)	11.1 (8.3-14.7)	7.8 (5.7-10.5)	5.8 (4.1-8.3)
Community Service												
Total	10.0 (7.9-12.6)	12.2 (9.8-15.2)	10.9 (8.5-13.8)	9.9 (7.8-12.9)	8.6 (6.5-11.3)	7.1 (4.7-10.4)	9.4 (6.7-13.0)	8.3 (6.1-11.3)	7.1 (4.9-10.3)	8.4 (6.0-11.8)	7.0 (4.8-10.0)	4.3 (3.0-5.9)
12-17	21.7 (18.1-25.8)	22.7 (18.5-27.6)	24.4 (19.5-30.2)	19.3 (14.9-24.7)	12.1 (9.1-16.0)	11.6 (8.6-15.1)	13.6 (10.0-18.2)	13.3 (9.4-18.3)	12.3 (8.9-16.8)	13.0 (9.0-18.4)	13.7 (9.6-19.2)	13.3 (9.9-17.6)
18-25	10.8 (7.7-14.9)	14.5 (11.2-18.5)	13.1 (10.0-16.8)	12.5 (9.1-17.1)	10.1 (7.1-14.2)	8.9 (6.0-13.2)	10.9 (7.9-15.0)	9.5 (6.5-13.6)	7.2 (4.2-11.9)	6.9 (4.7-10.1)	7.1 (4.7-10.8)	6.8 (4.7-9.6)
≥26	8.0 (5.2-11.9)	9.9 (6.9-14.0)	8.1 (5.4-12.0)	7.9 (5.1-12.1)	7.7 (5.1-11.7)	6.0 (3.3-10.5)	8.5 (5.3-13.5)	7.4 (4.9-11.2)	6.5 (4.0-10.4)	8.2 (5.1-12.9)	6.0 (3.4-10.5)	2.6 (1.4-4.9)
Possible Prison Sentence												
Total	33.6 (29.8-37.6)	34.5 (30.2-39.0)	34.0 (29.6-38.6)	30.0 (26.4-33.7)	26.7 (22.8-31.0)	24.0 (20.5-27.9)	21.2 (17.9-25.0)	20.0 (16.6-24.0)	16.7 (13.3-20.8)	16.6 (14.0-19.6)	13.2 (10.3-16.8)	6.7 (5.0-8.9)

12-17	36.8 (32.2-41.6)	39.1 (33.7-44.7)	35.4 (30.0-41.2)	31.6 (27.1-36.5)	26.0 (22.1-30.4)	23.5 (19.7-27.7)	25.9 (22.1-30.2)	29.4 (24.2-35.1)	26.8 (22.2-31.9)	24.6 (20.6-29.1)	24.5 (19.3-30.7)	20.0 (15.9-24.8)
18-25	33.2 (28.4-38.2)	34.7 (30.3-39.3)	36.5 (31.4-41.9)	30.7 (26.3-35.5)	28.0 (23.2-33.4)	23.6 (18.7-29.4)	20.3 (15.8-25.6)	21.7 (15.7-29.1)	19.0 (14.2-24.9)	16.3 (12.9-20.4)	10.7 (7.8-14.6)	8.0 (5.7-11.0)
≥26	33.1 (28.0-38.7)	33.6 (27.8-40.0)	33.1 (27.5-39.3)	29.5 (25.0-34.5)	26.5 (21.2-32.5)	24.2 (19.5-29.6)	20.8 (16.5-25.9)	18.4 (14.3-23.4)	15.0 (11.1-20.0)	15.6 (12.4-19.5)	12.2 (8.6-17.2)	4.8 (2.9-7.7)
Mandatory Prison Sentence												
Total	7.0 (5.0-9.6)	7.0 (5.1-9.4)	5.8 (4.0-8.5)	3.6 (2.3-5.6)	2.7 (1.8-4.1)	3.9 (2.7-5.5)	5.6 (4.1-7.7)	6.1 (4.2-8.7)	4.5 (2.7-7.4)	3.4 (2.1-5.5)	1.9 (1.1-3.2)	0.7 (0.5-1.2)
12-17	9.4 (6.7-13.1)	12.1 (8.7-16.5)	12.8 (9.2-17.5)	10.3 (7.4-14.2)	10.0 (6.8-14.3)	11.2 (8.4-14.8)	11.3 (8.8-14.3)	9.9 (7.4-13.2)	7.8 (5.3-11.2)	6.1 (4.0-9.3)	5.1 (3.0-8.6)	4.4 (2.5-7.7)
18-25	8.8 (5.6-13.7)	7.4 (4.5-12.0)	6.8 (4.4-10.2)	5.6 (3.5-8.9)	2.3 (1.2-4.7)	2.4 (1.2-4.7)	4.0 (2.5-6.3)	3.9 (2.3-6.4)	2.6 (1.3-4.9)	1.8 (0.9-3.5)	1.4 (0.6-3.2)	0.9 (0.3-2.6)
≥26	6.2 (3.6-10.4)	6.0 (3.6-9.8)	4.4 (2.3-8.5)	2.2 (0.8-5.6)	1.7 (0.8-3.7)	3.1 (1.8-5.4)	5.2 (3.3-8.1)	6.0 (3.6-9.9)	4.5 (2.3-8.6)	3.3 (1.7-6.3)	1.5 (0.6-3.6)	0.3 (0.1-0.8)
No penalty												
Total	DNA	DNA	DNA	3.5 (2.2-5.5)	5.6 (3.8-9.0)	6.0 (3.7-9.4)	7.3 (4.7-11.0)	11.1 (7.9-15.4)	13.5 (10.7-17.0)	11.6 (9.1-14.8)	28.4 (24.4-32.6)	54.3 (50.4-58.3)
12-17	DNA	DNA	DNA	2.6 (1.4-4.9)	6.6 (4.6-9.3)	7.3 (5.1-10.1)	7.5 (5.2-10.7)	7.4 (4.5-12.0)	5.9 (3.8-8.9)	5.6 (3.7-8.4)	10.1 (7.1-14.1)	16.7 (12.9-21.5)
18-25	DNA	DNA	DNA	1.7 (0.8-3.7)	3.1 (1.7-5.5)	4.8 (2.8-8.2)	6.3 (3.9-10.1)	7.9 (5.9-10.5)	10.5 (6.9-15.8)	9.4 (5.9-14.6)	19.7 (15.6-24.6)	37.3 (32.0-42.8)
≥26	DNA	DNA	DNA	4.1 (2.4-6.9)	6.0 (3.8-9.3)	6.0 (3.2-11.0)	7.4 (4.1-13.0)	12.3 (8.1-18.3)	15.1 (11.3-19.9)	12.9 (9.6-17.1)	32.5 (27.5-38.0)	62.4 (57.3-67.3)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.
Abbreviations: CI = confidence interval; DNA = data not available.
Perception of legal penalty for marijuana possession is defined as those who reported their perception of the maximum legal penalty in their state of residence for first offense possession of an ounce or less of marijuana for their own use.

Supplementary Material

Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Suggested Citation: Alejandro Azofeifa, Margaret E. Mattson, and Rob Lyster. Supplementary Material. State Level Data: Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, District of Columbia, 2002–2014. Center for Behavioral Health Statistics and Quality. (2016). Substance Abuse and Mental Health Services Administration, Rockville, MD.

Notes: A) Two years of data were combined to allow for reliable state-level prevalence estimates. Some state-level prevalence estimates for combined years have been suppressed (indicated by an asterisk (*) symbol in each cell) based on the following National Survey on Drug Use and Health (NSDUH) suppression criteria. Prevalence estimates are suppressed if any of the following occurred: 1) the prevalence estimate is $<0.005^*$ or $>99.995^*$; 2) the relative standard error (RSE) of the negative natural logarithm of the estimated proportion p (where p is the prevalence divided by 100) is >0.175 if the prevalence is $>50^*$; 3) the RSE of the negative natural logarithm of $(1-p)$ is >0.175 if the prevalence is $>50^*$; 4) the actual sample size is <100 ; or 5) the effective sample size (defined as the actual sample size divided by the design effect) is <68 . B) The estimates provided in this supplementary material for each state and the District of Columbia are from design-based direct estimation procedures (i.e., they are based directly on the respondent sample and the accompanying analytical weights with variances derived through a Taylor-linearization approach using SUDAAN software) as used for the national level estimates (available at: Azofeifa A, Mattson ME, Schauer G, McAfee T, Grant A, Lyster R. National Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, United States, 2002–2014. MMWR Surveill Summ 2016;65(No. SS-11):1–25. DOI: <http://dx.doi.org/10.15585/mmwr.ss6511a1>). Caution is advised when comparing data from other NSDUH reports that are based on different design-based estimation procedures. C) Interpretation of state-level trend data should be done with caution as some prevalence estimates are suppressed based on the above criteria and survey cycle differences may not be statistically significant.

BILAG 3

Table 1. Percentage of past month marijuana use* among all persons aged ≥ 12, by selected demographic characteristics — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Characteristics Age group [yrs]	2002–2003	2003–2004	2004–2005	2005–2006	2006–2007	2007–2008	2008–2009	2009–2010	2010–2011	2011–2012	2012–2013	2013–2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	9.5 (7.8–11.5)	6.6 (5.2–8.2)	6.5 (4.8–8.7)	8.1 (6.3–10.2)	9.6 (7.6–12.0)	9.5 (7.7–11.7)	8.5 (6.3–11.4)	10.1 (7.7–13.3)	11.0 (8.7–13.8)	11.2 (9.0–13.8)	12.3 (10.1–14.9)	12.6 (10.7–14.7)
12–17	7.8 (5.9–10.2)	5.9 (4.1–8.5)	7.8 (5.5–10.9)	7.3 (5.3–9.9)	7.4 (5.2–10.4)	9.3 (7.1–12.0)	9.7 (6.6–14.0)	11.8 (8.3–16.5)	12.9 (9.9–16.6)	9.8 (7.6–12.5)	10.7 (8.6–13.3)	12.7 (10.1–15.9)
18–25	24.7 (20.6–29.4)	19.4 (15.2–24.5)	17.6 (13.7–22.4)	19.0 (14.0–25.2)	23.6 (18.9–29.1)	22.6 (18.1–27.7)	17.8 (14.3–21.8)	18.7 (15.7–22.1)	20.8 (17.6–24.5)	23.8 (19.6–28.7)	24.2 (19.8–29.1)	22.7 (18.7–27.3)
≥26	6.7 (4.8–9.1)	4.2 (3.0–5.8)	4.3 (2.7–7.0)	6.0 (4.2–8.5)	6.7 (4.7–9.5)	6.6 (4.8–9.2)	6.3 (4.0–10.0)	8.1 (5.3–12.2)	8.6 (6.2–11.9)	8.3 (6.1–11.3)	9.7 (7.3–12.8)	10.4 (8.3–12.9)
26–34	14.3 (9.5–21.2)	6.3 (3.6–10.7)	8.8 (5.2–14.4)	14.4 (10.4–19.6)	16.9 (10.6–25.9)	15.8 (9.9–24.2)	12.6 (8.4–18.3)	16.3 (11.3–23.0)	16.2 (11.7–22.2)	13.4 (9.4–18.6)	15.5 (11.0–21.5)	12.4 (8.5–17.7)
35–44	6.6 (3.7–11.7)	4.0 (1.9–8.4)	4.1 (2.0–8.3)	4.8 (2.3–9.8)	7.1 (3.3–14.5)	6.1 (2.7–13.1)	1.5 (0.5–4.9)	5.8 (3.3–9.9)	11.0 (5.6–20.3)	11.2 (7.0–16.6)	12.7 (7.0–21.9)	9.5 (6.0–14.8)
45–54	7.8 (3.5–16.3)	8.5 (4.4–15.9)	2.1 (0.6–7.4)	4.2 (2.3–9.8)	4.2 (1.4–11.4)	4.7 (2.1–10.2)	4.7 (1.4–11.4)	4.7 (1.4–11.4)	4.6 (1.7–11.6)	4.6 (1.7–11.6)	4.6 (1.7–11.6)	20.5 (13.0–30.8)
55–64	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
≥65	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Sex												
Male	10.7 (8.1–14.0)	7.3 (5.2–10.0)	8.9 (5.8–13.4)	11.9 (8.3–16.7)	13.2 (9.6–17.9)	11.8 (8.9–15.5)	11.0 (7.3–16.2)	14.6 (10.3–20.2)	14.7 (10.7–19.8)	14.3 (10.8–18.6)	14.3 (10.7–18.7)	14.7 (11.8–18.2)
Female	8.5 (6.4–11.1)	6.0 (4.1–8.6)	4.4 (3.1–6.1)	4.8 (3.6–6.3)	6.4 (4.8–8.6)	7.6 (5.8–9.9)	6.4 (4.6–8.8)	6.3 (4.2–9.4)	7.8 (5.7–10.7)	8.5 (6.5–11.1)	10.5 (8.6–12.8)	10.7 (8.6–13.3)
Race/ethnicity												
Non-Hispanic white	10.6 (7.3–15.3)	8.6 (6.0–12.2)	8.0 (4.7–13.4)	9.1 (5.7–14.1)	8.9 (6.0–13.1)	8.9 (6.4–12.2)	8.0 (4.8–13.1)	8.3 (5.0–13.6)	9.1 (6.4–12.4)	10.5 (7.4–14.7)	10.6 (7.4–15.1)	11.6 (8.8–15.3)
Non-Hispanic black	9.4 (7.4–11.8)	6.2 (4.6–8.2)	6.0 (4.3–8.4)	7.8 (5.8–10.4)	9.8 (7.1–13.2)	9.7 (7.1–13.1)	9.1 (5.9–13.7)	11.2 (7.4–16.6)	11.3 (8.3–15.2)	10.0 (7.8–12.7)	12.5 (9.7–16.1)	14.1 (11.1–17.7)
Non-Hispanic American Indian/Alaska Native	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic Hawaiian/Other Pacific Islander	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic Asian	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic two or more races	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Hispanic	(*)	(*)	2.4 (0.9–6.0)	4.0 (1.5–10.5)	(*)	(*)	(*)	11.2 (5.7–20.8)	(*)	(*)	(*)	7.0 (3.1–15.0)
Current employment status	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)

Full-time	10.9 (7.9-14.9)	5.4 (3.6-8.2)	3.9 (2.5-5.9)	7.5 (5.6-9.9)	8.0 (5.9-10.8)	7.0 (4.9-9.8)	6.8 (4.4-10.4)	8.1 (5.4-12.0)	10.9 (8.2-14.4)	12.2 (9.0-16.2)	11.4 (8.6-14.9)	11.2 (9.1-13.8)
Part-time	9.5 (6.5-13.6)	12.2 (8.0-18.1)	(*)	(*)	12.3 (7.1-20.5)	12.4 (7.3-20.3)	(*)	(*)	11.2 (6.5-18.8)	11.2 (7.2-17.1)	16.2 (10.3-24.5)	18.1 (11.0-28.3)
Unemployed	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	20.1 (12.4-31.0)	20.7 (13.1-31.0)
Other ²	5.0 (3.3-7.5)	4.8 (2.6-8.7)	6.7 (3.8-11.5)	7.5 (4.3-12.8)	12.5 (8.0-19.2)	13.0 (8.6-19.0)	6.9 (4.6-10.1)	5.6 (3.4-9.0)	5.4 (3.6-8.0)	7.2 (4.9-10.6)	10.4 (6.7-15.9)	11.7 (7.8-17.2)
Highest level of education completed ¹												
< High school	7.6 (4.4-12.7)	4.5 (2.4-8.3)	6.9 (4.0-11.7)	9.0 (5.2-15.1)	11.8 (6.7-20.0)	10.8 (6.0-18.9)	6.5 (3.6-11.6)	11.0 (6.3-18.7)	(*)	(*)	12.3 (7.1-20.5)	18.4 (12.4-26.6)
High school graduate	11.9 (8.0-17.1)	7.9 (5.1-11.9)	5.7 (3.3-9.8)	7.6 (5.0-11.3)	12.4 (8.0-18.8)	15.1 (10.5-21.2)	9.4 (6.9-12.8)	6.6 (3.9-11.0)	8.6 (4.4-16.1)	14.2 (8.9-22.0)	16.7 (11.1-24.4)	13.9 (8.7-21.5)
Some college	10.7 (7.0-16.0)	8.0 (4.9-13.0)	7.6 (5.1-11.3)	9.9 (5.6-16.9)	14.5 (8.8-22.9)	11.5 (6.6-19.2)	(*)	(*)	20.3 (13.2-30.1)	15.1 (10.8-20.7)	13.0 (9.1-18.3)	12.7 (8.7-18.1)
College graduate	8.8 (5.7-13.4)	5.9 (3.4-10.1)	5.9 (3.0-11.2)	7.4 (4.5-11.9)	5.9 (3.7-9.1)	5.8 (3.8-8.6)	7.4 (4.6-11.6)	7.9 (5.0-12.4)	7.2 (5.5-9.5)	8.8 (6.3-12.2)	10.6 (7.7-14.5)	10.8 (8.6-13.6)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Abbreviation: CI = confidence interval.

* Past month use of marijuana is defined as those who reported use of marijuana within 30 days preceding the date of interview.

¹ Current employment and highest level of education completed are only defined for 18 or older.

² "Other" employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Table 2. Percentage of past year marijuana use among all persons aged ≥12, by age group — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	15.3 (12.9-18.0)	14.4 (12.1-17.0)	15.2 (12.9-17.8)	14.8 (12.7-17.1)	15.9 (13.5-18.6)	15.9 (13.7-18.3)	14.4 (11.7-17.6)	16.3 (13.2-19.9)	18.7 (15.4-22.5)	20.3 (17.3-23.7)	20.8 (17.7-24.4)	21.4 (18.9-24.1)
12-17	15.1 (12.3-18.4)	12.1 (9.7-15.1)	14.5 (11.3-18.4)	13.5 (10.7-17.0)	12.9 (10.1-16.2)	14.6 (11.9-17.8)	16.8 (13.0-21.6)	19.0 (14.9-24.0)	18.8 (15.1-23.1)	17.8 (15.1-20.9)	19.5 (16.2-23.4)	21.7 (18.1-25.7)
18-25	37.2 (32.7-41.9)	30.3 (25.4-35.8)	32.8 (28.0-38.1)	37.5 (32.8-42.6)	39.0 (33.6-44.6)	37.0 (32.0-42.4)	32.8 (28.6-37.2)	33.5 (29.9-37.4)	37.5 (33.0-42.2)	42.8 (37.6-48.2)	42.6 (37.1-48.3)	38.9 (34.1-44.0)
≥26	11.0 (8.3-14.3)	11.5 (9.2-14.4)	12.1 (9.6-15.1)	10.5 (8.5-13.0)	11.2 (8.6-14.3)	11.3 (9.0-14.1)	10.0 (7.2-13.8)	12.2 (8.9-16.4)	14.4 (10.8-19.0)	15.2 (11.9-19.2)	16.1 (12.6-20.2)	17.6 (14.6-20.9)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 Past year use of marijuana is defined as those who reported use of marijuana within 12 months preceding the date of interview. —these also includes those who reported use of marijuana within 30 days preceding the date of interview.

Table 3. Percentage of daily or almost daily marijuana use in the past year and past month among all persons aged ≥12, by age group — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Age group (yrs)	2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014	
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Daily or almost daily marijuana use in the past year¹																								
Total	1.6 (1.1-2.4)	0.5 (0.3-0.8)	1.0 (0.6-1.7)	1.1 (0.7-1.8)	1.0 (0.6-1.8)	1.0 (0.6-1.8)	1.7 (1.1-2.7)	1.4 (0.9-2.4)	1.6 (0.9-2.8)	2.2 (1.4-3.3)	1.6 (0.9-2.6)	2.1 (1.4-3.2)	3.0 (2.0-4.3)											
12-17	1.0 (0.5-2.1)	1.2 (0.5-2.9)	1.3 (0.5-3.2)	1.3 (0.7-2.4)	1.7 (1.0-2.9)	1.5 (0.7-3.3)	1.1 (0.3-4.0)	2.2 (1.2-3.9)	2.2 (1.2-4.1)	0.9 (0.4-2.3)	1.5 (0.7-3.0)	2.1 (0.9-5.1)												
18-25	5.2 (3.2-8.3)	2.1 (1.0-4.3)	3.4 (1.8-6.2)	3.6 (2.0-6.6)	3.1 (1.8-5.3)	4.7 (3.2-6.8)	3.1 (2.0-4.8)	2.6 (1.3-5.1)	4.8 (3.0-7.5)	5.3 (4.1-8.4)	5.9 (4.1-8.4)	6.0 (4.0-8.7)												
≥26	1.0 (0.5-2.0)	0.1 (0.0-0.5)	0.6 (0.2-1.5)	0.6 (0.3-1.6)	0.5 (0.1-1.8)	1.1 (0.5-2.4)	1.1 (0.5-2.4)	1.4 (0.6-3.0)	1.6 (0.8-2.9)	0.7 (0.2-2.3)	1.3 (0.6-2.8)	2.4 (1.4-4.0)												
Daily or almost daily marijuana use in the past month²																								
Total	2.4 (1.8-3.4)	1.2 (0.8-1.9)	1.9 (1.2-2.8)	2.3 (1.6-3.3)	2.1 (1.4-3.1)	2.0 (1.3-3.1)	1.6 (1.0-2.6)	1.9 (1.3-2.9)	2.4 (1.6-3.5)	2.3 (1.5-3.4)	3.4 (2.5-4.7)	4.4 (3.3-5.9)												
12-17	1.4 (0.6-3.3)	1.6 (0.8-3.1)	1.9 (1.0-3.7)	1.7 (1.0-2.8)	2.0 (1.2-3.3)	1.8 (1.0-3.1)	1.7 (0.8-3.6)	3.3 (1.9-5.8)	2.9 (1.7-5.0)	1.3 (0.6-2.6)	1.3 (0.5-2.9)	1.6 (0.5-5.1)												
18-25	7.5 (5.3-10.3)	5.2 (3.5-7.8)	5.4 (3.6-8.1)	5.4 (3.3-8.8)	5.5 (3.7-8.0)	4.5 (3.0-6.7)	2.6 (1.5-4.5)	3.4 (2.0-5.6)	5.1 (3.3-7.8)	7.3 (5.3-9.8)	8.0 (6.0-10.7)	7.0 (4.8-10.0)												
≥26	1.5 (0.8-2.9)	0.4 (0.1-1.4)	1.2 (0.6-2.5)	1.7 (1.0-3.1)	1.3 (0.6-2.8)	1.4 (0.7-3.0)	1.4 (0.7-2.7)	1.5 (0.8-2.8)	1.7 (0.9-3.0)	1.2 (0.5-2.8)	2.6 (1.6-4.0)	4.1 (2.9-5.8)												

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Past year daily or almost daily use is defined as having used marijuana on 300 or more days in the past year.

² Past month daily or almost daily use is defined as having used marijuana on 20 or more days in the past month.

Table 4. Percentage of daily or almost daily marijuana use in the past year and past month among marijuana users aged ≥12, by age group — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Age group (yrs)	2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014	
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Daily or almost daily marijuana use in the past year*																								
Total	10.6 (7.0-15.8)	3.4 (2.0-5.9)	6.8 (4.1-11.2)	7.7 (4.9-11.8)	6.4 (3.7-10.7)	11.0 (7.1-16.6)	10.0 (6.1-15.9)	10.1 (5.9-17.2)	11.7 (7.7-17.3)	7.7 (4.6-12.5)	10.1 (6.6-15.0)	13.8 (9.7-19.3)												
12-17	(*)	(*)	(*)	(*)	(*)	(*)	(*)	11.6 (6.8-19.2)	11.8 (6.9-19.9)	5.2 (2.2-12.0)	7.4 (3.7-14.5)	(*)												
18-25	14.0 (8.7-21.6)	6.9 (3.3-13.8)	10.2 (5.6-18.0)	9.6 (5.4-16.4)	8.0 (4.8-13.2)	12.6 (8.7-17.8)	9.5 (6.1-14.6)	7.7 (3.8-14.9)	12.8 (8.2-19.5)	12.3 (8.4-17.6)	13.8 (9.7-19.3)	15.3 (10.4-21.9)												
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	8.1 (3.7-17.0)	13.5 (8.0-21.9)												
Daily or almost daily marijuana use in the past month†																								
Total	25.7 (18.8-34.0)	18.9 (13.1-26.5)	28.7 (21.2-37.6)	28.5 (21.1-37.1)	21.7 (15.2-30.1)	20.8 (13.9-30.1)	18.9 (12.1-28.4)	18.9 (12.4-27.8)	21.4 (15.2-29.3)	20.5 (14.0-28.9)	28.1 (21.8-35.6)	35.3 (28.4-42.9)												
12-17	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)												
18-25	30.1 (22.5-39.0)	26.9 (19.7-35.6)	30.6 (21.5-41.6)	19.9 (14.0-27.6)	23.3 (16.2-32.2)	19.9 (14.0-27.6)	14.7 (8.9-23.5)	18.1 (11.2-27.8)	24.6 (16.7-34.6)	30.5 (24.7-37.0)	33.2 (26.1-41.3)	30.8 (22.5-40.5)												
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)												

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Analysis limited to marijuana users (i.e., estimate of daily or almost daily use in the past year is among past year marijuana users; estimates of daily or almost daily use in the past month is among past month marijuana users).
 † Past year daily or almost daily use is defined as having used marijuana on 300 or more days in the past year.
 ‡ Past month daily or almost daily use is defined as having used marijuana on 20 or more days in the past month.

Table 5. Percentage of past year marijuana initiation among persons aged 212 years at risk for initiation, and mean age at first use of marijuana among past year marijuana initiates, by age group — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Past Year marijuana initiation among persons at risk for initiation ¹												
Total	2.5 (1.5-3.9)	1.6 (1.1-2.5)	1.7 (1.1-2.6)	1.9 (1.1-3.2)	1.6 (1.0-2.5)	2.0 (1.3-3.0)	2.3 (1.6-3.2)	2.1 (1.3-3.2)	2.9 (1.5-5.7)	3.9 (2.2-7.0)	3.4 (2.2-5.3)	2.5 (1.6-3.9)
12-17	7.8 (5.5-11.0)	5.4 (4.1-7.2)	5.7 (3.9-8.2)	5.6 (3.8-8.1)	5.3 (3.7-7.4)	6.7 (4.6-9.5)	7.4 (5.2-10.6)	7.4 (5.0-10.8)	6.2 (3.9-9.8)	5.5 (3.8-7.8)	7.8 (5.7-10.6)	10.2 (7.1-14.4)
18-25	6.3 (3.7-10.6)	5.8 (3.0-11.0)	7.6 (4.2-13.6)	8.8 (4.6-16.1)	5.9 (2.8-12.0)	6.1 (3.7-9.9)	7.8 (5.5-11.0)	8.2 (5.1-13.1)	8.7 (5.3-14.1)	9.9 (6.1-15.5)	9.1 (5.3-15.2)	6.3 (3.2-11.9)
≥26	1.0 (0.3-3.7)	0.3 (0.0-2.0)	0.3 (0.0-1.9)	0.2 (0.0-1.9)	0.3 (0.0-1.9)	0.3 (0.0-1.9)	0.2 (0.0-1.9)	0.2 (0.0-1.9)	0.2 (0.0-1.9)	2.3 (0.7-7.8)	1.5 (0.5-4.3)	0.2 (0.0-1.9)
Estimated no. (in 1,000s) of past year marijuana initiation among those at risk for initiation ²												
Total	6 (4-10)	4 (3-6)	4 (3-6)	4 (3-7)	4 (2-6)	5 (3-8)	6 (4-9)	5 (4-8)	7 (4-15)	10 (5-17)	8 (5-13)	6 (4-9)
12-17	2 (2-3)	2 (1-2)	2 (1-2)	2 (1-3)	2 (1-2)	2 (1-3)	2 (2-3)	2 (1-3)	2 (1-3)	1 (1-2)	2 (1-3)	3 (2-4)
18-25	2 (1-3)	2 (1-4)	2 (1-4)	3 (1-5)	2 (1-4)	2 (1-4)	3 (2-5)	3 (2-5)	3 (2-5)	4 (3-6)	4 (2-6)	2 (1-5)
≥26	2 (1-7)	1 (0-4)	0 (0-3)	0 (0-3)	0 (0-3)	0 (0-3)	0 (0-4)	0 (0-4)	0 (0-4)	4 (1-14)	3 (1-8)	0 (0-3)
Mean age at first use among past year marijuana initiates												
Total	19.9 (17.3-22.4)	18.7 (16.4-20.9)	18.4 (16.8-20.0)	18.4 (16.7-20.0)	17.3 (16.1-18.6)	18.9 (15.8-21.9)	19.3 (16.7-21.9)	18.0 (17.1-18.9)	22.9 (16.4-29.4)	26.3 (20.0-32.7)	24.8 (17.8-31.9)	19.5 (15.2-23.9)
12-17	14.8 (14.1-15.5)	14.9 (13.9-15.8)	14.9 (14.2-15.5)	15.0 (14.4-15.7)	15.1 (14.6-15.7)	15.2 (14.7-15.6)	15.0 (14.5-15.5)	14.8 (14.4-15.2)	15.0 (14.6-15.4)	14.9 (14.6-15.2)	14.7 (14.3-15.1)	14.8 (14.4-15.2)
18-25	19.4 (18.1-20.7)	19.5 (18.4-20.7)	20.8 (19.4-22.2)	20.5 (19.1-21.9)	19.1 (18.0-20.3)	19.0 (17.8-20.3)	20.1 (18.9-21.2)	20.2 (19.0-21.3)	19.8 (18.9-20.8)	20.2 (19.3-21.2)	20.1 (19.2-21.1)	19.5 (18.5-20.3)
≥26	0.2 (0.0-0.4)	0.2 (0.0-0.4)	0.2 (0.0-0.4)	0.2 (0.0-0.4)	0.2 (0.0-0.4)	0.2 (0.0-0.4)	0.2 (0.0-0.4)	0.2 (0.0-0.4)	0.2 (0.0-0.4)	0.2 (0.0-0.4)	0.2 (0.0-0.4)	0.2 (0.0-0.4)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹Past year marijuana initiation (initiates), among persons at risk for initiation is defined as those who did not use marijuana in their lifetime or used marijuana for the first time within 12 months.

Table 6. Percentage of perceived great risk and no risk from smoking marijuana once a month and once or twice a week among all persons aged ≥12, by age group — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived great risk from smoking marijuana once a month												
Total	32.1 (27.8-36.6)	32.1 (28.0-36.4)	31.7 (27.4-36.3)	30.7 (26.3-35.4)	29.2 (25.7-33.0)	29.7 (25.1-34.6)	28.3 (24.3-32.6)	24.8 (20.8-29.2)	23.5 (20.2-27.2)	22.9 (19.4-26.9)	20.3 (17.6-23.4)	16.2 (13.8-18.9)
12-17	31.2 (26.9-35.9)	33.2 (28.6-38.0)	30.9 (26.8-35.3)	31.8 (28.0-35.8)	30.2 (26.3-34.4)	27.9 (23.4-33.0)	27.8 (23.5-32.4)	26.3 (21.8-31.4)	24.2 (20.0-28.9)	23.7 (19.9-28.0)	24.4 (20.8-28.5)	17.5 (14.2-21.4)
18-25	18.7 (14.8-23.3)	18.6 (14.9-23.2)	19.4 (16.0-23.4)	16.2 (12.4-20.7)	15.7 (12.2-19.9)	17.2 (13.8-21.2)	16.7 (13.3-20.7)	14.8 (11.8-18.5)	11.8 (9.3-15.0)	12.1 (9.6-15.2)	11.7 (9.4-14.4)	9.9 (6.9-14.1)
≥26	34.9 (29.7-40.4)	34.6 (29.7-39.9)	34.1 (28.8-39.7)	33.4 (28.0-39.3)	32.1 (27.7-36.7)	32.6 (28.8-39.0)	30.9 (25.8-36.6)	26.9 (21.9-32.5)	26.1 (22.0-30.7)	25.4 (20.8-30.6)	22.0 (18.4-26.0)	17.5 (14.6-20.8)
Perceived no risk from smoking marijuana once a month												
Total	13.9 (10.7-17.7)	15.0 (11.8-18.8)	14.8 (12.7-17.2)	14.6 (12.4-17.1)	16.1 (13.3-19.3)	17.3 (15.0-19.9)	15.9 (13.2-18.9)	18.3 (15.6-21.3)	20.4 (17.4-23.9)	19.6 (16.2-23.4)	23.2 (20.0-26.7)	28.4 (25.2-31.8)
12-17	11.4 (8.8-14.7)	8.8 (6.4-12.0)	10.4 (7.8-13.8)	11.3 (8.8-14.6)	10.9 (8.5-13.7)	14.1 (10.8-18.2)	14.6 (11.1-19.1)	12.5 (9.6-16.1)	15.6 (11.9-20.1)	17.1 (13.2-22.0)	16.8 (13.5-20.8)	21.4 (17.1-26.3)
18-25	20.5 (16.8-24.9)	19.4 (15.7-23.9)	22.3 (18.6-26.1)	22.8 (19.2-26.9)	24.3 (20.3-28.9)	24.5 (20.6-28.8)	22.1 (18.8-25.8)	23.5 (20.5-26.8)	26.9 (23.2-31.0)	31.1 (27.4-35.1)	37.3 (32.9-41.8)	41.0 (35.9-46.3)
≥26	12.7 (9.0-17.8)	14.7 (10.8-19.5)	13.9 (11.4-16.9)	13.3 (10.7-16.5)	14.8 (11.5-18.7)	16.0 (13.1-19.4)	14.6 (11.4-18.4)	17.6 (14.1-21.8)	19.3 (15.6-23.7)	17.0 (13.1-21.9)	20.5 (16.7-24.9)	26.1 (22.3-30.4)
Perceived great risk from smoking marijuana once or twice a week												
Total	40.0 (35.0-46.2)	41.7 (37.1-46.5)	40.7 (36.5-45.0)	40.4 (36.3-44.6)	38.1 (34.2-42.2)	34.3 (29.9-39.1)	33.4 (29.1-37.9)	32.0 (27.9-36.4)	31.5 (27.1-36.3)	29.8 (25.8-34.1)	26.1 (23.3-29.1)	20.5 (17.6-23.8)
12-17	42.7 (37.4-48.2)	45.2 (40.1-50.5)	43.2 (38.4-48.1)	41.9 (36.1-48.0)	42.4 (37.1-47.7)	43.8 (37.4-50.3)	40.6 (35.9-45.6)	36.0 (31.3-41.0)	33.8 (28.5-39.6)	30.3 (24.9-36.4)	30.6 (26.4-35.2)	26.5 (22.2-31.4)
18-25	27.3 (22.2-33.0)	27.4 (22.0-33.5)	25.5 (21.2-30.4)	24.9 (19.2-31.5)	24.8 (20.0-30.3)	24.1 (20.1-28.7)	23.2 (18.9-28.3)	21.8 (18.3-25.9)	18.9 (14.9-23.7)	17.5 (14.2-21.3)	14.7 (12.1-17.8)	11.3 (8.1-15.5)
≥26	42.3 (36.2-48.7)	44.3 (38.9-49.8)	43.2 (38.1-48.5)	43.2 (38.3-48.3)	40.6 (35.7-45.7)	35.7 (30.0-41.8)	35.0 (29.8-40.6)	34.0 (28.9-39.4)	34.2 (28.7-40.2)	32.6 (27.6-38.0)	28.4 (25.0-31.9)	22.1 (18.6-26.0)
Perceived no risk from smoking marijuana once or twice a week												
Total	6.9 (5.5-8.6)	6.7 (5.3-8.6)	7.0 (5.5-8.8)	8.1 (6.6-10.0)	9.4 (7.4-11.8)	8.9 (7.1-11.1)	7.7 (6.1-9.7)	10.4 (8.2-13.0)	11.8 (9.4-14.7)	12.0 (9.8-14.6)	16.3 (13.4-19.7)	19.9 (17.2-22.8)
12-17	6.7 (4.8-9.2)	5.3 (3.5-7.8)	7.3 (4.8-11.1)	8.6 (6.0-12.1)	8.5 (5.8-12.1)	7.8 (5.4-10.9)	8.4 (5.7-12.1)	9.9 (7.1-13.6)	10.9 (8.0-14.6)	12.5 (9.3-16.6)	13.4 (10.2-17.4)	16.5 (13.1-20.5)
18-25	11.8 (8.6-16.0)	9.9 (7.5-13.0)	11.7 (8.9-15.1)	13.7 (10.2-18.1)	12.8 (9.5-15.4)	12.1 (9.5-15.4)	12.9 (10.0-16.4)	13.7 (10.9-17.1)	15.0 (11.8-18.9)	19.0 (15.9-22.4)	24.7 (20.8-29.1)	27.2 (23.0-31.8)
≥26	5.9 (4.2-8.3)	6.2 (4.5-8.6)	6.1 (4.4-8.3)	7.0 (5.2-9.3)	8.7 (6.4-11.7)	8.3 (6.1-11.3)	6.4 (4.6-8.9)	9.6 (6.9-13.3)	11.1 (8.3-14.7)	10.3 (7.8-13.5)	14.6 (11.2-18.9)	18.6 (15.2-22.4)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
Abbreviation: CI = confidence interval.
Perceived great risk and no risk from smoking marijuana is defined as those who reported that smoking marijuana once a month and once or twice a week might cause harm (great risk) and might not cause harm (no risk).

Table 7. Percentage of perceived parental disapproval of trying marijuana once or twice and using marijuana once a month or more among persons aged 12 to 17 — National Survey on Drug Use and Health, District of Columbia, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
Perceived parental disapproval of trying marijuana once or twice ¹	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
12–17 years	94.4 (91.8-96.3)	94.2 (92.0-95.3)	92.9 (89.6-95.2)	90.1 (85.5-93.4)	90.6 (86.3-93.6)	91.9 (88.3-94.5)	91.4 (88.1-93.8)	94.1 (91.0-96.2)	94.3 (91.3-96.2)	91.6 (88.6-93.9)	91.0 (87.5-93.5)	92.2 (88.7-94.6)
Perceived parental disapproval of using marijuana once a month or more ²	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
12–17 years	95.2 (93.3-96.7)	95.2 (93.2-96.6)	94.0 (91.3-95.9)	91.5 (87.0-94.5)	92.1 (87.7-95.0)	93.7 (90.1-96.0)	92.6 (89.3-95.0)	94.1 (91.0-96.1)	96.0 (94.2-97.3)	93.8 (91.3-95.7)	91.8 (88.0-94.4)	92.6 (89.0-95.0)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
¹ Perceived parental disapproval of trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of trying marijuana once or twice.
² Perceived parental disapproval of using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of using marijuana once a month or more.

Table 8. Percentage of perceived parental disapproval of trying marijuana once or twice and using marijuana once a month or more among past month marijuana users aged 12 to 17 — National Survey on Drug Use and Health, District of Columbia, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
Perceived parental disapproval of trying marijuana once or twice ¹	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
12–17 years	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Perceived parental disapproval of using marijuana once a month or more ²	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
12–17 years	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
¹ Perceived parental disapproval of trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of trying marijuana once or twice.
² Perceived parental disapproval of using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of using marijuana once a month or more; indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.

Table 9. Percentage of perceived attitudes towards peers trying marijuana once or twice and using marijuana once or more among persons aged 12 to 17— National Survey on Drug Use and Health, District of Columbia, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
Perceived attitudes towards peers trying marijuana once or twice ¹	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
12–17 years	77.3 (74.1-80.2)	80.8 (77.4-83.7)	80.1 (76.8-83.1)	79.0 (74.2-83.0)	78.6 (73.0-83.4)	78.3 (73.9-82.1)	74.8 (69.2-79.7)	74.5 (69.6-79.6)	77.2 (73.7-80.5)	75.0 (71.9-77.9)	71.9 (68.5-75.1)	71.1 (66.9-75.0)
Perceived attitudes towards peers using marijuana once a month or more ²												
12–17 years	78.2 (74.4-81.7)	80.0 (75.8-83.5)	78.5 (74.5-82.1)	77.6 (72.4-82.1)	78.7 (73.0-83.5)	78.9 (73.7-83.3)	74.9 (69.6-79.6)	75.4 (69.9-80.2)	77.8 (73.9-81.2)	74.8 (71.7-77.6)	71.7 (68.2-74.9)	70.9 (66.1-75.3)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
¹ Disapproving attitudes toward peers trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers trying marijuana once or twice.
² Disapproving attitudes toward peers using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers using marijuana once a month or more.

Table 10. Percentage of perceived attitudes towards peers trying marijuana once or twice and using marijuana once a month or more among past month marijuana users aged 12 to 17— National Survey on Drug Use and Health, District of Columbia, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
Perceived attitudes towards peers trying marijuana once or twice ¹	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
12–17 years	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)
Perceived attitudes towards peers using marijuana once a month or more ²												
12–17 years	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
¹ Disapproving attitudes toward peers trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers trying marijuana once or twice; indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.
² Disapproving attitudes toward peers using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers using marijuana once a month or more; indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.

Table 11. Percentage of perceived availability among all persons aged ≥12 and mode of acquisition of marijuana among past year marijuana users aged ≥12 by age group — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived availability												
Total	64.5 (60.4-68.5)	62.2 (58.0-66.1)	63.0 (58.6-67.2)	63.0 (59.3-66.5)	64.1 (60.6-67.5)	63.2 (58.5-67.6)	60.9 (56.3-65.3)	63.1 (59.2-66.9)	66.4 (62.2-70.4)	67.7 (63.0-72.0)	68.3 (63.8-72.5)	68.5 (64.4-72.4)
12-17	51.5 (46.5-56.5)	49.3 (45.2-53.5)	52.9 (48.7-57.2)	54.4 (49.3-59.4)	55.5 (50.6-60.3)	55.0 (49.7-60.2)	51.6 (45.9-57.3)	49.7 (43.8-55.5)	53.8 (49.4-58.2)	52.5 (47.9-57.2)	48.9 (44.2-53.7)	49.4 (43.5-55.3)
18-25	83.8 (79.1-87.6)	78.7 (74.6-82.3)	76.9 (72.6-80.8)	82.7 (78.4-86.3)	83.0 (78.6-86.7)	82.0 (77.6-85.7)	77.9 (73.1-82.0)	80.5 (76.0-84.4)	84.9 (81.2-87.9)	82.3 (78.0-85.8)	80.5 (75.5-84.7)	81.6 (77.0-85.5)
≥26	61.8 (56.4-66.9)	60.1 (54.9-65.1)	61.4 (55.9-66.6)	59.9 (55.3-64.3)	60.7 (56.3-65.0)	59.7 (53.9-65.1)	57.8 (52.0-63.5)	60.3 (55.4-65.1)	63.1 (57.7-68.2)	65.2 (59.4-70.7)	66.9 (61.4-71.9)	67.0 (62.0-71.6)
Mode of acquisition of marijuana												
Bought it												
Total	40.3 (33.6-47.4)	33.5 (26.3-41.4)	34.6 (27.7-42.3)	36.2 (29.2-43.7)	38.4 (30.2-47.2)	41.4 (33.2-50.0)	40.1 (30.5-50.3)	32.1 (23.2-42.5)	28.7 (22.4-36.0)	31.5 (25.2-38.6)	41.1 (35.2-47.3)	50.0 (43.9-56.1)
12-17	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
18-25	39.5 (30.1-47.7)	31.5 (24.7-39.2)	31.9 (23.0-42.5)	35.7 (25.6-47.2)	35.6 (27.9-44.2)	34.4 (28.0-41.3)	32.0 (25.2-39.5)	36.3 (26.9-46.9)	39.0 (32.1-46.3)	39.5 (32.4-47.2)	42.9 (35.3-50.8)	47.9 (39.7-56.2)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	50.7 (42.4-59.1)
Traded something for it												
Total	1.2 (0.4-3.4)	0.7 (0.2-2.8)	1.3 (0.3-5.3)	1.6 (0.5-4.9)	1.2 (0.5-2.9)	1.1 (0.4-3.4)	0.6 (0.1-3.3)	(*)	0.5 (0.1-1.7)	1.8 (0.6-4.9)	1.8 (0.7-4.8)	0.8 (0.3-2.1)
12-17	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	1.0 (0.2-4.7)	2.0 (0.6-6.7)	1.8 (0.4-6.7)	2.6 (0.8-8.1)
18-25	1.3 (0.4-5.0)	(*)	(*)	1.2 (0.3-4.8)	1.4 (0.4-4.4)	0.7 (0.2-3.1)	0.4 (0.0-2.5)	(*)	0.3 (0.0-2.4)	1.3 (0.4-5.0)	1.7 (0.5-5.8)	1.1 (0.2-4.6)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Got it for free or shared with someone else												
Total	58.4 (51.5-64.9)	65.5 (57.6-72.6)	64.0 (56.3-71.1)	62.2 (54.5-69.4)	60.1 (51.3-68.3)	57.0 (48.3-65.2)	58.8 (48.5-68.4)	66.7 (56.3-75.0)	70.4 (63.0-76.0)	66.7 (59.5-73.2)	57.0 (50.8-63.0)	49.1 (42.9-55.3)
12-17	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
18-25	59.9 (51.1-68.0)	66.1 (58.8-72.6)	66.8 (56.2-76.0)	63.2 (51.4-73.6)	62.6 (53.9-70.6)	64.0 (57.0-70.5)	66.7 (58.9-73.7)	63.0 (52.4-72.6)	60.5 (53.0-67.5)	59.1 (51.7-66.1)	55.4 (47.3-63.1)	51.1 (42.7-59.4)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	56.7 (49.5-67.4)	48.8 (40.4-57.2)
Grew it yourself												

Total	0.2 (0.0-0.8)	0.3 (0.1-1.0)	0.1 (0.0-0.8)	*	0.3 (0.1-1.2)	0.5 (0.2-1.5)	0.5 (0.2-1.6)	0.8 (0.2-3.1)	0.5 (0.1-2.8)	*	0.1 (0.0-0.4)	0.1 (0.0-0.5)
12-17	(*) (0.0-1.7)	(*) (0.0-2.0)	(*) (0.0-0.8)	(*) (*)	(*) (*)	(*) (0.2-3.7)	(*) (0.2-3.9)	(*) (0.1-3.1)	(*) (0.0-1.3)	(*) (*)	(*) (*)	(*) (*)
18-25	(*) (0.0-1.7)	(*) (0.0-2.0)	(*) (0.0-0.8)	(*) (*)	(*) (*)	(*) (0.2-3.7)	(*) (0.2-3.9)	(*) (0.1-3.1)	(*) (0.0-1.3)	(*) (*)	(*) (*)	(*) (*)
≥26	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.
Abbreviation: CI = confidence interval.
Perceived availability is defined as those who reported that it would be fairly easy or very easy for them to obtain marijuana if they wanted some.
* Mode of acquisition is defined as those who reported how they got the marijuana they used for the last time [i.e., bought it, traded something for it, got it for free or shared with someone else, or grew it yourself]; this question is only asked of those respondents indicating use of marijuana in the past 12 months.

Table 12. Percentage of past year marijuana dependence and abuse* among all persons aged ≥12 by age group — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	2.4 (1.8-3.4)	1.9 (1.2-2.8)	2.6 (1.8-3.8)	2.6 (1.7-3.9)	2.8 (1.8-4.4)	3.1 (2.1-4.5)	3.1 (2.0-4.8)	3.0 (1.8-4.9)	2.7 (1.8-4.0)	2.4 (1.7-3.5)	2.4 (1.6-3.4)	2.4 (1.7-3.4)
12-17	3.2 (1.8-5.4)	2.6 (1.6-4.3)	3.2 (1.8-5.4)	2.9 (1.4-5.9)	2.7 (1.6-4.6)	2.6 (1.6-4.0)	2.6 (1.5-4.3)	2.6 (1.4-4.8)	4.1 (2.4-7.0)	4.5 (2.6-7.6)	3.8 (2.3-6.5)	4.2 (2.7-6.4)
18-25	6.6 (4.6-9.4)	7.4 (5.0-10.8)	7.2 (4.9-10.5)	5.9 (3.6-9.5)	6.9 (4.6-10.3)	8.7 (6.3-11.8)	6.6 (4.6-9.4)	4.1 (2.8-6.0)	5.3 (3.5-7.9)	6.9 (4.8-9.9)	6.5 (4.6-9.1)	7.2 (4.7-10.9)
≥26	1.6 (0.8-2.9)	0.7 (0.2-2.3)	1.7 (0.9-3.4)	1.9 (1.0-3.4)	1.9 (0.9-4.1)	1.9 (0.9-4.1)	2.4 (1.2-4.7)	2.7 (1.4-5.3)	1.9 (1.0-3.7)	1.2 (0.7-2.3)	1.3 (0.7-2.7)	1.3 (0.6-2.5)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Substance use disorder (also known as dependence or abuse) is defined as meeting criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association [APA], 1994) for either dependence or abuse for illicit drugs or alcohol.

Table 13. Percentage of past year marijuana dependence and abuse* among past year marijuana users† aged ≥12, by age group — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	16.0 (11.5-21.9)	12.9 (8.5-19.1)	17.1 (11.8-24.1)	17.3 (11.7-24.9)	17.8 (11.6-26.2)	19.4 (13.5-27.1)	21.4 (14.8-30.0)	18.2 (11.2-28.1)	14.2 (9.7-20.4)	12.1 (8.5-16.8)	11.3 (8.0-15.8)	11.4 (7.9-16.0)
12-17	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	19.7 (12.0-30.5)	19.4 (12.9-28.0)
18-25	17.8 (12.4-25.0)	24.3 (17.3-33.0)	22.0 (15.3-30.4)	15.7 (9.7-24.3)	17.7 (12.2-25.0)	23.5 (18.0-30.0)	20.1 (14.6-26.9)	12.2 (8.5-17.4)	14.0 (9.3-20.6)	16.1 (11.1-22.8)	15.3 (11.1-20.6)	18.5 (12.3-27.0)
≥26	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	8.3 (4.2-15.6)	7.2 (3.7-13.9)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Substance use disorder (also known as dependence or abuse) is defined as meeting criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association [APA], 1994) for either dependence or abuse for illicit drugs or alcohol.
 † Analysis limited to past year marijuana users (defined as those who reported use of marijuana within 12 months preceding the date of interview, including those who reported use of marijuana within 30 days preceding the date of interview).

Table 14. Percentage of perception of maximum legal penalty in your state for first offense possession of an ounce or less of marijuana for your own use among all persons aged ≥12, by age group — National Survey on Drug Use and Health, District of Columbia, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Fine												
Total	21.0 (17.0-25.6)	19.8 (16.7-23.4)	20.1 (16.5-24.3)	20.5 (17.1-24.4)	20.2 (16.1-25.0)	21.4 (17.9-25.2)	19.8 (15.9-24.3)	21.1 (16.8-26.0)	24.7 (20.1-29.9)	23.4 (19.6-27.7)	22.0 (18.1-26.3)	37.3 (33.4-41.3)
12-17	13.5 (9.9-18.2)	12.4 (9.4-16.2)	11.5 (8.5-15.3)	13.6 (10.6-17.4)	16.3 (13.1-20.1)	15.1 (11.2-20.1)	15.2 (10.6-21.3)	14.8 (10.7-20.1)	13.2 (10.0-17.2)	13.5 (10.5-17.1)	16.3 (13.2-19.9)	25.6 (21.1-30.7)
18-25	16.6 (12.0-22.3)	13.7 (9.3-19.7)	12.0 (9.3-15.3)	11.7 (8.8-15.3)	11.9 (8.5-16.4)	14.0 (10.1-19.1)	16.5 (13.1-20.7)	18.3 (14.8-22.4)	20.8 (16.5-25.9)	19.6 (15.8-24.0)	18.8 (14.9-23.6)	35.3 (29.4-41.7)
≥26	22.9 (17.7-29.1)	22.3 (18.1-27.0)	23.1 (18.3-28.6)	23.7 (19.2-28.9)	22.9 (17.2-29.7)	24.0 (19.4-29.3)	21.3 (16.2-27.5)	22.6 (16.9-29.4)	26.9 (21.1-33.6)	25.5 (20.5-31.3)	23.3 (18.1-29.6)	39.8 (34.0-43.8)
Probation												
Total	20.8 (17.1-25.0)	21.6 (18.1-25.6)	20.2 (16.8-23.9)	18.5 (15.5-22.0)	20.2 (16.5-24.3)	19.3 (16.1-23.0)	17.8 (14.4-21.9)	17.4 (13.6-21.8)	17.7 (14.2-21.8)	19.4 (15.8-23.6)	21.9 (18.0-26.5)	17.8 (15.0-21.1)
12-17	24.9 (20.0-30.5)	24.4 (18.6-31.2)	25.0 (19.8-30.9)	23.1 (18.6-28.2)	23.8 (19.9-28.2)	25.9 (22.1-30.1)	23.2 (18.2-29.2)	23.1 (17.5-29.9)	27.4 (23.1-32.1)	29.7 (25.8-34.1)	27.6 (22.4-33.4)	25.1 (19.6-31.5)
18-25	26.8 (21.5-32.8)	24.2 (19.9-29.0)	19.6 (15.7-24.3)	20.0 (14.9-26.3)	21.5 (16.4-27.8)	23.1 (17.9-29.2)	22.0 (17.7-27.0)	20.4 (16.8-24.5)	18.7 (14.9-23.1)	22.0 (18.6-25.9)	27.1 (23.1-31.6)	21.6 (17.8-26.0)
≥26 years	18.9 (13.9-25.0)	20.6 (15.8-26.4)	19.7 (15.4-24.8)	17.5 (13.8-22.0)	19.3 (14.7-25.0)	17.6 (13.5-22.6)	15.9 (11.9-21.0)	15.8 (11.4-21.7)	16.5 (11.9-22.4)	17.6 (12.8-23.7)	20.0 (15.0-26.2)	16.3 (12.6-20.7)
Community Service												
Total	9.5 (6.7-13.3)	11.8 (8.8-15.8)	10.7 (8.0-14.1)	8.8 (6.4-12.0)	9.2 (6.5-12.8)	8.9 (6.3-12.5)	9.4 (6.8-13.4)	8.0 (5.6-11.3)	7.4 (5.3-10.3)	8.6 (6.2-11.8)	7.8 (5.9-10.4)	5.9 (4.6-7.5)
12-17	11.5 (8.4-15.5)	13.5 (9.7-18.5)	15.2 (11.5-19.8)	12.0 (9.1-15.7)	6.9 (4.8-9.9)	7.6 (5.2-11.0)	11.0 (8.0-14.9)	9.6 (6.8-13.4)	7.1 (5.2-9.7)	7.4 (5.1-10.6)	8.2 (5.9-11.1)	8.1 (5.8-11.4)
18-25	10.1 (7.1-14.2)	8.0 (5.6-11.3)	11.7 (8.9-15.3)	11.8 (9.1-15.2)	8.4 (6.0-11.8)	7.8 (5.1-11.8)	9.9 (7.0-13.8)	10.7 (7.7-14.6)	10.1 (6.8-14.8)	8.8 (6.5-11.8)	8.5 (6.1-11.9)	6.6 (4.4-9.8)
≥26	9.2 (5.8-14.2)	12.6 (8.6-18.0)	9.9 (6.5-14.7)	7.6 (4.5-12.4)	9.6 (6.3-14.4)	9.3 (6.0-14.3)	9.1 (5.4-15.1)	7.1 (4.0-12.3)	6.7 (4.1-10.8)	8.6 (5.5-13.2)	7.6 (5.2-11.0)	5.5 (3.9-7.7)
Possible Prison Sentence												
Total	40.8 (36.0-45.7)	40.2 (34.8-45.9)	42.1 (37.6-46.7)	43.1 (39.1-47.2)	39.6 (35.7-43.7)	38.9 (33.8-44.3)	39.8 (34.5-45.4)	38.3 (33.6-43.3)	36.9 (32.0-42.2)	37.3 (33.1-41.6)	36.6 (32.5-40.9)	27.7 (24.7-30.8)

12-17	37.1 (31.9-42.7)	37.5 (32.3-42.9)	37.7 (32.5-43.2)	34.0 (28.6-40.0)	32.0 (28.0-36.2)	30.7 (25.7-36.3)	32.2 (25.6-39.6)	35.2 (30.1-40.7)	32.2 (27.1-37.7)	28.5 (24.8-32.7)	28.6 (23.5-34.3)	26.1 (21.3-31.5)
18-25	37.0 (31.3-43.2)	42.9 (37.7-48.1)	46.3 (41.4-51.3)	46.7 (42.2-51.4)	45.2 (40.0-50.5)	40.6 (35.1-46.4)	36.8 (32.3-41.4)	35.8 (32.3-39.6)	37.6 (32.9-42.5)	38.8 (34.6-43.2)	34.4 (29.8-39.3)	25.6 (21.3-30.3)
≥26	42.1 (35.9-48.6)	39.9 (32.7-47.6)	41.6 (35.6-47.9)	43.5 (37.6-49.5)	39.1 (33.3-45.2)	39.5 (32.5-46.9)	41.7 (34.5-49.3)	39.4 (32.8-46.4)	37.2 (30.8-44.1)	37.7 (32.2-43.5)	38.0 (32.7-43.6)	28.3 (24.7-32.2)
Mandatory Prison Sentence												
Total	7.9 (5.6-11.0)	6.6 (5.1-8.3)	7.0 (5.2-9.2)	6.7 (5.1-8.8)	6.6 (5.0-8.5)	6.8 (4.9-9.6)	8.2 (6.9-11.2)	8.6 (5.7-12.7)	5.9 (4.4-7.9)	5.2 (3.7-7.3)	4.9 (3.5-6.8)	2.8 (1.9-4.3)
12-17	13.0 (9.8-17.2)	12.2 (9.0-16.4)	10.6 (8.1-13.8)	12.5 (9.1-17.0)	13.5 (10.2-17.8)	13.5 (10.2-17.7)	11.0 (7.8-15.3)	8.9 (6.4-12.0)	11.0 (7.6-15.6)	9.7 (6.5-14.2)	7.9 (5.5-11.2)	6.4 (4.1-9.8)
18-25	9.6 (6.5-13.8)	11.3 (8.4-15.0)	10.4 (7.8-13.7)	9.3 (6.5-13.1)	9.6 (6.2-14.5)	9.5 (6.4-14.0)	9.4 (6.8-12.9)	9.4 (6.9-12.6)	7.5 (5.2-10.8)	5.5 (3.4-8.6)	5.7 (3.6-8.8)	3.6 (1.9-6.9)
≥26	7.0 (4.3-11.2)	4.7 (3.0-7.2)	5.7 (3.6-9.0)	5.3 (3.3-8.3)	4.9 (3.1-7.6)	5.3 (2.8-9.8)	7.5 (4.6-11.9)	8.3 (4.8-14.0)	5.0 (3.4-7.3)	4.8 (2.7-8.3)	4.4 (2.6-7.4)	2.3 (1.3-4.2)
No penalty												
Total	DNA	DNA	DNA	2.3 (1.3-4.0)	4.3 (2.8-6.6)	4.7 (3.1-7.1)	5.0 (3.4-7.1)	6.7 (4.6-9.6)	7.3 (5.1-10.5)	6.1 (4.0-9.1)	6.8 (4.7-9.7)	8.5 (6.5-11.1)
12-17	DNA	DNA	DNA	4.7 (2.6-8.2)	7.4 (5.0-10.9)	7.1 (4.4-11.4)	7.4 (5.1-10.7)	8.5 (5.9-12.0)	9.1 (6.4-12.8)	11.1 (8.1-15.1)	11.5 (8.5-15.4)	8.8 (6.4-11.8)
18-25	DNA	DNA	DNA	0.5 (0.1-2.6)	3.3 (1.7-5.2)	5.0 (3.0-8.2)	5.5 (3.3-8.9)	5.4 (2.9-9.9)	5.3 (2.8-10.0)	5.3 (3.4-8.3)	5.4 (3.9-7.6)	7.3 (4.9-10.6)
≥26	DNA	DNA	DNA	2.5 (1.3-4.7)	4.2 (2.3-7.4)	4.4 (2.4-7.8)	4.5 (2.6-7.6)	6.8 (4.0-11.3)	7.7 (4.8-12.2)	5.8 (3.3-10.1)	6.7 (4.1-10.8)	8.8 (6.2-12.4)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.
Abbreviations: CI = confidence interval; DNA = data not available.
Perception of legal penalty for marijuana possession is defined as those who reported their perception of the maximum legal penalty in their state of residence for first offense possession of an ounce or less of marijuana for their own use.

Supplementary Material

Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, Oregon, 2002–2014

Suggested Citation: Alejandro Azofeifa, Margaret E. Mattson, and Rob Lryerla. Supplementary Material. State Level Data: Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, Oregon, 2002–2014. Center for Behavioral Health Statistics and Quality. (2016). Substance Abuse and Mental Health Services Administration, Rockville, MD.

Notes: A) Two years of data were combined to allow for reliable state-level prevalence estimates. Some state-level prevalence estimates for combined years have been suppressed (indicated by an asterisk (*) symbol in each cell) based on the following National Survey on Drug Use and Health (NSDUH) suppression criteria. Prevalence estimates are suppressed if any of the following occurred: 1) the prevalence estimate is $<0.005\%$ or $>99.995\%$; 2) the relative standard error (RSE) of the estimated proportion p (where p is the prevalence divided by 100) is ≥ 0.175 if the prevalence is $\geq 50\%$; 3) the RSE of the negative natural logarithm of $(1-p)$ is ≥ 0.175 if the prevalence is $>50\%$; 4) the actual sample size is <100 ; or 5) the effective sample size (defined as the actual sample size divided by the design effect) is <68 . B) The estimates provided in this supplementary material for each state and the District of Columbia are from design-based direct estimation procedures (i.e., they are based directly on the respondent sample and the accompanying analytical weights with variances derived through a Taylor-linearization approach using SUDAAN software) as used for the national level estimates (available at: Azofeifa A, Mattson ME, Schauer C, McAfee T, Grant A, Lryerla R. National Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, United States, 2002–2014. *MMWR Surveill Summ* 2016;65(No. SS-11):1–25. DOI: <http://dx.doi.org/10.15585/mmwr.ss6511a1>). Caution is advised when comparing data from other NSDUH reports that are based on different design-based estimation procedures. C) Interpretation of state-level trend data should be done with caution as some prevalence estimates are suppressed based on the above criteria and survey cycle differences may not be statistically significant.

BILAG 4

Table 1. Percentage of past month marijuana use* among all persons aged ≥ 12, by selected demographic characteristics — National Survey on Drug Use and Health, Oregon, 2002–2014

Characteristics	2002–2003	2003–2004	2004–2005	2005–2006	2006–2007	2007–2008	2008–2009	2009–2010	2010–2011	2011–2012	2012–2013	2013–2014
Age group (yrs)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	9.8 (7.3-13.0)	8.4 (6.1-11.4)	8.7 (7.2-10.6)	8.1 (6.5-10.2)	8.1 (6.3-10.5)	9.3 (7.4-11.7)	9.7 (8.1-11.6)	10.5 (8.8-12.5)	11.8 (9.8-14.0)	13.0 (11.1-15.2)	12.9 (10.1-16.4)	12.7 (10.5-15.4)
12–17	8.5 (6.6-10.8)	9.9 (7.4-13.0)	10.3 (7.1-14.6)	9.3 (6.5-13.2)	7.9 (5.8-10.6)	7.5 (5.4-10.3)	8.9 (7.1-11.0)	10.0 (7.8-12.8)	10.1 (7.8-12.9)	9.4 (7.2-12.2)	9.7 (7.3-12.7)	11.0 (8.3-14.5)
18–25	22.7 (17.3-29.2)	21.3 (17.1-26.2)	20.6 (17.1-24.7)	17.8 (13.6-22.8)	17.9 (13.9-22.6)	21.8 (18.0-26.1)	25.2 (21.4-29.5)	25.1 (21.6-28.9)	26.0 (21.3-31.2)	26.2 (21.3-31.8)	22.9 (18.7-27.7)	24.3 (20.3-28.8)
≥26	7.7 (5.2-11.4)	6.0 (3.7-9.4)	6.5 (4.8-8.9)	6.4 (4.6-8.9)	6.6 (4.6-9.5)	7.6 (5.5-10.3)	7.4 (5.6-9.7)	8.3 (6.3-10.8)	9.7 (7.3-12.6)	11.3 (9.0-14.2)	11.7 (8.4-16.1)	11.1 (8.5-14.4)
26–34	9.9 (5.7-16.7)	11.5 (6.2-20.4)	14.0 (9.2-20.8)	13.6 (9.5-19.2)	12.0 (8.4-16.9)	13.0 (8.5-19.3)	11.5 (6.9-18.6)	10.4 (6.3-16.6)	18.9 (13.5-25.9)	21.3 (14.5-30.1)	16.0 (10.3-23.9)	20.8 (15.0-28.1)
35–44	8.3 (4.4-15.2)	8.2 (4.9-13.3)	9.9 (5.6-17.0)	7.4 (3.6-14.5)	3.4 (1.5-7.4)	6.5 (3.4-11.8)	7.4 (4.1-13.2)	9.8 (6.1-15.4)	11.9 (7.2-19.1)	12.2 (7.3-19.7)	12.6 (7.0-21.7)	10.6 (6.0-18.1)
45–54	10.9 (5.5-20.5)	4.5 (1.9-10.4)	4.9 (2.6-9.2)	7.8 (3.7-15.7)	7.8 (3.7-15.7)	6.8 (3.4-13.0)	6.1 (3.0-12.2)	10.0 (5.6-17.2)	7.5 (3.9-14.2)	8.5 (5.1-13.9)	11.0 (6.7-17.6)	10.6 (6.2-17.4)
55–64	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
≥65	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Sex												
Male	11.2 (8.0-15.6)	9.3 (6.7-12.8)	10.2 (7.9-13.1)	9.9 (7.3-13.3)	9.3 (6.7-12.7)	10.8 (7.9-14.7)	11.2 (8.5-14.6)	13.0 (9.9-16.8)	15.5 (12.3-19.4)	17.8 (14.6-21.5)	18.0 (13.6-23.5)	17.1 (13.4-21.5)
Female	8.4 (6.0-11.6)	7.5 (5.1-11.0)	7.3 (5.5-9.6)	6.4 (4.7-8.6)	7.0 (5.0-9.7)	7.8 (5.8-10.5)	8.3 (6.5-10.4)	8.2 (6.8-9.8)	8.1 (6.4-10.2)	8.5 (6.6-10.8)	9.0 (6.1-10.5)	8.6 (6.8-10.8)
Race/ethnicity												
Non-Hispanic white	10.2 (7.4-14.0)	8.7 (6.3-12.1)	9.2 (7.5-11.2)	8.6 (6.6-11.1)	8.6 (6.4-11.4)	9.5 (7.4-12.0)	9.5 (7.9-11.5)	10.3 (8.6-12.3)	12.6 (10.3-15.4)	14.3 (12.0-17.1)	13.4 (10.3-17.3)	13.2 (10.7-16.2)
Non-Hispanic black	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic American Indian/Alaska Native	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic Hawaiian/Other Pacific Islander	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic Asian	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic two or more races	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Hispanic	6.9 (2.9-15.5)	5.0 (2.2-10.9)	4.3 (1.9-9.5)	7.5 (3.4-15.6)	6.7 (3.2-13.5)	3.4 (1.8-6.4)	6.5 (2.7-14.6)	11.0 (6.1-19.2)	7.0 (3.7-12.8)	3.5 (1.7-7.0)	9.5 (4.8-18.0)	12.4 (6.9-21.4)
Current employment status												

Full-time	9.1 (5.9-13.8)	7.8 (4.9-12.2)	7.7 (5.5-10.6)	8.1 (5.8-11.2)	9.6 (6.4-14.1)	10.3 (7.2-14.6)	9.3 (7.2-11.9)	9.3 (6.8-12.7)	10.7 (8.1-14.1)	11.9 (9.0-15.7)	11.4 (8.4-15.4)	12.5 (9.5-16.2)
Part-time	12.4 (7.1-20.8)	11.1 (6.3-18.9)	12.9 (9.2-17.7)	9.6 (6.4-14.1)	9.7 (5.9-15.7)	13.4 (8.6-20.3)	13.3 (9.1-19.0)	15.1 (10.4-21.4)	17.7 (12.3-24.7)	16.6 (12.6-21.5)	13.3 (8.4-20.5)	12.4 (8.3-18.0)
Unemployed	(*)	(*)	(*)	(*)	(*)	11.9 (6.6-20.6)	20.9 (12.8-32.3)	26.1 (17.6-36.8)	(*)	(*)	18.0 (10.6-28.9)	(*)
Other ²	8.7 (5.2-14.2)	5.6 (2.6-11.4)	5.6 (3.2-9.7)	5.7 (3.1-10.1)	5.0 (2.6-9.1)	6.3 (3.7-10.6)	6.7 (4.1-10.8)	6.7 (4.2-10.5)	8.7 (5.2-14.3)	12.6 (8.9-17.5)	14.9 (9.0-23.5)	12.3 (7.9-18.8)
Highest level of education completed												
< High school	10.6 (7.7-14.4)	9.2 (5.5-15.1)	14.5 (9.1-22.5)	12.2 (7.1-20.1)	10.8 (5.9-19.1)	12.9 (6.8-23.1)	12.1 (6.6-20.9)	14.5 (9.1-22.3)	16.8 (10.5-25.6)	22.7 (15.7-31.5)	*	18.2 (11.8-27.1)
High school graduate	9.0 (5.0-15.6)	9.3 (5.4-15.5)	7.8 (5.6-10.8)	8.7 (6.0-12.5)	8.9 (5.9-13.4)	9.7 (6.3-14.7)	10.2 (6.9-14.8)	9.1 (6.0-13.4)	10.5 (7.0-15.5)	13.0 (8.9-18.4)	16.4 (10.2-25.5)	16.5 (11.2-23.7)
Some college	10.3 (6.5-16.0)	9.2 (6.2-13.4)	10.4 (7.1-15.0)	8.3 (4.9-13.7)	10.3 (6.7-15.3)	10.5 (6.8-15.8)	9.6 (6.8-13.4)	11.4 (8.3-15.4)	13.0 (9.7-17.2)	15.0 (11.4-19.5)	12.9 (9.5-17.4)	13.2 (9.5-18.1)
College graduate	10.2 (5.2-19.1)	5.7 (2.5-12.4)	5.4 (2.9-9.7)	5.3 (3.2-8.5)	4.1 (2.5-6.7)	6.7 (4.4-10.1)	8.6 (5.5-13.4)	9.9 (6.5-14.8)	10.2 (6.5-15.6)	7.9 (5.0-12.5)	6.5 (4.0-10.3)	7.9 (5.2-11.9)
Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.												
Abbreviation: CI = confidence interval.												
* Past month use of marijuana is defined as those who reported use of marijuana within 30 days preceding the date of interview.												
¹ Current employment and highest level of education completed are only defined for 18 or older.												
² Other employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.												

Table 2. Percentage of past year marijuana use among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Oregon, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	14.8 (11.8-18.4)	13.6 (10.7-17.1)	14.1 (12.1-16.5)	14.0 (12.2-16.2)	14.5 (11.9-17.4)	15.1 (12.6-18.1)	15.4 (13.4-17.7)	16.5 (14.3-18.9)	16.9 (14.7-19.3)	18.4 (15.8-21.4)	20.6 (17.2-24.6)	20.1 (17.3-23.1)
12-17	16.4 (13.3-20.1)	18.1 (14.8-22.0)	19.4 (15.1-24.5)	19.7 (15.3-25.1)	16.5 (13.5-20.1)	16.7 (13.0-21.1)	19.4 (16.0-23.4)	21.0 (17.2-25.4)	18.6 (15.0-22.9)	15.4 (12.6-18.8)	16.4 (13.3-20.1)	18.3 (14.7-22.7)
18-25	36.6 (30.0-43.7)	34.6 (29.3-40.4)	35.9 (31.3-40.8)	32.9 (28.1-38.1)	33.9 (28.5-39.8)	37.7 (33.3-42.3)	38.7 (34.4-43.1)	39.9 (35.4-44.6)	40.4 (35.5-45.4)	38.4 (33.2-43.8)	34.9 (29.5-40.7)	37.7 (32.9-42.8)
≥26	10.9 (7.9-14.8)	9.4 (6.6-13.2)	9.7 (7.5-12.6)	10.2 (8.1-12.8)	11.1 (8.5-14.4)	11.4 (8.9-14.6)	11.4 (9.3-13.9)	12.3 (9.8-15.2)	12.9 (10.4-15.9)	15.6 (12.6-19.2)	18.9 (14.9-23.6)	17.5 (14.4-21.1)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

Past year use of marijuana is defined as those who reported use of marijuana within 12 months preceding the date of interview. — these also include those who reported use of marijuana within 30 days preceding the date of interview.

Table 3. Percentage of daily or almost daily marijuana use in the past year and past month among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Oregon, 2002–2014

Age group (yrs)	2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014	
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Daily or almost daily marijuana use in the past year¹																								
Total	2.8 (1.6-4.9)	2.2 (1.3-3.8)	2.1 (1.5-2.9)	1.5 (1.0-2.4)	1.5 (0.9-2.4)	1.5 (0.9-2.4)	2.2 (1.4-3.4)	2.5 (1.6-3.8)	2.6 (2.0-3.5)	2.9 (2.3-3.7)	3.8 (2.8-5.1)	4.9 (3.4-7.1)	4.9 (3.4-7.1)											
12-17	1.0 (0.5-2.3)	2.5 (1.5-4.2)	2.4 (1.3-4.1)	1.1 (0.4-2.7)	0.7 (0.3-2.0)	0.4 (0.1-1.5)	0.4 (0.1-1.5)	1.2 (0.5-3.2)	3.6 (2.0-6.2)	3.0 (1.7-5.3)	1.5 (0.8-2.9)	1.3 (0.7-2.4)	1.3 (0.4-4.1)											
18-25	8.0 (5.1-12.4)	6.7 (4.9-9.2)	6.8 (4.6-9.9)	5.7 (3.7-8.8)	3.8 (2.4-5.9)	5.8 (3.6-9.1)	7.4 (5.5-10.0)	7.4 (5.5-10.0)	6.7 (4.7-9.4)	8.6 (6.2-11.9)	10.2 (7.6-13.6)	9.6 (7.4-12.3)	9.7 (7.2-12.9)											
≥26	2.1 (0.9-4.9)	1.4 (0.5-4.0)	1.2 (0.5-2.3)	0.9 (0.4-1.9)	1.2 (0.5-2.6)	1.9 (1.0-3.5)	1.9 (1.0-3.5)	1.9 (1.0-3.5)	1.9 (1.1-3.1)	2.0 (1.3-3.0)	3.0 (1.9-4.7)	4.6 (2.8-7.3)	4.6 (2.8-7.3)											
Daily or almost daily marijuana use in the past month²																								
Total	4.5 (2.9-6.9)	3.3 (2.1-5.2)	2.8 (2.2-3.6)	2.4 (1.8-3.2)	2.9 (1.9-4.4)	3.6 (2.4-5.2)	3.6 (2.5-5.0)	3.6 (2.5-5.0)	4.0 (3.1-5.2)	4.3 (3.6-5.2)	5.1 (4.0-6.4)	6.5 (4.7-9.0)	6.5 (4.8-8.6)											
12-17	1.6 (0.9-3.1)	2.8 (1.6-4.7)	2.4 (1.4-4.2)	1.0 (0.4-2.2)	1.1 (0.5-2.3)	0.7 (0.3-1.7)	1.6 (0.8-3.1)	1.6 (0.8-3.1)	4.8 (3.2-7.0)	4.4 (2.8-6.8)	2.6 (1.7-4.0)	2.7 (1.7-4.2)	2.9 (1.6-5.1)											
18-25	11.3 (8.4-15.1)	9.7 (7.1-13.2)	8.3 (5.9-11.5)	6.9 (4.7-10.2)	6.0 (4.2-8.5)	8.1 (5.7-11.4)	10.0 (7.9-12.7)	10.0 (7.9-12.7)	11.0 (8.4-14.4)	13.3 (10.0-17.4)	13.2 (10.7-16.3)	11.7 (9.2-14.8)	12.1 (9.4-15.4)											
≥26	3.7 (2.0-6.8)	2.3 (1.1-4.8)	1.9 (1.3-2.8)	1.8 (1.3-2.7)	2.7 (1.5-4.6)	3.2 (1.9-5.4)	2.8 (1.6-4.8)	2.8 (1.6-4.8)	2.8 (1.8-4.4)	2.8 (2.0-4.0)	4.1 (2.9-5.7)	6.1 (4.0-9.3)	6.0 (4.1-8.8)											

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Past year daily or almost daily use is defined as having used marijuana on 300 or more days in the past year.

² Past month daily or almost daily use is defined as having used marijuana on 20 or more days in the past month.

Table 4. Percentage of daily or almost daily marijuana use in the past year and past month among marijuana users aged ≥12, by age group — National Survey on Drug Use and Health, Oregon, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2010-2011	2011-2012	2012-2013	2013-2014	
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	
Daily or almost daily marijuana use in the past year*												
Total	16.8 (11.6-29.0)	16.1 (9.8-25.3)	14.5 (10.1-20.4)	10.9 (7.0-16.5)	10.1 (6.3-16.6)	14.5 (9.7-21.1)	16.1 (11.0-23.0)	15.9 (12.1-20.7)	17.3 (13.2-22.4)	20.5 (15.1-27.2)	23.8 (17.5-31.6)	24.5 (17.5-33.2)
12-17	6.4 (3.0-13.2)	13.8 (8.8-21.1)	12.3 (7.1-20.4)	5.5 (2.4-12.5)	4.5 (1.7-11.2)	(*) (*)	(*) (*)	16.9 (10.5-26.2)	16.0 (10.0-24.5)	10.0 (5.4-17.7)	(*) (*)	(*) (*)
18-25	21.9 (15.2-30.4)	19.4 (14.5-25.3)	19.0 (13.2-26.5)	17.4 (11.3-25.8)	11.2 (7.2-16.8)	15.4 (9.8-23.5)	19.2 (14.7-24.7)	16.7 (11.7-23.2)	21.3 (15.5-28.6)	26.6 (20.5-33.8)	27.5 (22.5-33.3)	25.6 (19.1-33.5)
≥26	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	19.3 (12.2-29.1)	24.2 (16.1-34.8)	(*) (*)
Daily or almost daily marijuana use in the past month*												
Total	45.9 (34.8-57.4)	39.8 (29.3-51.4)	32.0 (24.5-40.7)	29.7 (22.1-38.7)	36.0 (26.7-46.5)	38.4 (28.6-49.1)	36.6 (27.5-46.8)	38.4 (30.1-47.4)	36.6 (29.0-44.9)	39.1 (31.6-47.3)	50.5 (42.8-58.2)	50.8 (42.3-59.4)
12-17	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)
18-25	49.9 (42.1-57.6)	45.8 (36.2-55.7)	40.2 (30.6-50.7)	(*) (*)	33.8 (24.3-44.9)	37.2 (26.9-48.9)	39.7 (30.9-49.3)	43.9 (34.1-54.2)	51.1 (40.7-61.4)	50.4 (42.4-58.3)	51.1 (41.9-60.3)	49.7 (40.7-58.8)
≥26	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Analysis limited to marijuana users (i.e., estimate of daily or almost daily use in the past year is among past year marijuana users; estimates of daily or almost daily use in the past month is among past month marijuana users).
 † Past year daily or almost daily use is defined as having used marijuana on 300 or more days in the past year.
 ‡ Past month daily or almost daily use is defined as having used marijuana on 20 or more days in the past month.

Table 5. Percentage of past year marijuana initiation among persons aged ≥12 years at risk for initiation, and mean age at first use of marijuana among past year marijuana initiates, by age group — National Survey on Drug Use and Health, Oregon, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	%	%	%	%	%	%	%	%	%	%	%	%
	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)
Past Year marijuana initiation among persons at risk for initiation												
Total	1.3	1.8	1.7	2.1	2.5	2.1	2.3	2.2	2.0	2.7	2.8	1.8
	(1.0-1.8)	(1.2-2.8)	(1.2-2.4)	(1.5-3.0)	(1.7-3.7)	(1.3-3.4)	(1.6-3.3)	(1.6-3.1)	(1.5-2.8)	(1.7-4.4)	(1.7-4.7)	(1.3-2.6)
12-17	5.0	6.8	6.6	8.1	8.3	7.3	9.1	8.8	7.1	6.3	7.8	7.9
	(3.6-7.0)	(4.8-9.5)	(4.5-9.6)	(5.5-11.7)	(5.9-11.7)	(4.8-11.1)	(6.6-12.3)	(6.4-11.9)	(5.4-9.3)	(4.4-8.9)	(5.3-11.2)	(5.5-11.2)
18-25	5.3	5.2	4.4	4.6	7.5	8.2	7.2	8.3	9.2	8.4	5.1	4.1
	(3.1-8.9)	(3.1-8.7)	(2.5-7.7)	(1.8-11.3)	(3.7-14.7)	(4.5-14.6)	(4.0-12.6)	(4.7-14.2)	(5.6-14.8)	(5.0-13.6)	(2.4-10.6)	(2.2-7.5)
≥26	*	0.2	0.2	0.5	0.5	*	*	*	*	1.0	1.1	*
	(**)	(0.0-1.7)	(0.0-1.7)	(0.1-1.8)	(0.1-1.9)	(**)	(**)	(**)	(**)	(0.2-4.1)	(0.3-4.6)	(**)
Estimated no. (in 1,000s) of past year marijuana initiation among those at risk for initiation												
Total	21	29	26	33	40	32	34	35	33	41	40	27
	(15-28)	(19-43)	(18-38)	(23-47)	(27-59)	(20-51)	(24-48)	(25-49)	(25-45)	(26-66)	(24-66)	(19-38)
12-17	12	17	16	20	22	18	22	21	17	16	20	20
	(9-17)	(12-23)	(11-23)	(14-29)	(15-30)	(12-28)	(16-30)	(16-29)	(13-23)	(11-23)	(14-29)	(14-28)
18-25	8	9	7	7	13	14	12	14	16	15	9	7
	(5-14)	(5-15)	(4-12)	(3-18)	(6-26)	(8-25)	(7-20)	(8-24)	(10-25)	(9-24)	(4-19)	(4-13)
≥26	*	3	3	5	5	*	*	*	*	11	11	*
	(**)	(0-21)	(0-20)	(1-21)	(1-22)	(**)	(**)	(**)	(**)	(2-45)	(2-44)	(**)
Mean age at first use among past year marijuana initiates												
Total	16.3	17.6	17.2	17.9	18.0	16.6	16.6	16.7	17.1	27.8	27.5	16.1
	(15.6-16.9)	(15.5-19.7)	(15.0-19.5)	(15.3-20.5)	(15.9-20.1)	(15.7-17.5)	(15.8-17.4)	(15.8-17.6)	(16.4-17.9)	(16.0-39.6)	(15.0-40.0)	(15.2-17.0)
12-17	14.7	15.0	14.7	14.8	15.0	15.0	15.2	14.9	14.6	14.9	14.6	14.7
	(14.0-15.4)	(14.6-15.5)	(14.2-15.1)	(14.4-15.3)	(14.6-15.3)	(14.6-15.4)	(14.8-15.5)	(14.5-15.2)	(14.3-14.9)	(14.4-15.4)	(14.2-15.1)	(14.3-15.1)
18-25	18.5	19.4	19.3	18.6	18.6	18.7	19.2	19.5	19.9	20.0	19.8	19.9
	(17.7-19.3)	(18.2-20.6)	(17.9-20.7)	(17.9-19.4)	(18.0-19.2)	(17.9-19.5)	(18.4-20.0)	(18.7-20.3)	(19.0-20.8)	(18.7-21.2)	(18.0-21.5)	(18.4-21.4)
≥26	*	*	*	*	*	*	*	*	*	*	*	*
	(**)	(**)	(**)	(**)	(**)	(**)	(**)	(**)	(**)	(**)	(**)	(**)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

* Past year marijuana initiation (initiates) among persons at risk for initiation is defined as those having first used marijuana within the 12-month period before the date of the interview.

Table 6. Percentage of perceived great risk and no risk from smoking marijuana once a month and once or twice a week among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Oregon, 2002–2014

Age group (yrs)	2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014	
	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Perceived great risk from smoking marijuana once a month																								
Total	31.1	(26.9-35.6)	33.4	(29.1-38.0)	28.9	(25.3-32.8)	28.1	(25.2-31.3)	28.3	(25.3-31.5)	27.5	(23.3-32.2)	25.6	(22.1-29.5)	21.3	(17.7-25.4)	19.4	(16.2-23.0)	19.7	(16.5-23.3)	18.1	(14.7-22.1)	16.5	(13.4-20.3)
12-17	28.6	(24.8-32.8)	32.2	(28.0-36.6)	30.7	(26.3-35.5)	28.9	(25.1-33.2)	28.2	(24.0-32.7)	26.6	(22.5-31.2)	26.1	(22.0-30.7)	24.3	(19.8-29.4)	23.2	(19.6-27.1)	22.3	(18.8-26.2)	19.9	(16.2-24.1)	17.8	(14.3-22.0)
18-25	16.0	(13.0-19.4)	20.7	(16.2-26.0)	19.5	(15.3-24.5)	16.2	(12.9-20.1)	17.1	(14.1-20.6)	17.1	(14.0-20.6)	14.1	(11.3-17.4)	10.1	(7.5-13.6)	11.8	(8.6-15.9)	14.4	(11.5-17.8)	12.2	(9.8-15.1)	9.5	(7.1-12.6)
≥26	34.1	(29.0-39.5)	35.8	(30.6-41.3)	30.3	(25.9-35.2)	30.0	(26.3-34.1)	30.1	(26.4-34.0)	29.2	(24.0-35.0)	27.3	(22.9-32.2)	22.8	(18.3-28.0)	20.2	(16.4-24.6)	20.3	(16.4-24.7)	18.8	(14.6-24.0)	17.5	(13.6-22.4)
Perceived no risk from smoking marijuana once a month																								
Total	13.8	(11.1-17.2)	12.3	(10.1-14.9)	11.9	(9.8-14.4)	12.6	(10.7-14.7)	14.1	(11.9-16.8)	18.0	(14.5-22.2)	20.2	(16.9-23.9)	18.8	(16.1-22.0)	20.4	(17.9-23.1)	23.4	(20.4-26.6)	23.9	(20.8-27.4)	23.5	(20.7-26.6)
12-17	9.3	(7.1-12.2)	8.9	(6.5-12.1)	9.6	(7.2-12.8)	10.0	(7.6-13.1)	9.7	(7.5-12.4)	10.5	(8.0-13.6)	13.6	(10.6-17.3)	17.3	(14.0-21.2)	18.6	(15.7-21.9)	18.4	(14.8-22.5)	19.1	(15.8-23.1)	19.8	(16.4-23.7)
18-25	26.7	(21.9-32.1)	23.4	(19.3-28.2)	23.8	(19.9-28.2)	22.3	(18.2-27.0)	20.5	(17.1-24.2)	25.7	(21.5-30.3)	29.2	(25.2-33.6)	31.0	(25.7-36.8)	34.7	(29.9-39.9)	34.3	(28.8-40.2)	33.7	(28.1-39.7)	38.5	(33.6-43.7)
≥26	12.2	(9.0-16.2)	10.8	(8.3-14.0)	10.2	(7.6-13.4)	11.3	(9.0-14.0)	13.7	(10.9-17.0)	17.8	(13.4-23.1)	19.5	(15.6-24.2)	17.1	(13.9-20.8)	18.2	(15.2-21.8)	22.2	(18.8-26.2)	22.9	(19.1-27.3)	21.5	(18.2-25.2)
Perceived great risk from smoking marijuana once or twice a week																								
Total	41.7	(37.8-45.8)	47.7	(43.1-52.4)	44.9	(41.1-48.6)	41.9	(38.6-45.2)	41.9	(38.1-45.8)	38.0	(33.0-43.3)	33.6	(29.8-37.7)	31.9	(27.8-36.4)	31.0	(27.2-35.1)	29.5	(25.2-33.0)	24.2	(20.2-28.6)	23.0	(19.6-26.8)
12-17	51.6	(46.7-56.4)	51.8	(46.7-57.0)	50.5	(45.4-55.5)	47.9	(43.6-52.2)	51.7	(47.2-56.1)	50.5	(45.9-55.2)	45.8	(41.6-50.1)	43.4	(38.8-48.1)	40.5	(35.8-45.4)	41.6	(35.8-47.6)	38.3	(33.0-44.0)	31.6	(26.5-37.1)
18-25	25.1	(20.9-29.9)	32.4	(27.2-38.1)	31.5	(26.9-36.5)	28.6	(25.1-32.3)	25.9	(22.3-29.9)	23.4	(20.1-27.0)	22.1	(18.4-26.3)	19.8	(16.0-24.2)	21.4	(17.9-25.3)	21.8	(18.3-25.9)	18.4	(14.7-22.8)	15.9	(12.4-20.2)
≥26	43.3	(38.6-48.1)	49.8	(44.2-55.5)	46.4	(41.7-51.2)	43.4	(39.4-47.5)	43.3	(38.6-48.0)	38.8	(32.7-45.3)	34.0	(29.3-39.0)	32.6	(27.5-38.1)	31.5	(26.8-36.6)	29.4	(25.4-33.7)	23.5	(19.2-27.8)	23.2	(19.2-27.8)
Perceived no risk from smoking marijuana once or twice a week																								
Total	8.6	(6.2-11.6)	6.9	(5.0-9.4)	6.5	(5.1-8.3)	7.3	(6.0-8.9)	8.2	(6.7-10.1)	9.7	(7.6-12.5)	11.9	(9.9-14.2)	12.1	(9.7-15.1)	14.5	(12.0-17.4)	16.3	(13.9-19.1)	16.9	(14.5-19.7)	17.2	(15.0-19.7)
12-17	4.9	(3.2-7.3)	5.4	(3.6-8.1)	6.1	(4.0-9.1)	5.7	(3.6-8.7)	6.5	(4.9-8.6)	6.8	(5.1-8.9)	8.1	(6.1-10.8)	12.1	(9.6-15.3)	11.8	(9.2-15.0)	12.1	(9.3-15.7)	16.0	(13.3-19.1)	15.7	(12.5-19.4)
18-25	15.8	(12.0-20.5)	14.7	(11.3-19.0)	17.3	(13.7-21.7)	14.6	(11.6-18.4)	11.8	(9.0-15.4)	15.9	(12.0-20.8)	19.0	(15.4-23.3)	21.4	(17.2-26.4)	26.9	(22.2-32.1)	28.3	(22.8-34.4)	26.6	(21.9-31.9)	31.8	(27.0-37.0)
≥26	7.8	(4.9-12.2)	5.7	(3.7-8.7)	4.8	(3.2-6.9)	6.3	(4.7-8.3)	7.9	(6.0-10.3)	9.1	(6.5-12.6)	11.2	(8.7-14.2)	10.7	(7.9-14.2)	12.8	(9.8-16.5)	14.9	(12.0-18.3)	15.5	(12.5-19.0)	15.1	(12.5-18.0)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 Perceived great risk and no risk from smoking marijuana is defined as those who reported that smoking marijuana once a month and once or twice a week might cause harm [great risk] and might not cause harm [no risk].

Table 7. Percentage of perceived parental disapproval of trying marijuana once or twice and using marijuana once a month or more among persons aged 12 to 17 — National Survey on Drug Use and Health, Oregon, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	%	%	%	%	%	%	%	%	%	%	%	%
	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)
Perceived parental disapproval of trying marijuana once or twice ¹												
12–17 years	93.7 (91.3-96.5)	94.7 (91.5-98.7)	95.0 (91.9-98.9)	95.1 (92.2-97.0)	93.6 (91.1-95.5)	92.7 (89.3-95.1)	93.7 (90.6-95.8)	93.6 (90.4-95.8)	94.4 (91.5-96.4)	95.2 (93.1-96.6)	94.7 (92.0-96.8)	94.2 (91.6-96.0)
Perceived parental disapproval of using marijuana once a month or more ²												
12–17 years	95.9 (93.7-97.4)	96.3 (94.4-97.6)	96.5 (93.6-98.1)	96.4 (93.0-98.2)	94.3 (91.6-96.2)	93.8 (90.5-96.9)	95.0 (92.1-96.9)	94.0 (90.8-96.1)	94.7 (91.8-96.7)	96.1 (93.9-97.5)	96.2 (94.3-97.5)	96.0 (94.0-97.4)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Perceived parental disapproval of trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of trying marijuana once or twice.

² Perceived parental disapproval of using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of using marijuana once a month or more.

Table 8. Percentage of perceived parental disapproval of trying marijuana once or twice and using marijuana once a month or more among past month marijuana users aged 12 to 17 — National Survey on Drug Use and Health, Oregon, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	%	%	%	%	%	%	%	%	%	%	%	%
	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)
Perceived parental disapproval of trying marijuana once or twice ¹												
12–17 years	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)
Perceived parental disapproval of using marijuana once a month or more ²												
12–17 years	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)	* (**)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Perceived parental disapproval of trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of trying marijuana once or twice.

² Perceived parental disapproval of using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of using marijuana once a month or more. Indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.

Table 9. Percentage of perceived attitudes towards peers trying marijuana once or twice and using marijuana once or more among persons aged 12 to 17— National Survey on Drug Use and Health, Oregon, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived attitudes towards peers trying marijuana once or twice ¹												
12–17 years	79.6 (75.2-83.3)	76.7 (72.2-80.6)	78.0 (73.4-82.0)	79.5 (75.3-83.1)	76.6 (72.3-80.4)	77.3 (72.2-81.6)	78.3 (74.4-81.7)	74.6 (70.7-78.0)	77.5 (73.7-80.9)	80.3 (76.2-83.8)	76.4 (72.4-80.0)	73.7 (69.5-77.5)
Perceived attitudes towards peers using marijuana once a month or more ²												
12–17 years	79.4 (74.8-83.3)	77.7 (73.1-81.8)	77.6 (72.4-82.1)	79.7 (74.2-84.3)	78.8 (74.7-82.4)	78.0 (73.1-82.2)	78.8 (74.9-82.2)	76.3 (72.5-79.7)	78.8 (75.1-82.2)	80.4 (76.2-84.0)	77.2 (73.1-80.9)	73.8 (69.8-77.5)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
Abbreviation: CI = confidence interval.
¹ Disapproving attitudes toward peers trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers trying marijuana once or twice.
² Disapproving attitudes toward peers using marijuana once or a month or more is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers using marijuana once a month or more.

Table 10. Percentage of perceived attitudes towards peers trying marijuana once or twice and using marijuana once a month or more among past month marijuana users aged 12 to 17— National Survey on Drug Use and Health, Oregon, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived attitudes towards peers trying marijuana once or twice ¹												
12–17 years	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)
Perceived attitudes towards peers using marijuana once a month or more ²												
12–17 years	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)	• (•-•)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
Abbreviation: CI = confidence interval.
¹ Disapproving attitudes toward peers trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers trying marijuana once or twice; indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.
² Disapproving attitudes toward peers using marijuana once or a month or more is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers using marijuana once a month or more; indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.

Table 11. Percentage of perceived availability among all persons aged ≥12 and mode of acquisition of marijuana among past year marijuana users aged ≥12 by age group — National Survey on Drug Use and Health, Oregon, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived availability												
Total	63.1 (58.2-67.8)	61.2 (56.0-66.2)	62.4 (58.0-66.6)	62.7 (59.0-66.2)	63.1 (59.6-66.5)	64.2 (59.8-68.3)	61.0 (56.5-65.3)	62.3 (58.0-66.4)	66.3 (62.7-69.7)	68.7 (63.8-73.3)	70.3 (65.7-74.6)	71.3 (67.4-74.9)
12-17	56.3 (51.3-61.2)	54.3 (49.0-59.4)	54.8 (50.3-59.3)	54.7 (50.6-59.8)	51.1 (45.9-56.3)	51.3 (46.1-56.3)	53.5 (48.9-58.0)	53.2 (49.0-57.3)	52.5 (48.2-56.7)	50.3 (45.2-55.5)	52.8 (48.1-57.4)	59.8 (54.9-64.6)
18-25	83.6 (78.7-87.5)	80.7 (76.0-84.7)	81.0 (77.3-84.2)	81.2 (77.6-84.3)	80.2 (75.3-84.3)	82.0 (76.6-86.4)	80.8 (76.3-84.5)	78.5 (74.5-82.0)	80.7 (76.7-84.1)	81.3 (76.3-85.5)	82.6 (77.8-86.5)	79.7 (74.7-83.9)
≥26	60.4 (54.6-66.0)	58.7 (52.4-64.7)	60.1 (54.7-65.3)	60.5 (56.2-64.6)	61.8 (57.6-65.9)	62.9 (57.7-67.8)	58.7 (53.1-64.1)	60.7 (55.3-65.9)	65.6 (61.2-69.7)	68.8 (62.6-74.4)	70.3 (64.4-75.5)	71.2 (66.2-75.7)
Mode of acquisition of marijuana												
Bought it												
Total	46.9 (39.3-54.5)	40.3 (32.8-48.4)	43.7 (37.5-50.2)	40.3 (33.9-47.0)	33.7 (26.0-42.4)	35.7 (27.1-45.4)	33.5 (26.5-41.4)	33.1 (26.5-40.4)	34.7 (27.0-43.4)	32.2 (26.5-38.5)	29.3 (22.6-37.0)	35.9 (29.2-43.1)
12-17	(*) (*)	38.7 (30.5-47.7)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	20.5 (12.9-31.2)	29.5 (21.5-39.0)	34.0 (25.4-43.7)	(*) (*)	(*) (*)	(*) (*)
18-25	46.6 (38.4-54.9)	43.6 (36.0-51.4)	39.1 (31.4-47.3)	35.8 (27.6-45.0)	31.5 (24.2-39.9)	34.6 (27.1-42.8)	37.3 (32.1-42.7)	39.6 (33.5-46.0)	42.7 (34.3-51.6)	41.9 (34.9-49.3)	38.8 (32.1-46.0)	44.4 (37.4-51.7)
≥26	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	28.0 (20.2-37.4)	26.7 (18.1-37.5)	33.8 (25.0-44.0)
Traded something for it												
Total	0.8 (0.3-1.9)	1.3 (0.6-2.8)	1.4 (0.5-3.9)	1.0 (0.2-4.1)	(*) (*)	2.1 (0.6-7.0)	0.6 (0.3-1.5)	1.2 (0.4-3.5)	1.6 (0.6-4.5)	0.9 (0.3-2.8)	2.5 (1.0-6.0)	3.2 (1.5-6.7)
12-17	(*) (*)	6.2 (2.5-14.5)	4.0 (1.5-10.6)	(*) (*)	(*) (*)	(*) (*)	3.6 (1.1-11.0)	1.2 (0.3-4.0)	(*) (*)	(*) (*)	(*) (*)	(*) (*)
18-25	0.8 (0.2-3.2)	1.3 (0.3-5.7)	(*) (*)	(*) (*)	2.1 (0.6-7.0)	1.7 (0.6-4.5)	0.7 (0.2-2.7)	1.8 (0.5-6.1)	1.6 (0.4-6.0)	0.6 (0.1-2.5)	1.8 (0.7-4.8)	2.0 (0.8-5.1)
≥26	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	0.9 (0.2-4.4)	2.6 (0.8-8.1)	3.4 (1.3-9.0)
Got it for free or shared with someone else												
Total	47.1 (39.9-54.5)	51.4 (43.0-59.7)	51.8 (44.8-58.8)	55.4 (47.7-62.8)	60.4 (50.8-69.2)	59.4 (50.2-67.9)	63.2 (55.7-70.1)	63.8 (56.5-70.4)	61.9 (53.5-69.6)	62.4 (55.1-69.1)	58.0 (47.8-67.5)	52.1 (44.0-60.1)
12-17	(*) (*)	54.2 (43.6-64.4)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	75.9 (64.7-84.3)	69.3 (60.2-77.2)	65.4 (56.1-73.5)	(*) (*)	(*) (*)	(*) (*)
18-25	52.5 (44.1-60.7)	54.3 (46.5-61.9)	58.7 (50.4-66.5)	61.5 (52.6-69.7)	65.7 (57.4-73.0)	62.8 (55.0-69.9)	61.0 (55.6-66.2)	58.6 (52.0-65.0)	54.5 (45.9-62.9)	56.1 (49.2-62.7)	57.5 (51.2-63.5)	52.1 (45.1-59.1)
≥26	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	64.8 (54.0-74.3)	(*) (*)	50.4 (39.1-61.7)
Grew it yourself												

Total	• (**)	• (**)	3.0 (1.2-7.1)	3.3 (1.2-9.0)	4.0 (1.5-9.7)	2.8 (1.0-7.4)	2.6 (1.0-6.9)	1.9 (0.5-6.4)	1.8 (0.5-6.0)	4.8 (1.7-11.4)	• (**)	• (**)
12-17	• (**)	• (**)	• (**)	1.1 (0.3-4.7)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)
18-25	0.2 (0.0-1.3)	• (**)	1.4 (0.4-4.4)	1.5 (0.5-5.1)	• (**)	1.0 (0.2-4.5)	1.0 (0.2-4.3)	• (**)	• (**)	• (**)	2.0 (0.7-5.2)	1.5 (0.5-4.6)
≥26	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)	• (**)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Abbreviation: CI = confidence interval.

* Perceived availability is defined as those who reported that it would be fairly easy or very easy for them to obtain marijuana if they wanted some.

† Mode of acquisition is defined as those who reported how they got the marijuana they used for the last time [i.e., bought it, traded something for it, got it for free or shared with someone else, or grew it yourself]; this question is only asked of those respondents indicating use of marijuana in the past 12 months.

Table 12. Percentage of past year marijuana dependence and abuse¹ among all persons aged ≥12 by age group — National Survey on Drug Use and Health, Oregon, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	1.6 (1.1-2.4)	2.1 (1.4-3.1)	2.1 (1.5-3.0)	1.6 (1.1-2.3)	1.4 (1.0-2.1)	1.9 (1.2-2.9)	1.9 (1.2-3.0)	2.1 (1.5-2.9)	2.3 (1.6-3.0)	2.1 (1.6-2.9)	2.4 (1.6-3.6)	2.3 (1.6-3.3)
12-17	2.9 (1.8-4.5)	3.7 (2.7-5.2)	4.5 (3.1-6.5)	4.7 (3.1-7.0)	4.0 (2.4-6.4)	3.3 (2.0-5.4)	3.6 (2.3-5.4)	5.5 (3.7-8.1)	5.5 (3.7-8.2)	3.6 (2.3-5.7)	4.3 (3.0-6.3)	4.3 (2.8-6.3)
18-25	7.4 (5.4-10.0)	5.9 (4.2-8.3)	6.8 (4.5-10.2)	6.4 (4.2-9.6)	5.6 (3.8-8.1)	7.0 (4.9-9.9)	7.2 (5.2-10.0)	6.4 (4.4-9.4)	7.1 (5.1-9.7)	7.9 (5.9-10.5)	5.4 (3.7-7.9)	5.7 (3.7-8.6)
≥26	0.5 (0.2-1.4)	1.2 (0.6-2.4)	1.0 (0.5-2.1)	0.4 (0.2-1.2)	0.4 (0.1-1.4)	0.9 (0.3-2.5)	0.9 (0.3-2.7)	1.0 (0.5-2.0)	1.2 (0.8-1.9)	1.1 (0.5-2.1)	1.7 (0.8-3.3)	1.5 (0.8-2.8)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
¹ Substance use disorder (also known as dependence or abuse) is defined as meeting criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association [APA], 1994) for either dependence or abuse for illicit drugs or alcohol.

Table 13. Percentage of past year marijuana dependence and abuse¹ among past year marijuana users² aged ≥12, by age group — National Survey on Drug Use and Health, Oregon, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	11.0 (7.7-15.6)	15.2 (11.3-20.2)	15.1 (10.7-20.9)	11.4 (7.7-16.5)	9.7 (6.7-14.0)	12.5 (8.4-18.2)	12.4 (8.3-18.1)	12.7 (9.3-17.2)	13.9 (10.7-17.9)	11.6 (8.6-15.6)	11.5 (8.0-16.3)	11.3 (8.1-15.6)
12-17	17.4 (11.2-26.1)	20.7 (15.4-27.2)	23.1 (15.1-33.8)	23.8 (15.4-35.0)	24.0 (15.5-35.2)	19.6 (12.0-30.4)	18.4 (12.4-26.4)	26.3 (18.9-35.4)	29.5 (21.2-39.5)	23.6 (15.8-33.7)	*	*
18-25	20.2 (14.6-27.2)	17.1 (12.2-23.5)	19.0 (12.7-27.6)	19.3 (12.9-27.8)	16.5 (11.7-22.7)	18.6 (13.3-25.3)	18.7 (13.3-25.7)	16.2 (11.0-23.0)	17.5 (12.8-23.5)	20.7 (16.6-25.5)	15.5 (11.1-21.3)	15.0 (9.7-22.5)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	6.8 (3.3-13.3)	8.9 (4.7-16.1)	8.7 (4.8-15.3)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
¹ Substance use disorder (also known as dependence or abuse) is defined as meeting criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association [APA], 1994) for either dependence or abuse for illicit drugs or alcohol.
² Analysis limited to past year marijuana users (defined as those who reported use of marijuana within 12 months preceding the date of interview, including those who reported use of marijuana within 30 days preceding the date of interview)

Table 14. Percentage of perception of maximum legal penalty in your state for first offense possession of an ounce or less of marijuana for your own use among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Oregon, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Fine												
Total	43.6 (38.0-49.3)	41.2 (36.7-45.8)	46.6 (42.0-51.2)	46.5 (42.0-51.1)	47.7 (43.0-52.5)	48.0 (43.8-52.2)	46.7 (42.2-51.2)	50.2 (45.2-55.1)	49.7 (45.3-54.1)	48.1 (43.4-52.9)	51.8 (47.5-56.1)	50.0 (45.7-54.3)
12-17	16.0 (12.1-20.9)	19.8 (16.0-24.4)	22.3 (17.6-27.8)	21.7 (16.5-28.0)	20.6 (16.1-26.0)	20.4 (15.9-25.8)	25.2 (20.4-30.8)	23.3 (18.4-29.0)	23.3 (19.0-28.2)	25.2 (20.6-30.4)	25.3 (21.0-30.1)	26.7 (22.3-31.5)
18-25	40.2 (35.5-45.1)	40.0 (35.3-44.9)	38.1 (32.9-43.6)	35.8 (31.3-40.6)	35.2 (30.2-40.6)	39.5 (32.9-46.6)	41.3 (36.4-46.4)	44.9 (39.7-50.1)	48.7 (42.9-54.6)	40.7 (36.4-45.2)	38.8 (33.5-44.4)	44.4 (39.2-49.7)
≥26	48.7 (41.0-56.5)	44.9 (38.8-51.1)	52.2 (46.1-58.2)	52.3 (46.0-58.5)	53.6 (47.2-59.9)	52.8 (47.3-58.3)	50.2 (44.5-55.9)	54.5 (48.4-60.5)	53.2 (47.5-58.8)	52.5 (46.2-58.7)	57.9 (51.9-63.6)	54.2 (48.5-59.8)
Probation												
Total	15.1 (12.2-18.6)	14.5 (11.4-18.2)	14.9 (12.0-18.3)	14.2 (10.9-18.4)	11.3 (8.8-14.4)	11.2 (8.6-14.4)	11.1 (8.7-13.9)	10.0 (7.8-12.6)	11.7 (9.7-14.1)	12.4 (10.1-15.2)	12.1 (9.6-15.2)	13.2 (10.5-16.4)
12-17	25.8 (21.6-30.4)	21.1 (17.4-25.4)	17.9 (14.3-22.3)	18.5 (14.5-23.4)	18.2 (13.5-24.0)	19.2 (15.1-24.2)	17.7 (14.5-21.4)	17.1 (13.3-21.7)	16.2 (14.9-22.0)	21.3 (17.6-25.5)	25.3 (20.5-30.9)	23.6 (19.4-28.3)
18-25	13.1 (9.7-17.5)	13.7 (9.9-18.7)	11.8 (8.7-15.7)	14.0 (11.1-17.5)	14.1 (11.4-17.4)	12.7 (10.2-15.7)	15.2 (12.1-18.9)	13.1 (10.3-16.6)	11.8 (8.9-15.4)	16.2 (12.5-20.6)	17.8 (13.9-22.4)	14.8 (11.0-19.6)
≥26 years	13.9 (10.2-18.6)	13.6 (9.7-18.7)	15.1 (11.4-19.7)	13.6 (9.4-19.4)	9.8 (6.8-14.0)	9.9 (6.8-14.2)	9.5 (6.6-13.6)	8.4 (5.9-12.0)	10.9 (8.3-14.1)	10.6 (7.8-14.1)	9.3 (6.5-13.0)	11.5 (8.2-15.9)
Community Service												

Total	9.0 (6.1-13.3)	10.4 (7.1-15.0)	7.2 (5.4-9.4)	6.2 (4.5-8.6)	7.4 (5.3-10.2)	7.4 (5.6-9.6)	7.4 (5.6-9.7)	7.2 (4.9-10.6)	6.8 (5.0-9.4)	7.7 (6.1-9.6)	6.7 (5.2-8.5)	6.5 (4.8-8.8)
12-17	19.4 (15.2-24.5)	18.6 (14.9-22.9)	14.5 (11.1-18.7)	13.2 (9.1-18.7)	15.1 (11.1-20.1)	13.0 (9.4-17.8)	11.4 (9.0-14.4)	13.1 (10.0-16.9)	14.6 (11.0-19.2)	13.5 (10.4-17.2)	11.0 (7.3-16.3)	11.4 (7.9-16.2)
18-25	10.1 (7.5-13.7)	9.7 (6.8-13.5)	6.8 (4.9-9.3)	4.7 (3.0-7.2)	7.4 (5.0-10.8)	10.6 (8.1-13.9)	9.0 (6.4-12.5)	7.9 (5.3-11.6)	8.4 (5.5-12.7)	9.1 (6.7-12.1)	8.6 (6.5-11.2)	9.4 (6.7-12.9)
≥26	7.2 (3.6-13.9)	9.3 (5.3-16.0)	6.1 (4.0-9.2)	5.6 (3.5-8.9)	6.4 (3.8-10.5)	6.1 (4.1-9.0)	6.6 (4.4-9.9)	6.4 (3.6-11.0)	5.6 (3.4-9.0)	6.7 (4.8-9.4)	5.7 (4.0-8.1)	5.4 (3.4-8.3)
Possible Prison Sentence												
Total	27.1 (23.1-31.5)	28.3 (24.9-32.0)	27.4 (23.9-31.2)	24.7 (21.7-28.0)	22.3 (18.9-26.1)	22.3 (18.8-26.2)	23.2 (19.5-27.4)	21.6 (18.6-25.0)	19.9 (16.7-23.6)	19.9 (16.6-23.6)	19.3 (16.3-22.6)	18.8 (16.3-21.6)
12-17	31.1 (26.2-36.5)	33.0 (28.8-37.5)	37.2 (32.0-42.7)	36.2 (30.6-42.3)	31.5 (26.8-36.6)	31.6 (26.1-37.7)	30.8 (25.6-36.5)	29.8 (25.8-34.0)	29.8 (25.6-34.3)	29.0 (24.1-34.4)	25.0 (20.8-29.8)	25.4 (21.2-30.0)
18-25	30.1 (25.6-35.0)	33.2 (28.7-38.1)	36.7 (31.8-41.8)	35.4 (30.7-40.4)	32.7 (28.2-37.7)	29.6 (24.6-35.0)	27.9 (23.5-32.7)	27.5 (22.9-32.6)	24.3 (20.2-29.0)	26.1 (22.8-29.6)	26.7 (21.9-32.1)	22.9 (18.1-28.5)
≥26	25.8 (20.6-31.7)	26.5 (21.9-31.7)	23.8 (19.4-29.0)	20.9 (16.9-25.4)	19.1 (14.8-24.3)	19.9 (15.6-24.9)	21.4 (17.0-26.6)	19.5 (15.7-23.8)	17.9 (13.8-22.8)	17.5 (13.4-22.6)	17.0 (13.4-21.5)	17.1 (14.0-20.7)
Mandatory Prison Sentence												
Total	5.1 (3.4-7.8)	5.5 (3.5-8.6)	4.0 (2.8-5.8)	4.3 (3.1-5.9)	3.8 (2.7-5.3)	4.1 (2.8-6.8)	5.0 (3.8-6.5)	4.3 (3.1-5.9)	3.9 (2.5-6.1)	3.6 (2.1-5.9)	2.5 (1.6-4.0)	2.6 (1.6-4.1)
12-17	7.7 (5.1-11.4)	7.5 (5.3-10.4)	8.1 (5.8-11.2)	8.9 (6.3-12.4)	10.6 (7.1-15.5)	11.7 (8.3-16.2)	13.1 (10.5-16.2)	13.7 (10.5-17.6)	11.2 (8.3-14.8)	8.9 (6.1-12.7)	7.8 (4.7-12.5)	7.1 (4.8-10.5)
18-25	6.5 (3.7-11.0)	3.4 (2.1-5.3)	6.7 (4.3-10.3)	8.9 (6.0-13.1)	7.9 (5.1-12.0)	4.5 (3.0-6.8)	4.1 (2.6-6.4)	4.5 (2.8-7.2)	4.0 (2.5-6.4)	4.5 (3.2-6.2)	3.8 (2.5-5.8)	3.8 (2.2-6.6)

≥26	4.4 (2.4-8.1)	5.7 (3.1-10.2)	2.8 (1.5-5.5)	2.7 (1.4-4.9)	2.2 (1.1-4.3)	3.0 (1.7-5.5)	4.1 (2.7-6.3)	3.1 (1.9-5.3)	3.0 (1.4-6.2)	2.7 (1.2-6.4)	1.6 (0.6-4.0)	1.7 (0.7-4.3)
No penalty												
Total	DNA	DNA	DNA	4.0 (2.5-6.4)	7.5 (5.3-10.6)	7.1 (4.4-11.5)	6.7 (4.1-10.9)	6.7 (4.2-10.5)	7.9 (5.2-11.7)	8.3 (6.1-11.3)	7.6 (5.2-10.9)	8.9 (6.6-11.9)
12-17	DNA	DNA	DNA	1.4 (0.5-3.6)	4.1 (2.1-7.9)	3.9 (2.2-7.0)	1.7 (0.9-3.4)	3.1 (1.9-5.2)	3.0 (1.8-4.8)	2.3 (1.0-5.3)	5.5 (3.2-9.3)	5.9 (3.6-9.4)
18-25	DNA	DNA	DNA	1.2 (0.4-3.3)	2.7 (1.3-5.6)	3.0 (1.8-5.2)	2.6 (1.3-5.0)	2.1 (1.1-4.1)	2.7 (1.4-5.0)	3.5 (2.0-6.0)	4.3 (1.8-9.8)	4.8 (2.4-9.6)
≥26	DNA	DNA	DNA	5.0 (3.0-8.1)	8.9 (6.1-13.0)	8.3 (4.8-13.9)	8.1 (4.7-13.6)	8.1 (4.9-13.0)	9.5 (6.1-14.4)	10.0 (7.1-13.8)	8.5 (5.6-12.9)	10.1 (7.2-14.1)
Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.												
Abbreviations: CI = confidence interval; DNA = data not available.												
*Perception of legal penalty for marijuana possession is defined as those who reported their perception of the maximum legal penalty in their state of residence for first offense possession of an ounce or less of marijuana for their own use.												

Supplementary Material

Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, Washington, 2002–2014

Suggested Citation: Alejandro Azofeifa, Margaret E. Mattson, and Rob Lyster. Supplementary Material. State Level Data: Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, Washington, 2002–2014. Center for Behavioral Health Statistics and Quality. (2016). Substance Abuse and Mental Health Services Administration, Rockville, MD.

Notes: A) Two years of data were combined to allow for reliable state-level prevalence estimates. Some state-level prevalence estimates for combined years have been suppressed (indicated by an asterisk (*) symbol in each cell) based on the following National Survey on Drug Use and Health (NSDUH) suppression criteria. Prevalence estimates are suppressed if any of the following occurred: 1) the prevalence estimate is $<0.005\%$ or $>99.995\%$; 2) the relative standard error (RSE) of the negative natural logarithm of the estimated proportion p (where p is the prevalence divided by 100) is >0.175 if the prevalence is $>50\%$; 3) the RSE of the negative natural logarithm of $(1-p)$ is >0.175 if the prevalence is $>50\%$; 4) the actual sample size is <100 ; or 5) the effective sample size (defined as the actual sample size divided by the design effect) is <68 .
B) The estimates provided in this supplementary material for each state and the District of Columbia are from design-based direct estimation procedures (i.e., they are based directly on the respondent sample and the accompanying analytical weights with variances derived through a Taylor linearization approach using SUDAAN software) as used for the national level estimates (available at: Azofeifa A, Mattson ME, Schauer G, McAftee T, Grant A, Lyster R. National Estimates of Marijuana Use and Related Indicators — National Survey on Drug Use and Health, United States, 2002–2014. MMWR Surveill Summ 2016;65(No. SS-1):1–25. DOI: <http://dx.doi.org/10.15585/mmwr.ss6511a1>). Caution is advised when comparing data from other NSDUH reports that are based on different design-based estimation procedures.
C) Interpretation of state-level trend data should be done with caution as some prevalence estimates are suppressed based on the above criteria and survey cycle differences may not be statistically significant.

BILAG 5

Table 1. Percentage of past month marijuana use* among all persons aged ≥ 12, by selected demographic characteristics — National Survey on Drug Use and Health, Washington, 2002–2014

Characteristics Age group (yrs)	2002–2003	2003–2004	2004–2005	2005–2006	2006–2007	2007–2008	2008–2009	2009–2010	2010–2011	2011–2012	2012–2013	2013–2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	7.1 (5.6–9.1)	5.8 (4.5–7.5)	7.2 (5.7–9.1)	8.7 (6.8–11.1)	8.0 (6.1–10.3)	8.0 (6.0–10.7)	7.6 (5.8–9.9)	9.3 (7.2–11.8)	10.2 (8.0–12.8)	10.7 (8.6–13.2)	13.0 (10.9–15.5)	13.3 (11.2–15.7)
12–17	9.4 (6.9–12.8)	8.4 (5.8–11.9)	6.3 (4.4–8.9)	7.7 (5.8–10.1)	6.8 (5.2–8.8)	6.7 (4.7–9.6)	8.4 (6.2–11.3)	7.6 (5.5–10.5)	9.6 (6.7–13.6)	9.4 (6.5–13.6)	8.9 (6.7–11.6)	10.2 (7.6–13.6)
18–25	22.1 (18.1–26.7)	17.3 (13.5–21.8)	18.2 (14.3–22.7)	20.3 (16.5–24.6)	19.0 (15.3–23.4)	16.9 (13.7–20.5)	17.0 (13.8–20.7)	24.2 (19.7–29.4)	27.1 (22.5–32.3)	23.8 (20.5–27.4)	25.3 (21.7–29.2)	22.6 (18.2–27.7)
≥26	4.2 (2.8–6.3)	3.5 (2.2–5.4)	5.4 (3.9–7.5)	6.8 (4.8–9.7)	6.3 (4.2–9.3)	6.8 (4.4–10.2)	6.0 (3.9–9.1)	7.1 (4.8–10.2)	7.4 (5.2–10.6)	8.6 (6.4–11.5)	11.5 (9.1–14.5)	12.1 (9.7–15.1)
26–34	7.8 (4.3–13.7)	8.5 (4.5–15.4)	9.8 (5.2–17.6)	9.7 (5.3–17.1)	11.7 (7.2–18.4)	9.3 (5.3–15.6)	7.0 (3.3–14.1)	14.2 (8.5–22.6)	14.4 (9.0–22.2)	17.4 (13.5–22.1)	28.8 (22.4–36.2)	24.4 (18.4–31.7)
35–44	9.2 (5.1–16.0)	6.1 (3.3–11.1)	8.5 (5.3–13.5)	9.9 (6.4–14.9)	6.5 (3.5–11.8)	7.3 (4.5–11.7)	5.8 (3.1–10.4)	5.5 (3.3–8.9)	8.0 (4.5–14.1)	12.4 (7.5–19.6)	14.9 (9.7–22.3)	11.3 (7.6–16.6)
45–54	2.2 (0.7–6.3)	1.5 (0.4–5.6)	6.7 (3.5–12.5)	10.1 (5.6–17.8)	8.7 (4.2–17.1)	8.7 (4.2–17.1)	9.8 (4.7–19.5)	8.7 (4.8–15.2)	6.7 (2.8–15.2)	7.5 (3.7–14.6)	7.0 (3.6–13.1)	8.8 (5.0–15.1)
55–64	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	15.8 (9.6–24.7)
≥65	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Sex												
Male	8.8 (6.8–11.4)	7.3 (5.3–9.8)	9.9 (7.4–13.1)	12.3 (9.2–16.3)	11.7 (8.8–15.4)	10.4 (7.8–13.8)	8.7 (6.5–11.6)	11.5 (8.1–16.2)	12.4 (9.3–16.2)	12.8 (9.9–16.3)	16.2 (12.8–20.2)	16.9 (14.0–20.2)
Female	5.4 (3.9–7.6)	4.4 (3.1–6.1)	4.6 (3.3–6.2)	5.2 (3.5–7.5)	4.4 (2.8–6.7)	5.7 (3.2–10.0)	6.5 (4.0–10.4)	7.1 (5.2–9.7)	8.0 (5.9–10.9)	9.6 (6.4–11.5)	10.0 (8.0–12.5)	9.8 (7.4–12.7)
Race/ethnicity												
Non-Hispanic white	7.2 (5.5–9.4)	5.9 (4.4–7.9)	7.0 (5.5–9.8)	8.9 (6.8–11.5)	8.3 (6.1–11.1)	7.4 (5.7–9.5)	7.4 (5.8–9.5)	9.9 (7.4–13.1)	10.1 (7.5–13.4)	10.6 (8.1–13.9)	13.8 (11.3–16.6)	14.8 (12.2–17.9)
Non-Hispanic black	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic American Indian/Alaska Native	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic Hawaiian/Other Pacific Islander	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Non-Hispanic Asian	2.7 (1.4–5.2)	2.5 (1.3–4.7)	(*)	(*)	2.6 (0.8–8.0)	(*)	(*)	(*)	(*)	5.5 (2.5–11.6)	4.0 (2.1–7.7)	2.8 (1.3–5.7)
Non-Hispanic two or more races	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Hispanic	(*)	4.4 (1.7–11.0)	3.0 (1.3–7.1)	4.4 (2.3–8.2)	7.8 (3.7–15.4)	7.9 (3.7–16.1)	4.3 (2.4–7.3)	4.4 (2.7–7.1)	7.1 (4.6–10.7)	9.3 (5.1–16.4)	7.9 (4.2–14.5)	7.7 (5.0–11.7)
Current employment status [†]												

Full-time	6.3 (4.3-9.1)	5.2 (3.1-8.4)	8.6 (6.3-11.7)	10.2 (7.1-14.4)	9.5 (6.6-13.5)	7.4 (5.1-10.5)	6.1 (4.1-9.1)	10.3 (7.3-14.4)	10.6 (7.3-15.3)	9.4 (6.8-12.8)	13.6 (10.0-18.3)	14.7 (11.3-19.0)
Part-time	12.9 (8.5-19.1)	7.3 (4.4-12.0)	5.9 (3.5-9.7)	6.2 (3.6-10.5)	5.6 (3.1-9.9)	(*)	(*)	11.0 (6.6-17.6)	15.3 (9.7-23.2)	16.5 (11.0-24.2)	13.6 (9.3-19.4)	11.3 (7.6-16.5)
Unemployed	(*)	(*)	(*)	(*)	(*)	(*)	(*)	14.0 (8.4-22.4)	14.3 (8.4-23.1)	23.2 (15.6-33.1)	30.6 (21.9-40.9)	(*)
Other ²	3.1 (1.6-6.3)	4.3 (2.4-7.8)	5.6 (3.4-9.3)	6.9 (3.7-12.5)	6.5 (3.8-11.0)	6.6 (3.7-11.5)	5.8 (2.9-10.9)	6.6 (3.9-11.0)	6.7 (4.2-10.4)	7.8 (4.5-13.2)	9.7 (6.2-14.7)	10.4 (7.1-15.2)
Highest level of education completed ¹												
< High school	18.5 (12.9-25.7)	13.1 (8.7-19.2)	13.9 (8.6-21.7)	15.3 (10.1-22.5)	10.9 (7.1-16.2)	12.3 (6.8-21.3)	15.5 (9.2-24.9)	17.0 (11.6-24.2)	19.5 (12.0-30.2)	24.2 (16.7-33.7)	21.9 (14.8-31.2)	21.0 (14.7-29.1)
High school graduate	6.5 (4.0-10.3)	5.6 (3.1-9.8)	7.0 (4.2-11.4)	9.7 (6.0-15.2)	11.0 (6.4-18.4)	12.3 (7.0-20.9)	9.7 (5.5-16.6)	7.7 (5.0-11.9)	6.8 (4.7-9.7)	7.3 (4.8-11.1)	10.4 (7.5-14.3)	14.5 (10.8-19.1)
Some college	6.8 (4.0-11.2)	5.7 (3.3-9.7)	7.8 (5.1-11.7)	8.7 (5.5-13.5)	9.6 (6.6-13.8)	8.5 (5.7-12.6)	6.1 (3.6-10.1)	13.5 (8.8-20.1)	14.7 (10.0-21.0)	12.4 (9.0-16.8)	15.3 (11.1-20.6)	13.2 (9.8-17.4)
College graduate	1.5 (0.6-3.6)	1.5 (0.7-3.5)	3.8 (2.0-7.1)	5.2 (2.7-9.7)	3.7 (2.0-6.6)	3.3 (1.4-7.9)	3.8 (1.8-8.1)	3.0 (1.6-5.8)	4.9 (2.7-8.7)	8.2 (5.2-12.6)	11.8 (8.1-16.9)	10.8 (7.3-15.9)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Abbreviation: CI = confidence interval.

* Past month use of marijuana is defined as those who reported use of marijuana within 30 days preceding the date of interview.

¹ Current employment and highest level of education completed are only defined for 18 or older.

² "Other" employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Table 2. Percentage of past year marijuana use^a among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Washington, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	13.6 (11.2-16.6)	11.9 (9.9-14.2)	11.7 (9.8-13.8)	12.7 (10.4-15.5)	12.5 (10.0-15.4)	13.3 (10.6-16.4)	13.3 (10.9-16.2)	14.3 (11.8-17.3)	15.6 (12.9-18.7)	16.2 (13.5-19.3)	18.5 (15.7-21.6)	19.4 (16.9-22.3)
12-17	17.4 (14.0-21.3)	15.2 (12.0-19.0)	11.4 (8.7-14.8)	13.9 (11.0-17.5)	13.7 (11.4-16.3)	13.5 (10.9-16.6)	15.2 (12.3-19.5)	14.4 (11.8-17.4)	16.9 (13.5-21.0)	16.0 (12.7-20.0)	15.5 (12.4-19.2)	17.4 (13.8-21.7)
18-25	36.5 (31.8-41.6)	28.7 (23.3-34.8)	29.5 (24.3-35.4)	32.7 (27.2-38.6)	31.4 (26.1-37.2)	29.8 (25.4-34.5)	33.8 (29.6-38.4)	39.9 (34.4-45.7)	40.5 (34.6-46.7)	37.2 (32.4-42.3)	37.9 (33.0-43.1)	34.3 (29.2-39.8)
≥26	9.2 (6.8-12.3)	8.5 (6.2-11.5)	8.6 (6.6-11.1)	9.2 (6.9-12.2)	9.2 (6.6-12.6)	10.6 (7.7-14.4)	9.8 (7.0-13.5)	10.1 (7.4-13.7)	11.3 (8.6-14.7)	12.7 (9.7-16.5)	15.7 (12.4-19.6)	17.3 (14.2-20.8)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

Past year use of marijuana is defined as those who reported use of marijuana within 12 months preceding the date of interview; – these also include those who reported use of marijuana within 30 days preceding the date of interview.

Table 3. Percentage of daily or almost daily marijuana use in the past year and past month among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Washington, 2002–2014

Age group (yrs)	2007-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Daily or almost daily marijuana use in the past year ¹												
Total	2.0 (1.3-2.9)	1.5 (1.0-2.4)	1.7 (1.1-2.8)	2.1 (1.3-3.5)	1.9 (1.1-3.1)	1.6 (1.0-2.6)	1.6 (1.0-2.6)	2.0 (1.2-3.4)	2.4 (1.6-3.7)	3.2 (2.1-4.8)	4.2 (2.9-6.2)	4.7 (3.4-6.5)
12-17	1.6 (0.7-3.4)	0.7 (0.2-1.6)	0.7 (0.2-2.2)	0.7 (0.3-1.6)	0.9 (0.4-1.9)	0.9 (0.4-2.4)	1.3 (0.5-3.1)	0.9 (0.4-1.9)	1.0 (0.5-2.1)	1.1 (0.6-2.2)	1.6 (0.8-3.2)	1.8 (0.9-3.5)
18-25	6.1 (4.0-9.3)	6.1 (3.9-9.3)	5.0 (3.3-7.6)	4.6 (3.1-6.9)	4.4 (2.7-7.0)	4.9 (3.0-7.8)	4.4 (2.9-6.8)	5.4 (3.4-8.5)	8.0 (5.6-11.4)	7.4 (5.4-10.0)	7.5 (5.4-10.4)	7.9 (5.6-10.9)
≥26	1.3 (0.7-2.5)	0.7 (0.3-1.7)	1.3 (0.6-2.9)	1.9 (0.9-3.8)	1.6 (0.7-3.4)	1.1 (0.5-2.5)	1.2 (0.5-2.6)	1.6 (0.7-3.4)	1.7 (0.8-3.3)	2.8 (1.5-5.0)	4.0 (2.5-6.5)	4.5 (3.0-6.8)
Daily or almost daily marijuana use in the past month ²												
Total	2.4 (1.7-3.5)	1.9 (1.2-3.0)	2.5 (1.5-3.8)	3.3 (2.3-4.8)	3.2 (2.1-4.8)	2.9 (2.0-4.3)	2.8 (1.9-4.0)	4.0 (2.7-5.9)	4.5 (3.2-6.4)	4.9 (3.4-6.9)	5.7 (4.1-7.9)	6.0 (4.5-7.8)
12-17	2.1 (1.1-4.1)	1.8 (0.8-4.1)	1.6 (0.8-3.1)	1.5 (0.8-2.8)	1.1 (0.6-2.3)	1.4 (0.6-3.3)	2.1 (1.2-3.9)	2.1 (1.2-3.9)	2.5 (1.7-3.8)	2.8 (1.7-4.6)	3.0 (1.7-5.2)	2.3 (1.3-4.1)
18-25	8.6 (6.1-11.9)	6.5 (4.1-10.2)	5.5 (3.4-8.6)	7.4 (5.4-10.1)	8.1 (5.7-11.3)	6.6 (4.5-9.6)	6.5 (4.6-9.2)	7.8 (5.4-11.2)	10.1 (7.3-13.8)	10.7 (8.4-13.6)	10.6 (8.0-14.0)	9.7 (6.8-13.6)
≥26	1.4 (0.8-2.6)	1.2 (0.5-2.5)	2.1 (1.1-3.8)	2.9 (1.6-5.0)	2.6 (1.4-4.8)	2.5 (1.5-4.2)	2.3 (1.3-3.9)	3.6 (2.1-6.0)	3.9 (2.4-6.2)	4.2 (2.5-6.9)	5.2 (3.4-7.9)	5.8 (4.2-8.0)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Past year daily or almost daily use is defined as having used marijuana on 300 or more days in the past year.

² Past month daily or almost daily use is defined as having used marijuana on 20 or more days in the past month.

Table 4. Percentage of daily or almost daily marijuana use in the past year and past month among marijuana users aged ≥12, by age group — National Survey on Drug Use and Health, Washington, 2002–2014

Age group (yrs)	2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014	
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Daily or almost daily marijuana use in the past year																								
Total	14.5 (10.2-20.3)	12.9 (8.6-18.8)	14.9 (9.6-22.5)	16.7 (10.7-25.2)	15.0 (9.5-22.7)	12.0 (7.5-18.7)	12.1 (7.7-18.6)	14.1 (9.0-21.4)	15.5 (10.4-22.6)	19.8 (13.8-27.7)	22.9 (16.4-31.1)	24.1 (17.8-31.8)												
12-17	9.0 (4.1-18.8)	10.4 (5.0-20.2)	17.1 (11.0-25.6)	14.2 (8.8-22.1)	13.9 (8.4-22.1)	16.4 (10.1-25.5)	13.1 (8.4-20.0)	13.6 (9.0-20.1)	19.7 (14.0-27.1)	19.8 (14.7-26.2)	19.8 (14.7-26.2)	23.0 (17.0-30.4)												
18-25	16.8 (11.5-23.8)	21.2 (13.6-31.4)	17.1 (11.0-25.6)	14.2 (8.8-22.1)	13.9 (8.4-22.1)	16.4 (10.1-25.5)	13.1 (8.4-20.0)	13.6 (9.0-20.1)	19.7 (14.0-27.1)	19.8 (14.7-26.2)	19.8 (14.7-26.2)	23.0 (17.0-30.4)												
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)												
Daily or almost daily marijuana use in the past month																								
Total	34.2 (25.7-43.9)	33.4 (24.0-44.4)	34.4 (24.2-46.4)	38.4 (28.3-49.7)	39.6 (28.9-51.4)	42.8 (32.5-53.7)	44.6 (33.8-56.0)	46.0 (35.5-56.7)	43.7 (33.7-54.2)	45.0 (35.9-54.5)														
12-17	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)												
18-25	38.9 (29.8-48.8)	36.7 (27.5-46.9)	36.7 (27.5-46.9)	36.7 (27.5-46.9)	36.7 (27.5-46.9)	36.7 (27.5-46.9)	36.7 (27.5-46.9)	36.7 (27.5-46.9)	36.7 (27.5-46.9)	36.7 (27.5-46.9)	36.7 (27.5-46.9)	36.7 (27.5-46.9)												
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)												

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Analysis limited to marijuana users (i.e., estimate of daily or almost daily use in the past year is among past year marijuana users; estimate of daily or almost daily use in the past month is among past month marijuana users).
 † Past year daily or almost daily use is defined as having used marijuana on 300 or more days in the past year.
 ‡ Past month daily or almost daily use is defined as having used marijuana on 20 or more days in the past month.

Table 5. Percentage of past year marijuana initiation among persons aged ≥12 years at risk for initiation, estimated number at risk for initiation, and mean age at first use of marijuana among past year marijuana initiates, by age group — National Survey on Drug Use and Health, Washington, 2002–2014

Age group (yrs)	2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014	
	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Past Year marijuana initiation among persons at risk for initiation¹																								
Total	1.7	(1.1-2.6)	1.3	(0.8-1.8)	1.4	(1.0-2.0)	1.7	(1.2-2.5)	1.8	(1.1-3.0)	2.2	(1.3-3.4)	2.0	(1.5-2.6)	1.8	(1.3-2.6)	2.0	(1.4-3.0)	1.8	(1.3-2.5)	1.9	(1.5-2.5)	2.4	(1.8-3.2)
12-17	6.8	(4.3-10.5)	6.0	(4.2-8.7)	4.5	(2.9-7.0)	4.8	(3.1-7.5)	5.0	(3.3-7.4)	5.4	(3.6-7.9)	5.9	(4.1-8.3)	4.9	(3.4-6.8)	6.8	(4.6-10.0)	8.1	(5.8-11.4)	7.7	(5.7-10.3)	7.6	(5.2-11.1)
18-25	3.0	(1.8-4.9)	3.0	(1.2-7.2)	6.2	(3.3-11.3)	8.1	(4.8-13.3)	5.8	(3.4-9.9)	6.6	(4.1-10.5)	8.6	(5.5-13.2)	8.5	(5.7-12.5)	6.5	(4.4-9.5)	4.6	(2.6-7.8)	5.8	(3.7-9.1)	7.3	(4.4-11.8)
≥26	0.3	(0.0-2.1)	0.3	(0.0-1.3)	0.2	(0.0-1.3)	0.2	(0.0-1.3)	0.7	(0.1-3.3)	0.7	(0.1-3.3)	0.3	(0.0-1.3)	0.3	(0.0-2.2)	0.3	(0.0-1.3)	0.3	(0.0-1.3)	0.3	(0.0-1.3)	0.3	(0.1-1.3)
Estimated no. (in 1,000s) of past year marijuana initiation among those at risk for initiation¹																								
Total	42	(28-64)	36	(25-50)	41	(28-59)	48	(33-69)	52	(32-86)	60	(38-95)	56	(43-72)	52	(37-74)	57	(38-84)	53	(37-74)	55	(42-72)	67	(50-90)
12-17	30	(19-46)	27	(19-39)	21	(14-33)	22	(14-34)	23	(15-34)	25	(17-37)	26	(18-37)	21	(15-30)	30	(20-45)	38	(27-52)	36	(26-48)	35	(24-50)
18-25	8	(5-12)	9	(4-21)	20	(11-36)	26	(15-42)	19	(11-33)	21	(13-34)	26	(16-40)	25	(17-36)	20	(13-29)	15	(9-26)	19	(12-30)	25	(15-41)
≥26	5	(1-38)	0	(0-1)	0	(0-1)	0	(0-1)	14	(3-66)	14	(3-66)	4	(0-26)	6	(1-47)	0	(0-1)	0	(0-1)	0	(0-1)	7	(2-27)
Mean age at first use among past year marijuana initiates																								
Total	16.8	(14.4-19.3)	15.5	(14.7-16.3)	16.7	(15.8-17.6)	17.0	(16.1-17.9)	22.4	(12.4-32.4)	22.3	(13.5-31.0)	17.3	(16.0-22.1)	19.1	(16.4-21.5)	18.4	(15.4-21.5)	16.1	(15.3-16.9)	16.3	(15.3-17.3)	18.7	(16.1-21.3)
12-17	14.7	(14.2-15.1)	14.6	(14.0-15.1)	14.7	(14.2-15.3)	14.9	(14.1-15.6)	14.7	(14.0-15.5)	14.6	(14.1-15.1)	14.8	(14.5-15.6)	15.1	(14.5-15.6)	14.8	(14.3-15.4)	14.7	(14.3-15.2)	14.6	(13.8-15.3)	15.0	(14.3-15.7)
18-25	18.8	(17.8-19.9)	18.8	(17.8-19.9)	18.8	(17.8-19.9)	18.8	(17.8-19.9)	18.3	(17.7-18.9)	18.7	(18.1-19.4)	18.9	(18.3-19.4)	19.4	(18.3-20.5)	19.4	(18.3-21.3)	19.5	(18.4-20.7)	19.5	(18.3-20.7)	19.3	(18.3-20.3)
≥26	0.3	(0.0-2.1)	0.3	(0.0-1.3)	0.2	(0.0-1.3)	0.2	(0.0-1.3)	0.7	(0.1-3.3)	0.7	(0.1-3.3)	0.3	(0.0-1.3)	0.3	(0.0-2.2)	0.3	(0.0-1.3)	0.3	(0.0-1.3)	0.3	(0.0-1.3)	0.3	(0.1-1.3)
Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.																								
Abbreviation: CI = confidence interval.																								
¹ Past year marijuana initiation (initiates) among persons at risk for initiation is defined as those having first used marijuana within the 12-month period before the date of the interview.																								

Table 6. Percentage of perceived great risk and no risk from smoking marijuana once a month and once or twice a week among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Washington, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	%	%	%	%	%	%	%	%	%	%	%	%
	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)
Perceived great risk from smoking marijuana once a month												
Total	25.5	23.6	27.6	25.4	26.0	24.8	22.9	23.3	21.2	17.5	17.7	17.7
	(21.7-29.7)	(20.5-27.0)	(23.5-32.1)	(22.1-28.9)	(22.0-30.5)	(20.9-29.1)	(19.8-26.2)	(20.0-27.0)	(17.6-25.3)	(14.2-21.3)	(15.0-20.7)	(15.0-20.7)
12-17	26.7	30.2	30.1	27.0	27.0	29.4	30.2	25.7	22.7	18.4	13.8	13.8
	(22.9-30.8)	(26.3-34.3)	(25.5-35.1)	(23.8-32.0)	(22.2-32.4)	(25.5-33.6)	(25.9-34.9)	(22.3-29.5)	(18.5-27.6)	(14.6-22.8)	(10.8-17.3)	(10.8-17.3)
18-25	16.1	17.5	14.6	19.5	21.1	16.6	12.8	11.2	13.8	12.1	9.8	9.8
	(12.5-20.5)	(13.5-22.4)	(11.2-18.9)	(16.5-23.0)	(17.8-24.8)	(13.4-20.3)	(10.0-16.3)	(8.5-14.6)	(11.0-17.3)	(9.2-15.6)	(6.9-13.6)	(6.9-13.6)
≥26	26.9	23.7	29.5	26.0	26.7	25.6	22.2	22.2	22.2	18.3	19.4	19.4
	(22.2-32.2)	(19.8-28.2)	(24.6-34.9)	(22.1-30.4)	(21.8-32.2)	(20.8-31.0)	(19.9-27.9)	(21.0-29.6)	(17.8-27.5)	(14.4-23.0)	(16.2-23.1)	(16.2-23.1)
Perceived no risk from smoking marijuana once a month												
Total	11.9	12.6	13.3	13.5	14.6	16.6	18.1	20.7	22.0	23.8	23.8	23.8
	(9.5-14.8)	(10.0-15.8)	(10.6-16.6)	(11.1-16.3)	(11.9-17.9)	(13.9-19.7)	(15.9-20.5)	(17.8-24.0)	(18.7-25.6)	(21.2-26.7)	(21.4-26.3)	(21.4-26.3)
12-17	7.2	7.6	7.5	11.4	12.8	14.6	14.0	14.5	15.4	16.9	20.9	20.9
	(5.2-9.9)	(5.7-10.1)	(5.2-10.8)	(8.2-15.7)	(9.4-17.2)	(11.5-18.3)	(11.3-17.1)	(11.5-18.1)	(12.5-18.8)	(13.4-21.0)	(17.0-25.5)	(17.0-25.5)
18-25	23.2	19.3	21.9	19.3	19.3	28.1	31.8	38.1	38.7	41.3	40.2	40.2
	(19.3-27.5)	(15.3-24.0)	(17.1-27.1)	(15.1-24.4)	(19.2-28.8)	(24.5-32.0)	(27.3-36.6)	(33.1-43.4)	(34.4-43.1)	(36.1-46.6)	(34.9-45.7)	(34.9-45.7)
≥26	10.6	12.1	12.6	12.8	13.4	14.9	16.3	18.6	19.9	21.8	21.4	21.4
	(7.8-14.1)	(8.9-16.1)	(9.3-16.7)	(9.9-16.4)	(10.1-17.6)	(11.6-19.0)	(13.6-19.4)	(15.3-22.3)	(16.1-24.5)	(18.8-25.1)	(18.6-24.3)	(18.6-24.3)
Perceived great risk from smoking marijuana once or twice a week												
Total	43.0	42.5	41.6	39.7	40.6	39.3	36.0	32.1	29.5	27.7	26.1	26.1
	(39.4-46.6)	(38.6-46.6)	(37.9-46.4)	(36.0-43.6)	(35.2-46.2)	(35.3-43.4)	(32.3-39.9)	(28.7-35.8)	(26.1-33.1)	(23.7-32.2)	(22.8-29.6)	(22.8-29.6)
12-17	51.1	53.9	48.3	49.0	49.0	48.7	51.3	45.6	39.4	37.3	32.2	32.2
	(46.7-55.5)	(49.5-58.3)	(44.1-52.4)	(44.7-53.4)	(45.1-54.6)	(43.9-53.5)	(47.2-55.3)	(40.0-51.4)	(34.5-44.6)	(32.7-42.1)	(27.4-37.3)	(27.4-37.3)
18-25	26.1	31.2	27.9	31.3	30.6	25.3	19.7	16.7	18.2	15.5	14.2	14.2
	(21.8-30.9)	(24.7-38.4)	(22.8-33.7)	(25.5-37.8)	(26.3-35.3)	(21.8-29.2)	(15.9-24.2)	(13.1-21.0)	(14.5-22.6)	(12.4-19.2)	(10.9-18.4)	(10.9-18.4)
≥26	44.7	43.0	43.1	39.9	41.0	40.4	36.8	33.1	30.2	28.6	27.3	27.3
	(40.3-49.2)	(38.0-48.1)	(38.7-47.5)	(35.3-44.7)	(34.6-47.8)	(35.4-45.7)	(32.2-41.7)	(29.0-37.5)	(26.0-34.7)	(23.5-34.3)	(23.3-31.8)	(23.3-31.8)
Perceived no risk from smoking marijuana once or twice a week												
Total	6.7	6.8	6.4	6.8	8.9	9.3	10.7	14.6	15.7	16.9	17.8	17.8
	(5.1-8.8)	(4.8-8.5)	(4.9-8.4)	(5.3-8.7)	(6.7-11.8)	(7.1-12.1)	(8.8-13.1)	(11.9-17.9)	(12.7-19.3)	(14.3-20.0)	(15.7-20.2)	(15.7-20.2)
12-17	4.8	5.6	3.1	4.7	7.0	9.9	7.7	9.3	11.1	11.7	14.9	14.9
	(3.2-6.9)	(3.7-8.5)	(1.7-5.3)	(2.9-7.6)	(4.7-10.3)	(7.4-13.1)	(5.7-10.4)	(6.9-12.4)	(8.3-14.7)	(9.3-14.7)	(11.8-18.6)	(11.8-18.6)
18-25	13.7	11.4	10.2	10.7	14.0	16.3	20.6	27.1	29.4	30.3	29.3	29.3
	(10.7-17.4)	(8.4-15.2)	(7.5-13.7)	(8.0-14.1)	(10.8-18.0)	(13.3-19.7)	(16.7-25.1)	(23.6-31.0)	(25.2-34.0)	(25.2-35.9)	(24.9-34.1)	(24.9-34.1)
≥26	5.8	6.1	6.2	6.5	8.3	8.1	9.5	13.2	14.0	15.3	16.3	16.3
	(4.0-8.4)	(3.8-9.6)	(4.4-8.8)	(4.6-9.0)	(5.7-11.8)	(5.5-11.7)	(7.2-12.4)	(9.9-17.3)	(10.4-18.5)	(12.3-18.9)	(13.8-19.1)	(13.8-19.1)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
Abbreviation: CI = confidence interval.
Perceived great risk and no risk from smoking marijuana is defined as those who reported that smoking marijuana once a month and once or twice a week might cause harm [great risk] and might not cause harm [no risk].

Table 7. Percentage of perceived parental disapproval of trying marijuana once or twice and using marijuana once a month or more among persons aged 12 to 17 — National Survey on Drug Use and Health, Washington, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived parental disapproval of trying marijuana once or twice ¹												
12–17 years	95.6 (93.1-97.2)	95.7 (93.9-97.0)	96.9 (95.4-97.9)	96.5 (94.6-97.7)	95.7 (93.1-97.4)	95.5 (92.7-97.2)	95.2 (92.9-96.8)	95.1 (93.2-96.4)	94.9 (92.4-96.6)	94.6 (92.2-96.4)	94.5 (92.3-96.1)	93.1 (89.4-95.6)
Perceived parental disapproval of using marijuana once a month or more ²												
12–17 years	96.9 (95.1-98.1)	97.3 (95.7-98.4)	98.4 (97.2-99.1)	97.7 (96.2-98.6)	98.1 (96.2-99.1)	97.1 (95.0-98.3)	95.3 (92.9-96.9)	95.7 (93.7-97.0)	95.8 (93.7-97.3)	95.3 (92.8-97.0)	94.8 (92.1-96.6)	94.9 (92.0-96.7)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Perceived parental disapproval of trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of trying marijuana once or twice.

² Perceived parental disapproval of using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of using marijuana once a month or more.

Table 8. Percentage of perceived parental disapproval of trying marijuana once or twice and using marijuana once a month or more among past month marijuana users aged 12 to 17 — National Survey on Drug Use and Health, Washington, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived parental disapproval of trying marijuana once or twice ¹												
12–17 years	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Perceived parental disapproval of using marijuana once a month or more ²												
12–17 years	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.

Abbreviation: CI = confidence interval.

¹ Perceived parental disapproval of trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of trying marijuana once or twice.

² Perceived parental disapproval of using marijuana once a month or more is defined as those aged 12–17 years [youth] who reported perceived parental disapproval as strongly disapprove or somewhat disapprove of using marijuana once a month or more; indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.

Table 9. Percentage of perceived attitudes towards peers trying marijuana once or twice and using marijuana once a month or more among persons aged 12 to 17 — National Survey on Drug Use and Health, Washington, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
Perceived attitudes towards peers trying marijuana once or twice ¹	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
12–17 years	77.8 (73.3-81.7)	80.0 (75.9-83.9)	81.9 (77.6-85.6)	78.5 (74.7-81.8)	78.5 (74.9-81.9)	79.4 (75.0-83.2)	80.1 (76.3-83.4)	82.0 (78.4-85.1)	78.3 (74.0-82.1)	76.4 (72.0-80.3)	76.9 (73.2-80.1)	76.2 (72.1-79.9)
Perceived attitudes towards peers using marijuana once a month or more ²												
12–17 years	81.0 (77.1-84.4)	83.1 (79.2-86.4)	84.7 (80.6-88.1)	80.4 (76.8-83.6)	80.0 (76.4-83.2)	79.3 (75.9-83.9)	79.5 (75.3-83.2)	82.2 (78.9-85.1)	77.5 (73.2-81.3)	75.0 (70.5-79.1)	77.4 (73.4-80.9)	76.2 (72.0-80.0)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
Abbreviation: CI = confidence interval.
¹Disapproving attitudes toward peers trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers trying marijuana once or twice.
²Disapproving attitudes toward peers using marijuana once or a month or more is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers using marijuana once a month or more.

Table 10. Percentage of perceived attitudes towards peers trying marijuana once or twice and using marijuana once a month or more among past month marijuana users aged 12 to 17 — National Survey on Drug Use and Health, Washington, 2002–2014

	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
Perceived attitudes towards peers trying marijuana once or twice ¹	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
12–17 years	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)
Perceived attitudes towards peers using marijuana once a month or more ²												
12–17 years	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)	• (*•)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
Abbreviation: CI = confidence interval.
¹Disapproving attitudes toward peers trying marijuana once or twice is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers trying marijuana once or twice, indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.
²Disapproving attitudes toward peers using marijuana once or a month or more is defined as those aged 12–17 years [youth] who reported that they strongly disapprove or somewhat disapprove of peers using marijuana once a month or more, indicator was restricted to respondents aged 12–17 years who reported past month marijuana use.

Table 11. Percentage of perceived availability[†] among all persons aged ≥12 and mode of acquisition[†] of marijuana among past year marijuana users aged ≥12 by age group — National Survey on Drug Use and Health, Washington, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Perceived availability												
Total	62.1 (57.0-67.0)	61.6 (57.0-66.0)	62.1 (57.5-66.5)	61.7 (57.2-66.1)	59.7 (55.8-63.4)	58.5 (54.5-62.3)	58.0 (54.6-61.3)	58.8 (54.5-63.0)	62.3 (58.0-66.4)	63.6 (59.4-67.6)	67.2 (63.7-70.5)	74.6 (71.5-77.6)
12-17	59.6 (53.8-65.2)	53.8 (47.9-59.5)	52.8 (47.4-58.0)	52.7 (47.9-57.5)	49.4 (44.0-54.8)	49.0 (43.9-54.1)	51.9 (46.4-57.3)	50.3 (46.2-54.5)	50.0 (45.3-54.7)	50.1 (45.0-55.3)	52.3 (47.0-57.5)	57.1 (52.6-61.5)
18-25	63.5 (78.4-87.5)	78.3 (72.3-83.2)	76.7 (71.1-81.5)	82.1 (77.5-85.8)	81.6 (78.1-84.7)	80.8 (77.1-84.1)	82.5 (78.3-86.0)	81.1 (75.8-85.4)	81.8 (77.9-85.1)	80.1 (75.6-83.9)	79.7 (76.0-83.0)	81.5 (76.4-85.8)
≥26	58.7 (52.5-64.7)	59.6 (54.0-65.0)	60.7 (55.0-66.2)	59.4 (53.8-64.7)	57.4 (52.7-61.9)	56.0 (51.2-60.7)	54.7 (50.5-58.8)	56.1 (50.7-61.3)	60.4 (55.3-65.4)	62.4 (57.0-67.5)	66.8 (62.3-71.1)	75.5 (71.6-79.1)
Mode of acquisition of marijuana												
Bought it												
Total	34.6 (27.5-42.3)	34.7 (28.7-43.7)	40.7 (32.9-49.0)	45.1 (36.8-53.7)	46.8 (37.8-55.9)	44.0 (34.8-53.7)	42.0 (34.2-50.1)	49.1 (40.9-57.4)	46.4 (37.2-55.7)	41.5 (34.1-49.4)	43.6 (36.7-50.8)	44.5 (38.3-50.8)
12-17	23.7 (15.6-34.1)	18.3 (11.7-27.4)	(*)	(*)	(*)	(*)	(*)	(*)	35.7 (28.0-46.4)	(*)	(*)	(*)
18-25	41.5 (35.1-48.3)	41.1 (31.9-50.9)	36.7 (27.7-46.7)	38.5 (29.2-48.7)	40.4 (32.0-49.5)	44.2 (37.0-51.6)	45.2 (38.4-52.2)	43.4 (36.5-50.5)	46.7 (39.2-54.5)	49.5 (42.7-56.3)	49.2 (42.5-55.9)	48.3 (40.1-56.7)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	38.1 (27.2-50.4)	42.5 (33.1-52.4)	44.8 (36.8-53.2)
Traded something for it												
Total	0.3 (0.1-1.1)	0.8 (0.2-2.7)	0.7 (0.2-2.5)	0.3 (0.1-1.1)	(*)	2.6 (0.7-9.1)	(*)	2.0 (0.5-7.2)	0.9 (0.4-1.8)	0.6 (0.3-1.6)	0.9 (0.3-2.7)	0.9 (0.3-2.6)
12-17	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
18-25	0.4 (0.1-1.8)	(*)	1.6 (0.4-6.5)	(*)	1.2 (0.3-5.1)	1.3 (0.3-5.3)	(*)	1.3 (0.5-3.5)	2.2 (1.0-4.9)	2.1 (0.9-5.1)	1.2 (0.4-3.9)	(*)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	0.8 (0.2-4.2)	1.0 (0.3-3.7)
Got it for free or shared with someone else												
Total	64.4 (56.4-71.7)	64.2 (55.0-72.4)	59.6 (50.3-68.4)	53.3 (44.6-61.7)	49.4 (40.7-58.1)	52.7 (42.7-62.6)	55.1 (46.6-63.3)	48.1 (39.0-57.3)	51.6 (42.4-60.6)	56.3 (48.6-63.7)	52.1 (44.8-59.4)	50.9 (44.7-57.1)
12-17	74.5 (63.7-82.9)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	61.5 (51.3-70.8)	(*)	(*)	(*)
18-25	57.1 (49.8-64.1)	56.4 (47.3-65.0)	61.7 (52.5-70.1)	60.1 (50.5-69.0)	57.3 (48.1-66.0)	54.0 (46.4-61.3)	52.8 (45.2-60.3)	53.9 (46.5-61.2)	49.9 (42.3-57.6)	46.9 (40.1-53.8)	48.1 (41.4-54.9)	49.5 (41.1-57.9)
≥26	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	60.1 (48.0-71.0)	52.3 (42.3-62.1)	49.9 (41.6-58.2)
Grew it yourself												

Total	0.7 (0.2-2.2)	0.4 (0.1-1.7)	1.3 (0.4-4.3)	1.7 (0.6-4.6)	0.6 (0.2-2.2)	0.6 (0.2-1.6)	0.9 (0.3-2.5)	1.2 (0.4-3.2)	1.5 (0.5-4.3)	3.3 (1.3-8.2)	3.7 (1.7-8.0)
12-17	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)
18-25	0.9 (0.2-4.3)	1.1 (0.2-5.4)	1.0 (0.2-3.9)	1.0 (0.3-3.8)	1.5 (0.5-4.4)	1.4 (0.5-3.7)	1.4 (0.5-3.7)	1.1 (0.3-3.8)	1.4 (0.4-4.6)	1.5 (0.5-4.4)	1.6 (0.5-4.5)
≥26	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	1.8 (0.4-7.0)	4.4 (1.5-12.5)	4.2 (1.5-11.2)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.
Abbreviation: CI = confidence interval.
* Perceived availability is defined as those who reported that it would be fairly easy or very easy for them to obtain marijuana if they wanted some.
† Mode of acquisition is defined as those who reported how they got the marijuana they used for the last time [i.e., bought it, traded something for it, got it for free or shared with someone else, or grew it yourself], this question is only asked of those respondents indicating use of marijuana in the past 12 months.

Table 12. Percentage of past year marijuana dependence and abuse among all persons aged ≥12 by age group — National Survey on Drug Use and Health, Washington, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	2.1 (1.5-2.9)	2.1 (1.5-2.9)	2.6 (1.9-3.6)	2.4 (1.8-3.4)	1.7 (1.3-2.2)	1.3 (0.9-2.0)	1.4 (1.0-1.9)	1.9 (1.3-2.8)	2.2 (1.5-3.1)	2.2 (1.4-3.6)	2.4 (1.7-3.3)	2.4 (1.7-3.4)
12-17	5.1 (3.2-7.9)	3.5 (2.2-5.5)	2.2 (1.2-3.7)	4.1 (2.7-6.0)	4.5 (3.2-6.2)	4.1 (2.5-6.7)	3.9 (2.5-5.8)	3.3 (2.0-5.5)	3.7 (2.2-6.0)	4.0 (2.4-6.7)	3.6 (2.3-5.5)	2.9 (1.8-4.6)
18-25	8.8 (6.5-11.9)	8.2 (5.9-11.2)	7.8 (5.7-10.5)	7.1 (5.1-9.6)	6.5 (4.6-9.1)	4.3 (2.8-6.4)	5.6 (3.5-8.9)	7.8 (5.6-10.7)	7.5 (5.4-10.2)	5.9 (4.2-8.3)	6.1 (4.1-8.9)	5.4 (3.5-8.2)
≥26	0.6 (0.2-1.4)	0.9 (0.4-1.7)	1.8 (1.0-3.2)	1.4 (0.7-2.8)	0.5 (0.2-1.3)	0.5 (0.2-1.4)	0.4 (0.2-1.2)	0.8 (0.4-1.8)	1.1 (0.5-2.4)	1.4 (0.5-3.5)	1.6 (0.9-3.1)	1.9 (1.1-3.1)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Substance use disorder (also known as dependence or abuse) is defined as meeting criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association [APA], 1994) for either dependence or abuse for illicit drugs or alcohol.

Table 13. Percentage of past year marijuana dependence and abuse among past year marijuana users aged ≥12, by age group — National Survey on Drug Use and Health, Washington, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Total	16.5 (11.7-20.3)	17.7 (13.4-23.1)	22.4 (16.6-29.5)	19.1 (13.7-25.8)	13.6 (10.1-17.9)	10.1 (6.9-14.6)	10.7 (7.7-14.5)	13.4 (9.3-19.0)	14.0 (10.0-19.4)	13.7 (8.4-21.5)	12.9 (9.0-18.2)	12.4 (9.0-16.8)
12-17	(*) (*)	23.2 (15.6-33.0)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	21.8 (13.5-33.2)	(*) (*)	(*) (*)	(*) (*)
18-25	24.1 (18.6-30.5)	28.4 (21.3-36.9)	26.4 (19.5-34.7)	21.6 (15.6-29.1)	20.8 (15.3-27.7)	14.3 (10.1-19.8)	16.7 (11.0-24.6)	19.5 (14.6-25.6)	18.5 (14.1-24.0)	16.0 (11.7-21.4)	16.1 (11.3-22.3)	15.7 (10.6-22.7)
≥26	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	(*) (*)	10.5 (5.5-19.1)	10.8 (6.5-17.4)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002–2014.
 Abbreviation: CI = confidence interval.
 * Substance use disorder (also known as dependence or abuse) is defined as meeting criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association [APA], 1994) for either dependence or abuse for illicit drugs or alcohol.
 † Analysis limited to past year marijuana users (defined as those who reported use of marijuana within 12 months preceding the date of interview, including those who reported use of marijuana within 30 days preceding the date of interview)

Table 14. Percentage of perception of maximum legal penalty in your state for first offense possession of an ounce or less of marijuana for your own use* among all persons aged ≥12, by age group — National Survey on Drug Use and Health, Washington, 2002–2014

Age group (yrs)	2002-2003	2003-2004	2004-2005	2005-2006	2006-2007	2007-2008	2008-2009	2009-2010	2010-2011	2011-2012	2012-2013	2013-2014
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Fine												
Total	29.5 (25.8-33.4)	31.1 (25.9-36.8)	29.1 (24.5-34.2)	29.7 (25.6-34.3)	31.9 (28.1-35.9)	32.8 (28.6-37.3)	35.0 (30.6-39.5)	37.4 (32.7-42.4)	38.7 (34.4-43.1)	38.0 (33.2-43.0)	29.6 (25.4-34.1)	23.0 (19.8-26.8)
12-17	8.4 (6.0-11.8)	9.7 (6.7-13.8)	14.2 (10.1-19.7)	14.9 (11.4-19.1)	14.3 (10.7-19.0)	16.4 (12.5-21.2)	17.6 (14.2-21.5)	18.5 (15.0-22.7)	22.5 (18.5-27.1)	21.3 (17.1-26.1)	20.3 (16.2-25.1)	26.0 (21.2-31.5)
18-25	23.5 (20.3-27.0)	22.8 (19.2-27.0)	24.5 (19.8-30.0)	25.9 (20.7-32.0)	25.0 (20.6-30.0)	24.4 (20.0-29.4)	28.5 (23.6-33.9)	34.1 (30.1-38.4)	34.6 (30.7-38.6)	32.3 (27.9-37.0)	30.5 (25.9-35.6)	31.1 (25.5-37.3)
≥26	34.2 (29.2-39.5)	36.2 (29.6-43.4)	32.6 (26.5-39.4)	32.9 (27.4-39.0)	36.0 (31.0-41.3)	36.9 (31.6-42.6)	38.6 (33.1-44.3)	40.7 (34.7-47.0)	41.9 (35.9-48.1)	41.6 (35.3-48.2)	30.5 (25.1-36.5)	21.2 (17.3-25.8)
Probation												
Total	16.0 (13.1-19.4)	16.1 (13.2-19.5)	17.4 (14.4-20.8)	16.8 (13.9-20.3)	14.7 (11.7-18.4)	14.7 (11.8-18.0)	14.5 (11.8-17.6)	14.7 (12.0-17.8)	14.0 (11.4-17.1)	14.3 (11.6-17.5)	9.7 (7.8-12.4)	5.8 (4.5-7.3)
12-17	18.4 (14.7-22.8)	18.5 (14.7-23.1)	19.2 (14.5-25.1)	19.8 (15.0-25.7)	19.6 (15.3-24.7)	20.6 (15.9-26.3)	20.3 (16.4-24.8)	18.7 (15.1-22.9)	17.8 (15.1-20.9)	18.8 (15.3-23.0)	18.7 (14.6-23.7)	19.5 (15.0-25.1)
18-25	13.1 (10.0-17.0)	17.3 (13.5-21.9)	18.1 (14.4-22.7)	15.8 (12.3-20.0)	15.5 (12.2-19.6)	16.5 (12.9-20.9)	18.6 (15.3-22.3)	19.4 (15.8-23.7)	18.9 (14.1-24.8)	20.0 (16.3-24.2)	16.5 (12.7-21.2)	10.8 (7.8-14.9)
≥26 years	16.3 (12.4-21.1)	15.5 (11.7-20.2)	16.9 (13.2-21.4)	16.6 (12.7-21.4)	13.8 (9.9-18.9)	13.4 (9.9-17.9)	12.9 (9.6-17.0)	13.1 (9.8-17.5)	12.4 (9.4-16.3)	12.5 (9.2-16.8)	7.3 (5.0-10.5)	3.3 (2.1-5.3)
Community Service												

Total	10.5 (8.1-13.4)	9.3 (7.0-12.3)	8.8 (6.5-11.7)	7.5 (5.7-9.9)	7.8 (5.9-10.3)	8.3 (6.3-10.9)	8.4 (6.3-11.1)	9.2 (6.5-12.9)	9.1 (6.5-12.7)	10.7 (8.0-14.3)	6.8 (4.6-9.8)	2.4 (1.7-3.4)
12-17	20.1 (15.8-25.1)	20.9 (17.6-24.7)	20.6 (16.8-25.1)	17.3 (13.3-22.2)	13.7 (10.5-17.5)	12.4 (9.5-15.9)	13.0 (9.6-17.4)	10.5 (7.9-13.8)	11.3 (8.7-14.5)	13.5 (10.4-17.3)	12.0 (8.7-16.3)	10.0 (6.6-14.9)
18-25	13.4 (10.3-17.4)	11.7 (8.5-15.9)	10.8 (7.7-14.9)	9.7 (6.8-13.7)	9.0 (6.7-12.1)	9.5 (7.0-12.8)	9.1 (6.3-12.9)	8.6 (6.2-11.8)	9.2 (7.1-11.9)	9.4 (7.1-12.2)	6.7 (4.5-9.7)	4.1 (2.4-7.0)
≥26	8.3 (5.6-12.2)	7.0 (4.3-11.3)	6.4 (3.8-10.6)	5.5 (3.4-8.8)	6.7 (4.4-10.1)	7.5 (5.0-11.1)	7.6 (5.0-11.2)	9.2 (5.8-14.3)	8.8 (5.5-13.7)	10.6 (7.2-15.4)	6.2 (3.7-10.1)	1.2 (0.6-2.4)
Possible Prison Sentence												
Total	37.9 (33.6-42.5)	36.8 (32.7-41.1)	39.3 (33.4-43.3)	36.6 (32.5-40.8)	34.2 (29.8-38.9)	34.8 (30.4-39.6)	31.8 (27.5-36.3)	27.9 (24.6-31.5)	26.3 (22.2-30.9)	24.8 (21.3-28.7)	16.0 (13.3-19.2)	8.2 (6.4-10.4)
12-17	38.8 (33.2-44.6)	36.1 (31.8-40.6)	36.4 (31.1-42.0)	35.7 (30.1-41.8)	36.5 (30.9-42.5)	35.1 (29.3-41.4)	31.9 (26.2-38.2)	34.2 (29.6-39.0)	32.5 (28.1-37.2)	32.2 (26.9-37.9)	30.1 (25.0-35.7)	18.7 (14.8-23.3)
18-25	42.0 (37.8-46.2)	39.6 (34.4-45.1)	40.0 (34.8-45.5)	41.2 (36.1-46.6)	39.5 (34.0-45.2)	39.2 (34.3-44.3)	34.4 (28.8-40.5)	30.1 (25.3-35.4)	30.1 (24.9-35.8)	28.8 (25.2-32.6)	20.1 (16.8-23.9)	13.1 (10.0-16.9)
≥26	37.0 (31.1-43.2)	36.3 (30.8-42.2)	38.1 (31.8-44.9)	35.7 (30.2-41.6)	32.8 (27.0-39.1)	33.9 (28.2-40.2)	31.3 (26.0-37.1)	26.6 (22.2-31.5)	24.7 (19.3-31.0)	22.9 (18.4-28.2)	13.5 (10.3-17.6)	6.1 (4.2-8.9)
Mandatory Prison Sentence												
Total	6.0 (4.4-8.2)	6.7 (4.9-9.1)	6.5 (4.5-9.3)	6.7 (4.7-9.3)	6.6 (4.8-8.8)	4.8 (3.4-6.7)	5.2 (3.6-7.3)	5.3 (3.8-7.3)	5.4 (3.8-7.7)	5.0 (3.0-8.0)	2.9 (1.9-4.4)	2.5 (1.6-3.7)
12-17	14.4 (11.1-18.3)	14.7 (11.2-19.0)	9.5 (6.7-13.3)	11.2 (8.1-15.4)	13.3 (10.2-17.3)	11.4 (8.0-16.0)	11.6 (8.6-15.5)	13.5 (10.3-17.6)	11.8 (8.6-16.0)	9.0 (6.9-11.7)	7.7 (5.4-10.7)	8.3 (5.7-11.9)
18-25	8.0 (5.8-10.9)	8.6 (5.7-12.7)	6.5 (3.8-11.0)	6.4 (4.1-9.6)	8.0 (5.5-11.3)	7.5 (5.5-10.1)	6.4 (4.6-8.8)	4.3 (2.6-7.2)	3.6 (2.4-5.4)	4.8 (3.2-7.1)	4.6 (3.2-6.7)	3.7 (2.0-6.7)
≥26	4.3 (2.3-7.8)	5.0 (2.9-8.5)	6.0 (3.5-10.0)	6.0 (3.6-9.8)	5.3 (3.3-8.3)	3.3 (1.7-6.1)	4.1 (2.3-7.0)	4.3 (2.5-7.4)	4.8 (2.3-9.7)	4.4 (2.1-9.0)	1.9 (0.9-4.2)	1.6 (0.8-3.3)

No penalty												
Total	DNA	DNA	DNA	2.7 (1.4-5.1)	4.7 (3.1-7.0)	4.6 (2.9-7.1)	5.3 (3.8-7.3)	5.5 (3.7-9.2)	6.5 (4.3-9.7)	7.2 (5.2-9.9)	35.1 (30.4-40.0)	58.2 (53.9-62.4)
12-17	DNA	DNA	DNA	1.0 (0.5-2.0)	2.5 (1.4-4.5)	4.1 (2.3-7.4)	5.6 (3.4-9.1)	4.6 (2.6-8.0)	4.2 (2.4-7.3)	5.2 (2.3-11.2)	11.2 (7.6-16.4)	17.5 (13.1-22.9)
18-25	DNA	DNA	DNA	1.0 (0.4-2.5)	3.0 (1.5-5.9)	2.9 (1.5-5.9)	3.1 (1.9-4.8)	3.4 (2.0-5.8)	3.7 (2.1-6.3)	4.8 (3.0-7.7)	21.5 (17.5-26.1)	37.2 (31.3-43.5)
≥26	DNA	DNA	DNA	3.3 (1.6-6.8)	5.4 (3.3-8.7)	5.0 (2.9-8.3)	5.6 (3.8-8.3)	6.1 (3.7-9.8)	7.4 (4.5-11.9)	7.9 (5.3-11.7)	40.5 (34.8-46.6)	66.5 (61.4-71.3)

Source: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Abbreviations: CI = confidence interval; DNA = data not available.

*Perception of legal penalty for marijuana possession is defined as those who reported their perception of the maximum legal penalty in their state of residence for first offense possession of an ounce or less of marijuana for their own use.



January 2016

The Legalization of Marijuana in Colorado: *The Impact*

Latest Results for Colorado

Youth and Adult Marijuana Use

ROCKY MOUNTAIN HIGH INTENSITY
DRUG TRAFFICKING AREA
www.rmhidta.org



Introduction

- This report on marijuana use in Colorado is an update of the publication *The Legalization of Marijuana in Colorado: The Impact Volume 3*.
- This report is a summary of some of the data from the National Survey on Drug Use and Health (NSDUH), an annual survey sponsored by the Substance Abuse and Mental Health Services Administration (SAMHSA).
- The NSDUH data reported by SAMHSA is an average of two consecutive years.
- Colorado legalized recreational marijuana in 2013 and retail marijuana businesses began operation in 2014.
- The findings are between the two-year average of full legalization (2013/2014) compared to the two-year average just prior to legalization (2011/2012).


Youth Findings

Past Month (Current) Marijuana Use for Colorado Youth Ages 12 to 17 Years Old:

- In the two year average (2013/2014) since Colorado legalized recreational marijuana, youth past month marijuana use increased 20 percent compared to the two year average prior to legalization (2011/2012).
 - Nationally youth past month marijuana use declined 4 percent during the same time.
- The latest 2013/2014 results show Colorado youth ranked #1 in the nation for past month marijuana use, up from #4 in 2011/2012.
- Colorado youth past month marijuana use for 2013/2014 was 74 percent higher than the national average compared to 39 percent higher in 2011/2012.

Youth Data


Youth Ages 12 to 17 Years Old Past Month Marijuana Use National vs. Colorado



Annual Averages of Data Collection

SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014

Past Month Usage by 12 to 17 Years Old, 2013/2014




SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014

NOTE:

*Oregon and Alaska voted to legalize recreational marijuana in November 2014

**States that had legislation for medical marijuana signed into effect during 2014

Average Past Month Use by 12 to 17 Years Old, 2013/2014



SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014

States for Past Month Marijuana Use Youth Ages 12 to 17 Years Old, 2013/2014

<u>Top 10</u> (Medical/Recreational States)	<u>Bottom 10</u> (Non-Medical or Recreational States)
National Rate = 7.22%	
1. Colorado – 12.56%	41. Mississippi – 5.60%
2. Vermont – 11.40%	42. West Virginia – 5.60%
3. Rhode Island – 10.69%	43. North Dakota – 5.60%
4. Oregon – 10.19%	44. Louisiana – 5.55%
5. Washington – 10.06%	45. Nebraska – 5.54%
6. Maine – 9.90%	46. Oklahoma – 5.52%
7. New Hampshire – 9.83%	47. Utah – 5.42%
8. Alaska – 9.19%	48. South Dakota – 5.32%
9. Massachusetts – 8.88%	49. Iowa – 5.17%
10. California – 8.74%	50. Alabama – 4.98%

SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014


College Age Findings

Past Month (Current) Marijuana Use for Colorado College Age Adults Ages 18 to 25 Years Old:

- In the two year average (2013/2014) since Colorado legalized recreational marijuana, college age past month marijuana use increased 17 percent compared to the two year average prior to legalization (2011/2012).
 - Nationally college age past month marijuana use increased 2 percent during the same time.
- The latest 2013/2014 results show Colorado college age adults ranked #1 in the nation for past month marijuana use, up from #3 in 2011/2012.
- Colorado college age past month marijuana use for 2013/2014 was 62 percent higher than the national average compared to 42 percent higher in 2011/2012.

College Age Data


College Age (18 to 25 Years Old) Past Month Marijuana Use



Annual Averages of Data Collection

SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014

Past Month Usage by 18 to 25 Years Old, 2013/2014




SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014

NOTE:

*Oregon and Alaska voted to legalize recreational marijuana in November 2014

**States that had legislation for medical marijuana signed into effect during 2014

Average Past Month Use by 18 to 25 Years Old, 2013/2014



SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014

States for Past Month Marijuana Use College Age Adults 18 to 25 Years Old, 2013/2014

<u>Top 10</u> (Medical/Recreational States)	<u>Bottom 10</u> (Non-Medical or Recreational States)
National Rate = 19.32%	
1. Colorado – 31.24%	41. Oklahoma – 15.76%
2. Vermont – 30.60%	42. Kansas – 15.11%
3. New Hampshire – 30.09%	43. Texas – 15.06%
4. Rhode Island – 28.90%	44. Alabama – 15.04%
5. Massachusetts – 28.74%	45. Tennessee – 14.72%
6. Maine – 28.38%	46. Idaho – 14.28%
7. Oregon – 24.85%	47. North Dakota – 14.05%
8. Washington – 24.47%	48. Iowa – 14.01%
9. Maryland – 23.42%	49. South Dakota – 13.02%
10. Michigan – 23.17%	50. Utah – 11.55%

SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014


Adult Findings

Past Month (Current) Marijuana Use for Colorado Adults Ages 26+ Years Old:

- In the two year average (2013/2014) since Colorado legalized recreational marijuana, adult past month marijuana use increased 63 percent compared to the two year average prior to legalization (2011/2012).
 - Nationally adult past month marijuana use increased 21 percent during the same time.
- The latest 2013/2014 results show Colorado adults ranked #1 in the nation for past month marijuana use, up from #7 in 2011/2012.
- Colorado adult past month marijuana use for 2013/2014 was 104 percent higher than the national average compared to 51 percent higher in 2011/2012.

Adult Data


Adult (26+ Years Old) Past Month Marijuana Use



Annual Averages of Data Collection

SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014

Past Month Usage by 26+ Years Old, 2013/2014




SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014

NOTE:

*Oregon and Alaska voted to legalize recreational marijuana in November 2014

**States that had legislation for medical marijuana signed into effect during 2014

Average Past Month Use by 26+ Years Old, 2013/2014



SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014

States for Past Month Marijuana Use Adults Ages 26+ Years Old, 2013/2014

Top 10 (Medical/Recreational States)	Bottom 10 (Non-Medical or Recreational States)
National Rate = 6.11%	
1. Colorado – 12.45%	41. Louisiana – 4.42%
2. Washington – 11.21%	42. Utah – 4.25%
3. Maine – 10.77%	43. Texas – 4.21%
4. Oregon – 10.68%	44. Alabama – 4.03%
5. Alaska – 10.42%	45. Tennessee – 4.01%
6. Vermont – 10.42%	46. Nebraska – 3.97%
7. Rhode Island – 9.92%	47. North Dakota – 3.95%
8. Massachusetts – 9.08%	48. Mississippi – 3.95%
9. New Hampshire – 8.78%	49. Iowa – 3.40%
10. Montana – 8.49%	50. South Dakota – 3.30%

SOURCE: SAMHSA.gov, National Survey on Drug Use and Health 2013 and 2014

