

Ministry of Environment of Denmark Department

Derogation Report 2020

Danish Report in accordance with the Commission Decisions 2005/294/EC, 2008/664/EC, 2012/659/EU, 2017/847/EU, 2018/1928/EU and 2020/1074/EU

July 2021

Ministry of Environment of Denmark Department Slotsholmsgade 12 DK-1216 Copenhagen K

Table of contents

1.	Introduction	4
2.	Maps of cattle holdings, arable land and livestock in kg N in 2018/2019	5
2.1	Map of derogation holdings 2018/2019	5
2.2	Map of arable land 2018/2019	5
2.3	Map of livestock in kg N in 2018/2019	5
2.4	Use of the derogation	6
2.5	Trends in livestock	10
3.	Controls at farm level	12
3.1	Control of compliance with the Danish derogation	12
3.2	Summary of inspection results 2020	12
3.3	Inspection of compliance within the derogation year	12
3.4	Results	13
3.5	General inspection of the harmony rules	14
3.6	Control of fertilizer accounts	17
4.	Water quality	19
4.1	Introduction	20
4.2	Development in agricultural practices at the national level from 2005 to 2019	23
4.3	Modelled nitrate leaching for farm types and geographical areas and the	
	impact of derogation farms at the national level – 2019 data	26
4.4	Development in modelled nitrate leaching in the Agricultural Catchment	
	Monitoring Programme 1990-2018	34
4.5	Measurements of nitrate in water leaving the root zone	36
4.6	The nitrogen flow to surface water in agricultural catchments	40
5.	Reinforced monitoring in areas characterized by sandy soils	45
5.1	Introduction	45
5.2	Method	45
5.3	Characterization of monitoring stations and data analysis	48
5.4	Results and Discussion	49
6.	Indicator and monitoring system for application of phosphorus in	
	Denmark	55
6.1	Introduction	55
6.2	Results from the P monitoring system	55
6.3	Results from P indicator system	56
7.	Targeted catch crops scheme and targeted nitrogen regulation	59
7.1	Results from 2017 to 2020	60
8.	Conclusions	61
8.1	Cattle holdings and controls on farm level	61
8.2	Water quality	61
8.3	Targeted catch crops and targeted nitrogen regulation	62
8.4	The reinforced monitoring	62
8.5	The phosphorus indicator and monitoring system	62

1. Introduction

With Commission Decisions 2005/294/EC, 2008/664/EC, 2012/659/EU, 2017/847/EU, 2018/1928/EU, and 2020/1074/EU Danish cattle holdings are allowed to derogate from the general rules in the Nitrates Directive (91/676/EEC).

The relevant decisions for the data reported in this report are 2018/1928/EU and 2020/1074/EU. According to these decisions, cattle holdings could apply for authorisations to apply livestock manure corresponding to up to 230 kg N per hectare per year if more than 80 per cent of the area available for manure application was cultivated with beets, grass or grass catch crops. Furthermore, derogation holdings have to comply with several other conditions laid down in the decision.

The aim of this report is to present maps showing the percentage of farms and percentage of agricultural land encompassed by the derogation in each Danish municipality for the planning period 2018/2019.

According to the decisions 2018/1928/EU and 2020/1074/EU, the Danish authorities shall submit the following information to the Commission for the derogation period 2018/2019:

- According to Article 10 (1) and 12 (a): maps, showing the percentage of cattle farms, percentage of livestock and percentage of agricultural land covered by the derogation for each municipality of Denmark.
- According to Article 12 (g), an evaluation of the implementation of the derogation conditions, on the basis of controls at farm level and information on non-compliant farms, based on the results of the administrative and field inspections.
- According to Article 12 (b, c, e), the results on ground and surface water monitoring as regards nitrate and phosphate, including information on water quality trends as well as the impact of derogation on water quality. Further results of model-based calculations from farms benefiting from an individual derogation.
- According to Article 12 (d and f), the results of the surveys on local land use, crop rotations and agricultural practices including tables showing the percentage of agricultural land under derogation covered by clover or alfalfa in grassland and by barley/pea, undersown with grass.
- According to article 12 (h), trends in livestock numbers and manure production for each livestock category in Denmark and in derogation farms.

The derogation decision 2018/1928/EU and 2020/1074/EU requires according to Articles 10 (2) and 12 (b), reporting of water quality data from reinforced monitoring on sandy soils and in an area, where at least 3% of all derogation farms are located. The monitoring data is updated with data from 2019 in this report.

Various Danish authorities and institutions have contributed to this report, edited by the Ministry of Environment of Denmark. The respective authors, and hence responsible institutions for the different chapters, can be found under the heading to the respective chapters.

2. Maps of cattle holdings, arable land and livestock in kg N in 2018/2019

Lars Paulsen & Lene Kragh Møller, the Danish Agricultural Agency, Ministry of Food, Agriculture and Fisheries of Denmark, November 2020

For the planning period 2018/2019, the Danish Agricultural Agency received 33,056 fertilizer accounts containing key figures on the use of nitrogen (commercial fertilizer and livestock manure). The accounts were registered and reviewed. The maps (Figure 2.1 - Figure 2.3) are based on the number of agricultural holdings, kg N per hectare per year and arable land used by derogation farms in 2018/2019. The fertilizer accounting year runs from 1st of August to 31st of July. Accounts for 2018/2019 were to be submitted to the Danish Agricultural Agency no later than 31st of March 2020.

In the fertilizer account, the farmer states whether the derogation was used. This means that the individual farmer needs to apply for the use of the derogation when the farmer submits the fertilizer quota and catch crops plan (at the latest 21st of April each year). The information about the application is automatically transferred to the fertilizer accounting system. The maps of cattle holdings, arable land and kg N from organic fertilisers per hectare per year are based on the data reported by the farmers. In reports before 2019, a map with livestock units per year was presented. This has from 2019 been replaced by a map showing kg N from organic fertilisers, including livestock manure per hectare and year at municipal level. In Danish regulation, it has generally from 2019 been changed to limit livestock density at farm level via a maximum allowable N application from organic fertilisers (instead of number of livestock). However, since one livestock unit corresponds to 100 kg N (ex storage), the data is directly convertible and hence does not present any change in the limitation per area.

2.1 Map of derogation holdings 2018/2019

The map (Figure 2.1) shows derogation holdings in percentage of the total number of agricultural holdings registered in each respective Danish municipality.

In 2018/2019, 1,284 derogation holdings were encompassed by the derogation. This corresponds to 3.9 % of all registered fertilizer accounts. The applied amount of manure on these farms ranged from 170 to 230 kg N per hectare per year. If the production of manure on a derogation farm corresponds to more than 230 kg N per hectare, the farmer is obliged to deliver the excess manure to one or more contractual partner-farmers.

2.2 Map of arable land 2018/2019

The map (Figure 2.2) shows the share of arable land on derogation holdings in relation to the total agricultural area in each Danish municipality.

In 2018/2019, the arable land on cattle holdings encompassed by the derogation was 195,804 hectare at national scale. This corresponded to 8.1 % of the registered area used for agriculture in Denmark.

2.3 Map of livestock in kg N in 2018/2019

The map (Figure 2.3) shows the share of kg N distributed from cattle holdings encompassed by the derogation holdings in relation to the total kg N from organic fertilisers in each Danish municipality.

In 2018/2019, the kg N from organic fertilisers distributed from cattle holdings encompassed by the derogation was 39.1 million kg N in total. This corresponded to

17.8 % of all kg N in organic fertilisers distributed on the agricultural area in Denmark.

2.4 Use of the derogation

Over the first three planning periods in which the derogation was used, i.e. 2002/2003, 2003/2004 and 2004/2005, an increase in the use of the derogation was registered both regarding the number of farms, the number of hectares and the number of livestock units (**Table 2.1**). This tendency was broken in 2005/2006, where a decrease was observed for all three measured parameters and the decreasing trend continued until the period 2008/2009. Between 2009/2010 and 2015/2016, an overall increase in the agricultural area using the derogation was observed, whereas the number of farms remained at a more constant level. The general trend of Danish farms becoming bigger is reflected in these numbers and from 2016/2017 there has been a decrease in the number of farms and the number of hectares encompassed by the derogation. From 2017/2018, the number of livestock unit was replaced by produced kg N per year in the Danish legislation.

Table 2.1: Development in use of the derogation for number of farms, agricultural area and kg N in organic fertilisers per year (livestock units (LU) until 2016/2017) from 2002/2003 until 2018/2019 (One LU=100 kg N (ex storage)).

Year	Number of derogation farms	Share of total farms (%)	Area of derogation (hectare)	Share of total Area (%)	Number of LUs	Share of total LUs (%)
2002/2003	1,845	4.0	123,068	5.0	213,617	10.6
2003/2004	1,927	4.0	128,523	5.0	225,586	10.6
2004/2005	2,331	5.0	134,780	5.0	277,330	12.9
2005/2006	1,779	3.4	115,336	4.2	220,839	10.3
2006/2007	1,610	3.2	111,845	4.0	211,765	9.5
2007/2008	1,296	2.8	92,282	3.9	186,313	8.3
2008/2009	1,115	2.4	90,647	3.6	176,588	8.2
2009/2010	1,507	3.3	134,698	6.1	276,765	11.9
2010/2011	1,607	3.9	164,353	7.4	341,781	14.1
2011/2012	1,652	4.0	175,783	7.1	365,887	15.5
2012/2013	1,481	3.7	162,176	6.7	334,508	14.5
2013/2014	1,482	3.8	189,495	7.7	397,014	17.1
2014/2015	1,500	4.0	205,165	8.2	425,102	18.6
2015/2016	1,466	4.2	210,061	8.6	443,134	19.4
2016/2017	1,378	3.9	205,874	8.4	439,114	19.3
					Mill. kg N (org. fert.)	Share of total kg N (%)
2017/2018	1,312	3.9	198,195	8.2	39.6	18.1
2018/2019	1,284	3.9	195,804	8.1	39.1	17.8

The livestock density on derogation farms has remained at an approximately constant level, compared to the periods 2009/2010-2016/2017 and the average number of livestock units per farm has increased over the same period. From 2017/2018, the average livestock size and the average livestock density were measured in kg N (from organic fertilisers) per holding and in kg N (from organic fertilisers) per hectare per year.

By comparison, a total number of 10,236 Danish agricultural holdings had cattle as livestock in 2018/2019. These holdings housed in total 116.4 million kg N from organic fertilisers and covered an agricultural area of 853,899 hectare. This gave an average of 11,372 kg N from organic fertilisers per cattle holding and an average livestock density of 136 kg N from organic fertilisers per hectare on all cattle Danish farms. Consequently, approximately 12.5 % of all cattle farms were derogation farms in 2018/2019, and the derogation (cattle) farms housed 33.6 % of all cattle-kg N in Denmark, covering 22.9 % of the total Danish cattle farm area.

Table 2.2: Average number of spread livestock units¹ (LU) per holding and per hectare under the derogation until 2016/2017. From 2017/2018 the number of livestock is expressed by kg N from organic fertilisers (One LU = 100 kg N (ex storage)).

Year	Average livestock size (LU/holding)	Average livestock den- sity (LU/ha)
2002/2003	115.78	1.74
2003/2004	117.07	1.76
2004/2005	118.97	2.06
2005/2006	124.14	1.91
2006/2007	131.53	1.89
2007/2008	143.76	2.02
2008/2009	158.37	1.95
2009/2010	183.65	2.05
2010/2011	212.68	2.08
2011/2012	221.48	2.08
2012/2013	225.86	2.06
2013/2014	267.89	2.10
2014/2015	283.40	2.07
2015/2016	302.27	2.11
2016/2017	318.66	2.13
	Average livestock size (kg N/holding) ²	Average livestock den- sity (kg N/ha)
2017/2018	30,171	199
2018/2019	30,475	200

¹ "Spread LU" is the term used to describe the amount of livestock manure, which is being applied to agricultural land within the farm, as this amount can be different from the amount of livestock manure produced at farm level due to import or export of livestock manure from/to other farms. One LU corresponds to 100 kg manure-N (ex storage) in the Danish system.

² From 2017/2018, the number of livestock units (LU) is replaced by produced kg N from organic fertilisers per year in the Danish legislation (One LU = 100 kg N).




Figure 2.1: Derogation holdings in percent of total number of agricultural holdings in Denmark in 2018/2019. The location of each holding is determined by address of the owner.




Figure 2.2: Agricultural land encompassed by the derogation in 2018/2019 in percent of the total agricultural area in Denmark. The location of each holding is determined by address of the owner.

The maps (**Figure 2.1 - Figure 2.3**) illustrate that derogation cattle holdings are concentrated in the western parts of Jutland. A few holdings are located on Zealand and even fewer on Funen and the island of Bornholm.




Figure 2.3: Kg N from organic fertilisers per hectare per year spread on derogation farms in percent of total kg N from organic fertilisers in 2018/2019 in Denmark. The location of each holding is determined by address of the owner.

2.5 Trends in livestock

According to decision 2018/1928/EU, and 2020/1074/EU the Danish authorities shall submit information about trends in livestock numbers and manure production for each livestock category in Denmark and in derogation farms according to Article

12 (h). All numbers have been brought to a round number in order to have a clearer picture.

The trends in livestock numbers (i.e. number of herds³) and manure production in kg N (until 2016/2017 in number of LUs⁴) for each livestock category and in derogation farms can be derived from the data shown in table 2.3. Over the planning periods from 2014/2015 to 2018/2019, the number of herds have decreased for each livestock category. The total number of Danish herds of livestock has decreased by ca. 16 % in between the planning periods of 2014/2015 and 2018/2019. From 2017/2018 the LUs is replaced by kg N.

Table 2.3: Number of Danish herds of livestock and production of manure in LUs
or in kg N per livestock category, rounded to the closest unit of 100 (1 LU=100 kg
N (ex storage))

Livestock category Year	Cattle total	Dero- gation cattle ⁵	Pigs	Fur and poultry	Sheep and goats	Others	Total	
2014/2015								
No. herds	12,300	1,500	4,100	2,000	2,400	6,100	26,900	
No. LUs	1,164,700	425,100	905,300	190,500	12,200	19,100	2,291,800	
2015/2016								
No. herds	11,800	1,500	3,900	2,000	2,300	5,800	25,800	
No. LUs	1,193,400	443,100	881,300	178,000	10,500	18,800	2,282,000	
2016/2017								
No. herds	11,500	1,400	3,600	2,100	2,200	5,600	25,000	
No. LUs	1,186,800	439,100	883,700	183,000	10,600	18,100	2,282,200	
2017/2018								
No. herds	10,800	1,300	3,400	2,000	2,000	5,500	23,700	
Kg N, mill.	115.2	39.6	80.0	20.2	1.0	2.2	218.6	
2018/2019								
No. herds	10,200	1,300	3,300	1,900	2,000	5,300	22,700	
Kg N, mill.	116.4	39.1	78.5	18.6	1.0	2.2	216.7	

³ The total number of herds does not coincide with total number of holdings in Denmark. A herd includes only one type of livestock and some holdings keep more than one herd, e.g. cattle and pigs.

⁴ One livestock unit is defined as 100 kg nitrogen in the livestock manure ex storage.

⁵ The amount of derogation cattle herds and LUs/kg N (organic fertiliser) are a part of "cattle total" and, thus, is not included in the summarization of herds and LUs/kg N (organic fertiliser) in "total".

3. Controls at farm level

Lars Paulsen & Lene Kragh Møller, the Danish Agricultural Agency, Ministry of Food, Agriculture and Fisheries of Denmark, November 2020

3.1 Control of compliance with the Danish derogation

According to Article 12 of Commission Decisions 2018/1928/EU, and 2020/1074/EU Denmark must submit a concise report on the evaluation practice, i.e. control at farm level, to the Commission every year.

The control of compliance with the Commission Decisions 2018/1928/EU and 2020/1074/EU follows two strategies:

- 1. Inspection of compliance with farm management, which is carried out during the year the farmer uses the derogation. This includes field inspections.
- 2. Control of the amount of livestock manure applied per hectare per year (control of compliance with the harmony rules), which is carried out after the derogation year has ended. This control is carried out as an administrative control of submitted fertilizer accounts.

3.2 Summary of inspection results 2020

Compliance with management conditions:

• Inspection at the farm in January and February 2020: 85 inspections were carried out. 85 holdings complied with the derogation management conditions, zero holdings got a remark in 2020 (**Table 3.1**).

Compliance with the harmony rules for holdings using the derogation:

- Administrative inspections of the submitted fertilizer accounts for 86 inspected farms in January and February 2019: 60 holdings complied with the specific rules for derogation holdings. Three holdings had minor violations and they received a warning. 21 holdings are still under investigation. Two holdings have gone bankrupt (**Table 3.2**).
- Administrative control of the submitted fertilizer accounts: 55 inspections were carried out, out of which 29 holdings complied with the rules and 26 holdings are still under investigation (**Table 3.5**).

3.3 Inspection of compliance within the derogation year

The farmers are required to fulfil certain conditions in order to use the derogation. The Danish Agricultural Agency has inspected the fulfilment of the Danish derogation conditions on derogation holdings from 2002/2003 through 2019/2020. Some conditions have to be checked on site at the farm (physical inspection), for example certain ploughing conditions, which are checked in January and February.

During the inspection at the farm, the inspector asks the following questions:

- 1. Does the farm have a yearly production of nitrogen in livestock manure above 300 kg of which at least 2/3 are from cattle (2/3 of the livestock units), i.e. is the farm mainly a cattle holding?
- 2. Has a plan been made for crops grown in the actual planning period?
- 3. Has the manager stated that the farm intends to comply with the 230 kg nitrogen per hectare per year derogation in the crop rotation plan?
- 4. Does the plan contain leguminous crops, e.g. red and white clover?
- 5. Has a declaration about (omitted) manure application been made?

- 6. Does the plan include ploughing grassland or grass catch crops in the planning period?
- 7. If the answer is "yes" in question 6: Have the fields already been ploughed by the time of inspection?
- 8. Does 80 % or more of the acreage available for manure application cultivated with crops with high nitrogen uptake and long growing season?

The inspection is based on 1) an interview with the farmer, 2) an inspection of the farms crop rotation plan for the previous and coming growing season and 3) a visual inspection of fields designated for ploughing.

At the inspection, the inspector draws up a report, which includes answers to the abovementioned questions. At the end of the inspection, the farmer is informed whether the holding is allowed to apply manure corresponding to 230 kg N/ha/year, i.e. whether the derogation can be used or not. If the holding is not complying with the derogation conditions, the holding is only allowed to apply livestock manure up to 170 kg N/ha/year. In this case, the farmer has to find other legal means of disposing the surplus manure produced on the farm.

If a farmer informs the inspector that the derogation will not be used, the field inspection is not carried out. An administrative control of the farm is carried out instead by the time the fertilizer account has been submitted. This control is carried out to secure that no more than 170 kg N/ha/year was applied.

The inspection report is submitted by the inspector to the headquarters of the Danish Agricultural Agency for possible further administrative inspection. The Danish Agricultural Agency verifies the data. Additional remarks made by the inspector, if any, are examined. This includes a process where the parties of interest are allowed to make statements on the case if an infringement is discovered.

3.4 Results

From 1 January until 1 March 2020, the Danish Agricultural Agency carried out 85 inspections on derogation holdings to inspect whether the conditions requirements were met. The control refers to the fertilizer accounts for the planning year 2018/2019 where some conditions are controlled in the next planning period 2019/2020. **Table 3.1** shows the results of the inspection for the last 17 years. Only very few remarks have been given and in general a good compliance with the rules has been noted.

Control planning- period ⁶	Total number of inspections	Inspections without re- marks	Inspections with remarks
2003/2004	35	29	6
2004/2005	46	46	0
2005/2006	50	49	1
2006/2007	50	49	1
2007/2008	54	54	0
2008/2009	47	46	1
2009/2010	51	49	2
2010/2011	50	50	0
2011/2012	54	52	2
2012/2013	49	49	0
2013/2014	47	46	1
2014/2015	49	49	0
2015/2016	48	48	0
2016/2017	49	48	1
2017/2018	90	87	3
2018/2019	86	86	0
2019/2020	85	85	0

Table 3.1: Results of on-site inspection of compliance within the derogation years during winter.

3.5 General inspection of the harmony rules

Harmony rules

Control of the harmony rules (i.e. the amount of livestock manure applied per hectare per year) on derogation farms is carried out after the derogation year has ended⁷. This control is carried out within the general inspection of the Danish harmony rules. The inspector visits the farm to inspect the production based on various production and fertilizer account documents. Violation of the harmony rules is sanctioned. For minor violations, the farmer receives a notification and recommendation or a warning. For more severe violations, the farmer is reports to the police and receives a fine. Farmers that receive a warning or a fine are reported for not complying with the cross compliance criteria.

Concerning the year 2017/2018, 84 derogation farms have been inspected for violation of the harmony rules. Holdings were automatically selected for inspection, based on a previously agreed set of "risk criteria". The Danish Agricultural Agency has therefore no direct influence on how many derogation holdings ware selected for "harmony rules inspection". Out of these derogation controls, 71.4 % (60 holdings)

⁶ The respective controls during the planning period 2019/2020, which have been performed in January and February 2020 are related to the fact that the farmer has made use of the derogation in the previous planning period, i.e. 2018/2019. This applies also to all previous control years.

⁷ This control population includes all holdings and not just cattle farms using the derogation. In this sample check, 55 cattle farms using the derogation have been selected. As consequence, further 55 cattle farms using the derogation have been controlled in relation to harmony rules in addition to the cattle farms controlled in accordance with Article 11.

were closed without remarks. Three holdings (3.6 %) had minor violations and they received a warning. 21 holdings (25.0 %) are still under investigation (**Table 3.2**).

Control planning- period	Total number of in- spec- tions	Inspec- tions without re- marks	Inspec- tions with minor viola- tions	Inspec- tions with fines	Inspec- tions still un- der in- vestiga- tion ⁸
2006/2007	65	59	0	5	1
2007/2008	27	22	2	2	1
2008/2009	32	26	1	5	0
2009/2010	27	24	1	2	0
2010/2011	37	35	0	0	2
2011/2012	52	50	0	2	0
2012/2013	43	40	0	3	0
2013/2014	29	27	0	1	1
2014/2015	30	29	0	0	1
2015/2016	28	24	0	2	2
2016/2017	86	85	0	0	1
2017/2018 ¹	84	60	3	0	21

Table 3.2 Results of inspection of compliance with the harmony rules for farms using the derogation.

¹ From 2017/2018 this is an administrative inspections of the submitted fertilizer accounts for the 86 inspected farms in January and February 2019. Two holdings have gone bankrupt.

Soil analysis

If the derogation is used for four consecutive years, the farmer must provide a soil analysis where phosphorous and nitrogen levels in the soil are examined. One sample per five hectares must be provided.

In Denmark, the soil analysis for phosphorus (the "P-tal") indicates the soil's phosphorus status and hence approximates the level of phosphorus in the soil available for uptake by the crop. Internationally, the soil analysis is referred to as "Olsen-P". Olsen-P is often expressed in mg P per kg soil. In Denmark, however, the "P-tal" is expressed in mg P per 100 g soil. Olsen-P in Danish agricultural soil is in average around 40 mg P per kg soil (P-tal = 4.0). Only a part of the inorganic phosphorus available for the crop is extracted from the soil sample, when the phosphorus status is determined. This extractable part accounts for approximately 5 to 10 per cent of the total phosphorous content of the soil. A P-tal between 2 and 4 is generally accepted as a sufficient level for most crops and 2-2.5 is the lower critical soil P level. A P-tal above 6 is considered very high.

The N-total analysis is used to determine the amount of extra fertilizer to be added to meet the nutrient demand of the crop. The total soil N content (N-total) describes the N pool in the soil, which potentially is available to the crops as a result of slow mineralization. In Denmark, depending on the C/N ratio in the soil, the standard N-total is 0.13 %. The farmer cannot expect any N-supply from mineralization, if the level of 0.13 % N-total is found. If the value is above 0.22 %, the level is high and expected

⁸ I.e. inspections still under investigation at time of reporting. Numbers of inspections still under investigation prior to 2017/2018 are not updated. Thus, these inspections may have been finalized.

mineralization is (accounted for with) 40 kg N in maize and cereals per hectare. The N-total standard for grass fields is 0.18-0.22 %, and if the value is above 0.22 %, the expected mineralization is (accounted for with) 10 kg N per hectare.

Results of soil analyses from derogation farms

The sampling and analyses must be carried out at least once every four years (prior to 2012/2013, the requirement was at least once every three years). The results of the development of compliance with the requirement of soil analysis are shown in Table 3.3.

The inspection of derogation farms for 2017/2018 showed that 45.3 % provided soil analysis. No holdings got a remark regarding soil analysis.

Control planning- period	Number of in- spections for soil analysis	Inspections without remarks	Inspections with remarks/still under investiga- tion
2004/2005	74	71	3
2005/2006	18	16	2
2006/2007	39	34	5
2007/2008	16	12	4
2008/2009	22	18	4
2009/2010	11	9	2
2010/2011	14	13	1
2011/2012	35	35	0
2012/2013	30	27	3
2013/2014	15	14	1
2014/2015	22	21	1
2015/2016	11	11	0
2016/2017	41	41	0
2017/2018	39	39	0

Table 3.3: Results of inspection of compliance with the soil analysis requirement.

The results of the soil analyses for phosphorus and nitrogen on derogation farms are shown in **Table 3.4**.

Control pl period	anning-	2011/ 2012	2012/ 2013	2013/ 2014	2014/ 2015	2015/ 2016	2016/ 2017	2017/ 2018
P tal	Aver- age	4.36	4.60	4.33	4.60	4.62	4.29	4.22
(mg P/100 g soil)	Mini- mum	2.00	2.90	2.90	2.87	3.10	2.39	2.20
	Maxi- mum	6.40	6.10	8.40	6.08	6.14	6.95	7.05
N-total	Aver- age	0.60	0.33	0.25	0.25	0.23	0.21	0.20
(%)	Mini- mum	0.11	0.12	0.15	0.13	0.13	0.11	0.12
	Maxi- mum	2.39	1.71	0.41	0.58	0.41	0.59	0.34
N in grass	Aver- age	0.36	0.24	0.48	0.24	0.24	0.22	No soil analy- ses
(%)	Mini- mum	0.01	0.17	0.16	0.16	0.17	0.13	
	Maxi- mum	1.10	0.35	2.00	0.51	0.33	0.36	

Table 3.4: Phosphorus ("P-tal" after Olsen-P-extraction) and nitrogen levels in soil analyses, given as average of all inspected holdings (n=39 in 2017/2018) and with the lowest and highest average values at holding scale, respectively.

3.6 Control of fertilizer accounts

Each year, the farmers submit their fertilizer accounts to the Danish Agricultural Agency. The accounts include key data on:

- total arable land on the farm
- arable land available for application of livestock manure
- data on catch crops
- type and number of livestock
- production of livestock manure (kg N and P)
- usage of livestock manure including manure from contractors
- usage of fertilizers and organic matter other than livestock manure
- the farms nitrogen quota and the average phosphorus ceilings for different livestock manure, fertilizers and organic matter other than livestock manure
- information on whether the farmer has used the derogation or not

For the year 2017/2018, 1,461 (4.4 %) of the submitted fertilizer accounts were subject to administrative inspection. 365 fertilizer accounts remain to be investigated. The data was verified and the parties of interest were allowed to comment on their cases. The accounts were selected based on different risk criteria. In 2017/2018, 55 (3.8 %) derogation holdings were selected for control. The holdings were asked to

submit their updated and valid fertilization plan and to state their manure application. It was checked whether the crop rotation plan included at least 80 % crops with high N-uptake and long growing season as well as whether leguminous plants were included. If the derogation had been used for four consecutive years, the farmer also had to submit the results of the soil analysis. The share of cattle- and other animal kg N on the farm was also controlled.

Results

Out of the 55 harmony controls, 29 holdings (52.7 %) were closed without remarks and 26 (47.3 %) inspections are still under investigation (**Table 3.5**).

Table 3.5: Results of administrative inspection of compliance with the harmony	
rules of farms using the derogation.	

Control plan- ning- period	Number of inspections	Inspections without remarks	Inspections with remarks	Inspections still under investigation
2009/2010	38	34	0	-
2010/2011	68	68	0	-
2011/2012	40	39	1	-
2012/2013	62	58	1	3
2013/2014	34	24	4	6
2014/2015	62	30	4	28
2015/2016	61	46	6	9
2016/2017	46	31	3	12
2017/2018	55	29	0	26

4. Water quality

Jonas Rolighed, Gitte Blicher-Mathiesen, Department of Bioscience, Aarhus University, June 2021

With Commission Decisions 2005/294/EC, 2008/664/EC, 2012/659/EU, 2017/847/EU, and 2018/1928/EU Danish cattle holdings are permitted to derogate from the general rules in the Nitrates Directive (91/676/EEC). Cattle holdings encompassed by derogation shall cover 80% or more of the acreage available for manure application by cultivated crops having high nitrogen uptake and a long growing season.

According to Article 10(1), Article 10(2), Article 10(3), and Article 10(4) of Commission Decision 2018/1928/EU, Denmark shall each year:

- Deliver maps at municipality level, showing the percentage of farms, the percentage of the livestock and the percentage of agricultural land with derogation.
- Provide continuous data about crop rotations and agricultural practices from farms with derogation.
- Provide continuous analysis of level and trends in nitrate and phosphorus concentrations in root zone water, surface waters and groundwater within the framework of the agricultural national monitoring programme on sandy and loamy soils for farms under both derogation and non-derogation conditions.
- Quantify the percentages of the land under derogation covered by: (a) clover or alfalfa in grassland; (b) barley and pea under sown with grass.

According to Article 10(2), the monitoring sites shall be representative of the main soil types, the prevalent fertilization practices and the main crops. Reinforced monitoring shall be conducted in agricultural catchments on sandy soils. In addition, nitrate concentrations in surface and groundwater shall be monitored for at least 3% of all farms benefiting from authorisation of derogation.

The competent authorities shall carry out surveys and continuous nutrient analyses in the agricultural catchment national monitoring programme and shall provide data on local land use, crop rotations and agricultural practices on cattle farms benefiting from an authorisation.

In Article 10(3), it is stated that competent authorities shall carry out surveys and continuous nutrient analyses in the agricultural catchment within the framework of the national monitoring programme and provide data on local land use, crop rotations and agricultural practices on cattle farms benefiting from an authorisation of derogation.

Moreover, information and data collected from nutrient analyses and from monitoring shall be used for model-based calculations of nitrogen and phosphorus losses from cattle farms benefitting from an authorisation of derogation.

In Article 10(4), it is stated, as mentioned before, that competent authorities shall quantify the percentages of the land under derogation covered by: (a) clover or alfalfa in grassland and (b) barley and pea under sown with grass.

This chapter covers the requested reporting in Article 12 (b-f) on:

- Results of ground and surface water monitoring as regards nitrate and phosphorus concentrations, including information on water quality trends, for farms under both derogation and non-derogation conditions, as well as the impact of derogation on water quality, as referred to in Article 10(2).
- The results of soil monitoring as regards nitrogen and phosphorus concentrations in the root zone water for farms under both derogation and non-derogation conditions, as referred to in Article 10(2).
- Results of the surveys on local land use, crop rotations and agricultural practices, as referred to in Article 10(3).
- Results of model-based calculations of the magnitude of nitrogen and phosphorus losses from farms benefitting from an authorisation of derogation, as referred to in Article 10(3).
- Tables showing the percentage of agricultural land under derogation covered by clover or alfalfa in grassland and by barley/pea under sown with grass, as referred to in Article 10(4).

So far, model-based calculations of phosphorus losses from farms benefitting from an authorisation of derogation are not available, but measured phosphorus concentrations in root zone water on fields with average application of less and more than 170 kg organic N ha⁻¹ are presented.

As data in this chapter are from the year 2019, the Commission Decision 2018/1928/EU covers this period.

4.1 Introduction

Since the late 1980s, Denmark has yielded a comprehensive and efficient effort to improve the environmental state of groundwater and surface water by lowering nitrate concentrations, especially through reductions in nitrate leaching from agricultural sources. The first Action Plan on the Aquatic Environment was adopted in 1987 and has since then been followed by subsequent action programmes to ensure efforts are made to reduce the loss of nitrogen and phosphorus to the aquatic environment.

In 1998, the Action Plan on the Aquatic Environment (APAE) II was accepted by the EU Commission as the Danish Nitrate Action Plan implementing the Nitrates Directive (1998-2003). In 2003, a final evaluation of Action Plan II was performed. The results showed a 48% reduction of the nitrate leaching from the agricultural sector, thus fulfilling the reduction target set in 1987. In the 5-year period 2001-2005, the total flow-normalised nitrogen load to marine waters ranged within the interval 62,000 to 70,000 t N.

In the subsequent action plans, the Green Growth Agreement from 2009, the first and the second River Basin Management Plan from 2014 and 2016 as well as the Food and Agricultural Agreement in December 2015, further mitigation measures were suggested in order to fulfil reduction targets for the N load to marine areas and the targets of the Water Framework Directive.

In 2015, Denmark implemented the EU Greening component under CAP direct payments (REG EU 1307/2013), implying that at least 5% of the arable land of farm holdings shall be appointed an ecological focus area with a greening element such as set aside, catch crops etc.

Establishment of 50,000 ha of obligatory buffer zone placed approximately 10 m from the edge of open streams and lakes larger than 100 m^2 was decided to be imple-

mented from the autumn 2012. In these buffer zones, application of fertilizer is prohibited and soil cultivation must not take place. The area with buffer zones was adjusted from 50,000 to 25,000 ha later in 2014, and from the beginning of 2016 the additional buffer zones are no longer mandatory and restricted to the former requirements of 2 m buffer zones along target streams and lakes larger than 100 m², amounting to approximately 6,000 ha.

The Political Agreement on Food and Agricultural Package from December 2015 includes a diverse package of measures aimed to change the environmental regulation of the agricultural sector. The first part of this political agreement was implemented from 2016.

In 2016, farmers were allowed to use more fertilizer. According to the APAE II agreement, farmers were restricted in the application of fertilizer at a level that was lower than the economic optimum. This measure in APAE II was set to reduce the fertilizer application of nitrogen to 10% below this optimum. This rule was regulated so that the total national nitrogen quota was set to a fixed level but with the possibility of an adjustment relative to changes in crop cover. This adjustment made sense as crops having a high application standard also have a higher nitrogen uptake. If crops such as grass increase in cover, then the fertilizer application and N quota will increase as well. However, due to the suspension of set aside in 2008, higher yields and increases in prices of cereals and proteins, the gap between the economic optimum and the national N quota increased, especially after 2008, amounting to 18% in 2015.

According to the Political Agreement on Food and Agricultural Package implemented in 2016, extra fertilizer application, amounting to 2/3 of the gap between the economic optimum and the reduced N quota, were allowed. From 2017, famers were allowed to apply nitrogen up to the economic optimum. Additional cover of catch crops and the greening element, for instance more catch crops and set-aside, were, among other measures, introduced to counteract the potential increase in leaching due to the extra application of fertilizer from 2016 and onwards.

Additionally, targeted catch crops of 145,000 ha were implemented in 2017 to counteract the potential increase in leaching due to the extra application of fertilizer in 2017. In 2018 and 2019, the need for targeted catch crops was approximately 114,000 and 139,000 ha, respectively. The targeted catch crops scheme was introduced to ensure that the status of coastal waters and groundwater does not deteriorate. Therefore, targeted catch crops are established in catchments where reduction of the nitrogen load is needed. Applicants for targeted catch crops could be all farmers who either own or lease fields in such small catchments for cultivation.

The second River Basin Management Plans (RBMPII) was adopted in June 2016, proposing schemes for implementation of mitigation measures, such as re-establishment of riparian areas, construction of wetlands, set aside of organic soils, afforestation and adjustment of greening elements, to obtain an annual reduction in the marine N load of 6,900 t N in the period 2015-2021 (SVANA 2016). However, the actual decision on which measures to initiate to reach an annual reduction in the nitrogen load of 6,200 t N has been postponed to after 2021.

The N load to marine waters has been reduced incrementally along with the successful implementation of measures from point sources and agriculture. Approximately half of the Danish land area is located within catchments equipped with stream water gauging stations where the N load to marine areas is regularly measured (Kronvang et al., 2008). The nitrogen load for ungauged catchments has been modelled using an empirical model (Windolf et al., 2011) and shows that the annual load to marine waters varied between 55,000 and 59,000 t N, yielding an average of 57,000 t N for the five years (2010-2014) used as reference level in the RBMPII (SVANA (2016), Wiberg-Larsen (2015)). However, the calculation of this total nitrogen load to coastal areas has been updated and now includes a higher proportion of gauged catchments as well as an improved and more detailed calculation of discharge in ungauged catchments (Thodsen et al., 2020). The updated calculation yields an annual flow-normalized nitrogen load ranging between 51,900 and 57,700 t N, with an average of 53,500 t N for the period 2010-2014. For the period 2015-2019, the flow-normalized total nitrogen load has varied between 53,400 and 61,000 t N with the lowest level in 2015 and highest value in 2019 following a year with drought related low crop harvest.

The regulation and effects described in this chapter cover the period until and including 2019. Additional agricultural regulation, such as requirement to increase the utilization efficiency of nitrogen in manure and reduced fertilizer application norm on soil with high content of organic matter, are implemented from 2020 and 2021.

The remaining part of this chapter is divided into three parts:

<u>First</u>, the general development in agricultural practices at national level is presented for the period 2005-2019. This analysis is based on national register datasets from the Ministry of Food, Agriculture and Fisheries (until November 2020 part of the Ministry of Environment and Food), i.e. the single payment register and the fertilizer accounts.

<u>Second</u>, modelled nitrate leaching, including crop distribution and nitrogen balances, is presented for various farm types and geographical areas, and the impact of derogation farms is analysed based on a dataset derived by linking data from the single payment register, including data on the crops on each field comprised by the holdings, and the fertilizer accounts. Both datasets cover agriculture in the year 2019.

<u>Third</u>, measurements of water quality from the National Monitoring Programme are presented for the period 1990/91-2018/19, with particular reference to the Agricultural Catchment Monitoring Programme (Blicher-Mathiesen et al., 2021). This section includes:

- Modelling of nitrate leaching in the agricultural monitoring catchments.
- Measurements of nitrate in water leaving the root zone, including fields receiving more than 170 kg N ha⁻¹ in organic manure.
- Nitrogen in surface water, draining from agricultural catchments.

Modelling of nitrate leaching in this report is carried out by means of the empirical model N-LES (version 4) from 2008 (Kristensen et al., 2008). This model is partly based on data from the Agricultural Catchment Monitoring Programme. The model requires input data for agricultural practises (N fertilization, cropping system), soil data and water percolation from the root zone. Percolation is calculated using the Daisy model and a standard climate from a 10 km grid net (Danish Meteorological Institute), representing weather measurements from the period 1990-2010. The climate dataset contains dynamic correction factors for rainfall (Refsgaard et al., 2011). Thus, modelled nitrate leaching represents the leaching in a standardised climate (water percolation). In contrast, all measurements from the Agricultural Catchment Monitoring represent nitrate leaching under the actual climatic conditions.

4.2 Development in agricultural practices at the national level from 2005 to 2019

Crop distribution

The development in crop distribution for 2005-2019 was analysed on the basis of the single payment registration. **Figure 4.1** presents the results for cash crops, fodder crops and non-cultivated areas. The year 2005 was the first year with single payment, and it was anticipated that the reporting of areas for this first year would be overestimated. Hereafter, the total reported agricultural area, including set aside, decreased from approximately 2,757,000 ha in 2006 to 2,600,000 ha in 2019.

The decrease in agricultural area of about 12,000 ha per year is due to road construction, afforestation, urbanisation etc. During the years 2006-07, set-aside comprised about 160,000 ha. As from 2008, the set aside obligation was suspended, and in 2008 and 2009 set aside areas were converted to cash crop, fodder crops and nature-like areas. Set aside covered between 23,000 and 32,000 ha in the period 2015-2019 as set aside is an element in the Danish implementation of the EU Greening. The area with cash crops and fodder crops has decreased slightly since 2012.

Catch crops

In Action Plan III, the requirement for growing catch crops was carried over from the former Action Plan, stipulating that farmers in 2005-2009 should grow catch crops on at least 6% of the potential catch crop area if they applied less than 80 kg organic manure N ha⁻¹ and on 10% of the area if they applied more than 80 kg organic manure N ha⁻¹. In 2008, the requirement for growing catch crops was raised to counterbalance the effects of the set aside suspension. From autumn 2009, an additional catch crop area, equivalent to an extra 4% of the potential catch crop area, was implemented, yielding a total requirement for the growing of catch crops of 10% or 14%, respectively. Hence adjusted to 10.7 and 14.7% from 2020.

During this period (2005-2010), farmers growing winter crops (wheat, rye, winter barley, oilseed rape) preventing fulfilment of catch crop requirements, were granted a reduction in the required catch crop area. From 2011, this possibility ceased.




Figure 4.1 Development in crop distribution at the national level from 2005 to 2019, data from the single payment register.

At the same time, voluntary alternatives to catch crops were introduced such as:

- Reduction of the farm nitrogen quota
- Growing of special crops between harvest and sowing of winter crops
- Growing catch crops on other farms
- Establishment of perennial energy crops
- Separation and treatment of animal manure (biogas and burning of the solid fraction of manure)
- From 2015, substitution of one ha of catch crop by four ha of set aside near open streams and lakes above 100 m² and located next to agricultural areas in rotation
- From 2014, substitution of one ha of catch crop by four ha of winter cereals, if sown earlier than September 7

• From 2017, substitution of one ha of catch crop by one ha of set aside

Data from the fertilizer accounts show that establishment of catch crops increased from about 118,600-138,000 ha in 2005/06-2007/08 to about 355,600 ha of catch crop equivalents in 2019/20 (**Table 4.1**). The introduction and use of catch crop alternatives were equivalent to the effect of 13,900-43,000 ha catch crops in the period 2011/12-2019/20.

Table 4.1 Area with catch crops and catch crop alternatives (1,000 ha of catch crop equivalents) reported by the farmers in the annual fertilizer account in the period 2005/06-2019/20.

	05/06	06/07	07/08	08/09	09/10	10/11	11/12	12/13	13/14	14/15	15/16	16/17	17/18	18/19	19/20
Catch crops	138.0	118.6	127.2	196.6	183.0	211.0	211.0	224.0	295.7	321.1	390.0	353.1	415.2	366.5	355.6
Catch crop alternatives	0	0	0	0	0	0	28.6	44.0	13.9	43.3	37.6	36.1	28.5	42.8	16.2

In 2017, a new regulation of animal husbandry was implemented. With this regulation, additional catch crops were to be established in certain areas on certain farms using organic fertilizers, including livestock manure. The regulation applies only to farms cropping more than 10 ha and with the use of organic fertilizer of > 30 kg N ha⁻¹. In addition, the cropped area must be located in catchments with an increasing use of organic fertilizers that drains into nitrate sensitive types of nature habitats of Natura 2000 area or to specific voluntary areas in near coastal waters with a need for N load reduction according to the River Basin Management Plans. The additional catch crops in certain areas on certain farms using organic fertilizer can replace all or a part of the need for 80% fodder crops on derogation farms and catch crops grown to fulfil the EU greening requirements.

Consumption of nitrogen fertiliser and nitrogen in manure

Data on the annual use of inorganic fertilizers and the use of nitrogen in animal manure are obtained from the fertilizer accounts (**Table 4.2**). In previous derogation reports, data on animal manure were based on the manure production for different categories of livestock. As these data are no longer recorded, the manure application data are instead based on the data from the fertilizer account.

The application of animal manure variated between 216,000 and 227,000 t N from 2005 to 2019. The use of inorganic fertilizers amounted to about 181,000-202,000 t N year-1 in 2005-2007 and increased to 205,000 and 209,300 t N year-1 in 2008 and 2009, probably due to the cultivation of previous set aside areas. This was expected to be a temporary effect as the procedure for setting the crop nitrogen standards implies that an increase in agricultural area with fertilizer requirements must be followed by an equivalent reduction in nitrogen standards. Administratively, however, this reduction is based on statistical data on the cultivated area, resulting in a delay of two years. Thus, in 2010-2014, the use of inorganic fertilizers decreased again, reaching the same level as in 2005-2007. The use of inorganic fertilizer increased from 210,000 t N in 2015 to 242,000, 237,000 and 224,000 t N in 2016, 2017 and 2018, respectively, after the implementation of the Food and Agricultural Package, according to which farmers were allowed to use more fertilizer after 2015. The lower use of inorganic fertilizer in 2018 compared to the two former years is caused by an increase in organic farming, farms that do not use inorganic fertilizer, as well as a decrease in the cultivated area. A change in the crop distribution with

higher cover of spring cereal on the cost of winter cereals also contribute to a lower use of inorganic fertilizer of approximately 20,000 t N in 2018, as winter cereal have a higher N uptake, higher harvest yield and therefore a higher economic optimal standard than spring cereal. The use of inorganic fertilizer amounted to 223,000 t N in 2019 as almost the same level as in 2018, hence taking into consideration that farmers were recommended to apply app. 4 kg N ha⁻¹ lower amount of fertilizer as a relevant amount of nitrogen was still remained in the soil in spring due a very dry autumn and winter.

Table 4.2 Development in the use of inorganic nitrogen fertilizer and of nitrogen in animal manure as reported by the farmers in the annual fertilizer status accounts for the period 2005-2019 (1,000 t N a⁻¹).

	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Ferti- lizer	191	181	202	205	209	198	203	198	199	203	210	242	237	224	223
Animal ma- nure	227	218	236	230	226	224	223	220	215	212	216	219	218	224	219

4.3 Modelled nitrate leaching for farm types and geographical areas and the impact of derogation farms at the national level – 2019 data

Modelled nitrate leaching demonstrates the effect of crop distribution, nitrogen input, soil type and water percolation through the soil. This section includes a presentation of all of these parameters. The analyses are based on the national datasets from the single payment register and the fertilizer accounts. However, before data can be used for this purpose, a detailed compilation of the two datasets must be made (Børgesen et al., 2009). The single payment register contains information on crops at field-block level, and the fertilizer accounts contain information on the use of nitrogen (inorganic fertilizer and organic manure) at farm level. The two datasets are linked by means of the common farm identity number or a common farm address, and the reported amounts of fertilizer and manure from the individual accounts are distributed on the fields of each farm according to the crop nitrogen standards. Hereby, we obtain a dataset with coherent data on crops and nitrogen application at field level. We have no information on grass-ley from either dataset. Therefore, we estimate this parameter based on the area with rotation grass, assuming a conversion rate of three years. If there is not sufficient space in the crop rotation, the area with grass-ley is reduced accordingly. Data on catch crops is derived from the fertilizer accounts.

The field-blocks are geographically mapped, implying that each field can be linked to soil maps and to the meteorological grid net. Having established the soil type for each field-block, the standard harvest yield may be estimated. Furthermore, nitrogen fixation is included using standard values for each crop. This final dataset now contains all information necessary for geographically distributed computation of crop coverage and field nitrogen balances and for modelling nitrate leaching.

Farm type

The data is divided into three main groups of farm type – arable farms, pig farms and cattle farms. A pig farm is defined as a farm where more than 2/3 of the used organic

N including manure originate from pigs, and a cattle farm is defined as a farm where at least 2/3 of the used organic N including manure originate from cattle. An arable farm is a farm with a production of organic N including manure of less than 20 kg N. The farm may import animal manure, which will appear in the fertilizer account and is therefore included in this analysis. Other farm types are not included in this analysis.




Figure 4.2: Crop distribution for three main farm types in 2019. Combined dataset from the single payment register and the fertilizer status accounts.

Figure 4.2 shows that arable farms and pig farms grew cereals, particularly winter wheat, on the majority of the agricultural area (48 and 64%) in 2019. Other major cash crops were oilseed rape, peas, root crops (potatoes and sugar beet) and grass for seeds (17-18%). Cereal silage, grass and maize constituted a minor part of the area (4-10%). Catch crops were grown on 9-13% and grass-ley on 2-3% of the agricultural area on arable and pig farms.

Cattle farms have a different crop rotation. Cereals and other cash crops were grown on 39% of the area, whereas cereal silage, grass and maize were grown on 60% of the area. Fodder beet was grown on 1.2% of the area. In addition, grass-ley was found on 10% and catch crops on 15% of the area.

On arable farms, an average amount of about 47 kg N ha⁻¹ from animal manure was applied. For pig and cattle farms, the amounts were, respectively, 100 kg N ha⁻¹ and 124 kg N ha⁻¹ (**Table 4.3**).

			Root	Root zone water								
	Comm. ferti- lizer	Ani- mal ma- nure	Other org.	N fix.	N depos.		Total input			Percol.	Nitrate leach- ing	
				(kį	g N ha-1 a-1)					(mm a-1)	(kg N ha-1)	(mg l-1)
Arable	104	47	5.6	8.7	12	2.0	179	115	64	335	54	71
Pigs	85	100	1.8	5.5	12	2.2	207	120	87	376	65	76
Cattle	72	124	1.4	29	12	1.5	240	147	93	408	64	70

Table 4.3 N inputs, N balances and nitrate leaching and nitrate concentration at
the bottom of the root zone for three main farm types in 2019. Combined dataset.

The use of inorganic fertilizers decreased with increasing application of animal manure. Total inputs of nitrogen from inorganic fertilizer, manure, other organic sources, N fixation and atmospheric deposition amounted to 179, 207 and 240 kg N ha⁻¹ for arable farms, pig farms and cattle farms, respectively. N balances, calculated as the difference between the total input of nitrogen and removal by harvested crops, were 64, 87 and 93 kg N ha⁻¹ for arable farms, pig farms and cattle farms, respectively. As expected, modelled nitrate leaching was lower from arable farms (on average 54 kg N ha⁻¹) than from animal husbandry farms (65 kg N ha⁻¹ from pig farms and 64 kg N ha⁻¹ from cattle farms). N leaching was, on average, 1 kg N ha⁻¹ lower for cattle farms than for pig farms. A major reason for this is that cattle farms grow a high proportion of fodder crops that have a long growing season and therefore a larger N uptake.

On arable farms, the modelled nitrate leaching amounted to 84% of the N balance, which is high relative to the 75% recorded for pig farms and the 69% observed for cattle farms. An explanation may be that leaching on these soils with low input of organic manure is affected by mineralisation of the organic pool, i.e. depletion of the total soil N content. However, the high leaching fraction may also be caused by the uncertainties associated with the two separate calculations of the N leaching and N balance.

Water percolation through the soil is considerably higher on cattle farms than on arable and pig farms. However, this is not due to the differences in farm type but the fact that the cattle farms are located mainly in the western part of the country with more sandy soil and higher rainfall and a consequently higher percolation. The higher percolation on cattle farms leads to dilution of the nitrate concentration in the soil water. Thus, the modelled average nitrate concentrations in soil water were 71 and 76 mg NO₃ l⁻¹ on arable and pig farms, respectively, and 70 mg NO₃ l⁻¹ on cattle farms for the year 2019.

Geographical areas

Farm types are not evenly distributed throughout the country because of variations in farming conditions. For the following analysis, Denmark has therefore been divided into five farming regions (**Figure 4.3**).




Figure 4.3 Farming regions in Denmark with different soil types, farming practices and rainfall and the location of the six monitored agricultural catchments.

Table 4.4 shows that Zealand is dominated by arable farming, whereas Eastern (E) Jutland and Funen are dominated by arable farming and pig production. Finally, North (N), North-West (NW) and West (W) Jutland have the highest density of cattle farming. Thus, arable and pig farms are located mainly in the eastern part of Denmark on loamy soils and with low rainfall, whereas cattle farms are located mainly in the northern and western parts of Denmark on sandy soils and with higher rainfall, the rainfall increasing from north to south.

	Arable	Pig	Cattle	Other	Sand	Loam	Organic soils	Percol.
		% of agric	ultural area		% o	f agricultu	ral area	mm/year
Zealand	61	14	15	10	4	93	3	199
Jutland E + Funen	38	26	24	13	25	71	4	334
Jutland N	31	18	35	16	79	10	11	363
Jutland NW	25	22	38	15	61	33	6	449
Jutland W	29	16	40	15	75	19	6	540

Table 4.4 Distribution of farm types and soil types and water percolation through the soils in Denmark divided into five main geographical areas – 2019.




Figure 4.4 Crop distribution for five farming regions in Denmark in 2019. Combined dataset from the single payment register and the fertilizer accounts.

The crop distribution within the five farming regions of Denmark follows the same pattern as for farm types, i.e. mainly cereals and other cash crops on the islands and in Eastern Jutland and cereals and fodder crops in West and North Jutland (**Figure 4.4**).

The input of nitrogen with animal manure, the total nitrogen input and the field nitrogen balances are lowest on Zealand, higher in E Jutland and on Funen and highest in W, NW and N Jutland (**Table 4.5**). In the latter three areas, the average nitrogen input varied between 204 and 214 kg N ha⁻¹. The average modelled nitrate leaching generally increased from east to west due to increases in nitrogen input and percolation. Within the three western and northern parts of Jutland, the nitrate leaching increased from northern to southern Jutland, mainly due to increased water percolation through the root zone. Higher water percolation led to dilution of the nitrate concentrations of the soil water, resulting in an average nitrate concentration in soil water of 79, 72, 75, 70 and 62 mg NO₃ l⁻¹ on Zealand, Funen + E and N Jutland, and NW and W Jutland, respectively.

Table 4.5 N inputs and N balances, nitrate leaching and nitrate concentration at
the bottom of the root zone calculated for five geographical areas in Denmark in
2019. Combined dataset from the single payment register and the fertilizer ac-
counts

				N	balanc	e				Root	t zone wa	ater
	Comm. fertiliser	Animal manure		N-fix.	N- depos.	Seeds	Total input	Har- vest	N bal- ance	Percol.	Nitrate leaching	NO ₃ - conc
			0	kg	N ha-1 a	-1	•			mm a-1	kg N ha-1	mg l-1
Zealand	107	35	5.1	9.3	10	1.8	168	119	49	199	35	79
Jutland E +Funen	90	71	2.8	12.0	12	1.9	189	122	67	334	54	72
Jutland N	71	97	1.8	20.4	12	1.6	204	125	79	363	61	75
Jutland NW	70	105	0.8	17.7	12	1.7	207	127	80	449	71	70
Jutland W	68	107	3.5	19.8	14	1.9	214	133	81	541	76	62

Derogation farms

Derogation farms are mainly located in N, NW and W Jutland where cattle farming is dominant. The effect of the derogation was evaluated for these three geographical areas. The cattle farms were grouped into four livestock density groups depending on the application of organic N including manure: 0-100, 100-140, 140-170 kg N ha⁻¹ and derogation farms with the use of organic N including manure of 170-230 kg N ha⁻¹.

The crop distributions for the three geographical areas were found to be almost identical, with some differences in cover between spring and winter cereals as well as larger proportion of maize and catch crops in W Jutland (**Figure 4.5**). There is a clear trend indicating a decrease in areas with cereals and increase in the areas with cash crops with increasing livestock density. In addition, the area with fodder crops increases with increasing livestock density. The area with roughage amounted to 54, 65 and 73% for the three groups, 0-100, 100-140, 140-170 use of organic N including manure ha⁻¹, respectively, whereas derogation farms grew roughage on an average of 90% of the area.

The effect of derogation on nitrate leaching was evaluated separately for the three geographical areas. The nitrogen input as well as the field nitrogen balances increased with increasing livestock density (**Table 4.6**). Modelled nitrate leaching is generally a combined effect of two opposing mechanisms – an increase in leaching due to increased nitrogen input and a decrease in leaching due to an increased area with roughage and catch crops. **Table 4.6** shows that the modelled nitrate leaching generally increased with increasing livestock density and hence with increasing nitrogen input. Thus, differences occurred in the modelled annual nitrogen leaching of 8, 6 and 9 kg N ha⁻¹, respectively, between derogation farms and farms using 140-170 kg N ha⁻¹ of organic N in the three Jutland regions. Similarly, modelled nitrate concentrations in the soil water leaving the root zone were 10, 10, and 6 mg NO₃ l⁻¹ higher for derogations farms than for cattle farms using 140-170 kg organic N ha⁻¹ in W, NW and N of Jutland, respectively.




Figure 4.5 Average crop distribution for four groups of livestock density calculated for the total use of organic nitrogen including manure (kg N ha 1) in N, NW and W Jutland in 2019. Combined dataset from the single payment register and the fertilizer accounts.

The use of legumes (clover, alfalfa, peas) in grass and cereal silage is shown in **Table 4.7**. The general trend is that derogation farms grow less legumes than non-derogation farms (**Table 4.7**). Thus, clover or alfalfa (max. 50% share) in rotation grass was used on 76% of the rotation grass area for derogation farms and on 81-90% for non-derogation farms. For permanent grass, the equivalent values were 20% for derogation farms and 27-39% for non-derogation farms. Cereal silage with peas amounted to 6% of the silage area for derogation farms and 13-18% for non-derogation farms.

Table 4.6 N inputs, N balances and nitrate leaching and nitrate concentration at the bottom of the root zone calculated for four groups of livestock density at cattle farms and for three geographical areas in Jutland, Denmark, 2019. Combined dataset from the single payment register and the fertilizer accounts.

					N	balanc	e				Root	zone w	ater
Re- gion	Annual use of organic N	Comm. ferti- lizer	Ani- mal ma- nure	Other org. N		N depos.		To- tal in- put	Har- vest	Bal- ance	Percol.	Nitrate leach- ing	NO ₃ - conc
	kg N ha-1				kg	N ha-1 a-1	1				mm a-1	kg N ha-1	mg l-1
Jut- land	0-100	81	57	1.6	22	12	1.2	175	115	60	359	54	67
Ν	100-140	61	119	0.5	34	12	1.3	228	138	89	366	60	72
	140-170	58	153	0.3	44	12	1.2	268	159	109	354	63	79
	170-230	72	199	0.0	39	12	1.4	323	187	136	348	71	89
Jut- land	0-100	83	63	2.5	20	12	1.4	183	116	66	423	57	60
NW	100-140	58	122	0.2	31	12	1.3	224	133	91	445	65	65
	140-170	59	151	0.2	37	12	1.3	261	155	106	449	74	73
	170-230	70	193	0.0	35	12	1.5	312	180	132	425	80	83
Jut- land	0-100	73	60	6.4	21	14	1.6	176	118	59	521	58	49
W	100-140	51	123	2.5	36	14	1.3	228	142	86	541	71	58
	140-170	57	153	0.5	37	14	1.5	263	158	105	543	81	66
	170-230	77	201	0.2	29	14	1.6	323	187	137	548	90	72

Table 4.7 Use of legumes in grass and cereal silage at cattle farms for derogation and non-derogation farms 2019.

	<u>Use of</u>	organic N, ind	cluding manur	e (kg N ha ⁻¹ a ⁻¹)
	0-100	100-140	140-170	170-230
		share of ag	ricultural area (%	6)
Rotation grass	13.8	23.8	28.0	32.6
		share of	rotation grass (%)
No clover/alfalfa	17	9	10	24
< 50% clover/alfalfa	81	90	90	76
> 50% clover/alfalfa	2	1	0	0
		share of a	gricultural area (S	%)
Permanent grass	16.1	11.3	8.7	5.9

		share of p	ermanent grass ((%)
No clover/alfalfa	60	71	72	80
< 50% clover/alfalfa	39	29	27	20
> 50% clover/alfalfa	0	0	0	0
	gricultural area (%)		
Cereal silage	1.9	5.6	6.8	9.0
		share of	cereal silage (%))
No legumes	82	63	68	94
< 50% legumes	13	18	15	6
100% legumes	5	18	17	0

4.4 Development in modelled nitrate leaching in the Agricultural Catchment Monitoring Programme 1990-2018

This section deals with the general development in nitrate leaching from 1990/91 to 2018/2019. Information on agricultural practises is derived from the Agricultural Catchment Monitoring Programme. This programme includes six small agricultural catchments situated in various parts of the country in order to cover the variation in soil type and rainfall and hence in agricultural practises (**Figure 4.3**). The farmers are interviewed every year about livestock, crops and fertilization and cultivation practises. Nitrate leaching is modelled for all fields in the catchments based on the information on agricultural practises and standard percolation values that are calculated on the basis of the climate for 1990-2010.

In 2019, 129 farmers participated in the investigation. Of all the investigated farms, 24 were cattle holdings, and five of these were registered as derogation farms. These derogation farms covered 15% of the total area in the Agricultural Monitoring catchments in 2018/19.

The modelled nitrate leaching from the agricultural area in the catchments was calculated for the period 1990 to 2019 (representing the hydrological years 1990/91 to 2019/20). The modelled leaching is shown in **Figure 4.6** as an average for sandy and loamy catchments, respectively.




Figure 4.6 Modelled nitrate leaching in a standard climate for the fields of the Agricultural Catchment Monitoring Programme 1990/91-2019/20.

Seen relative to the distribution of the main soil types in Denmark, the modelled nitrate leaching decreased by 43% during the period 1991 to 2003 due to the general improvement in agriculture and fertilization practises (Action Plan I+II). After 2003, there was a small increase in nitrate leaching, particularly on sandy soils, probably caused by suspension of the set aside obligation. At the national level, about 120,000 ha of set aside were cultivated in 2008 and 2009, leading to a change in crop rotation towards a higher leaching potential and a temporary increase in fertilizer application. For the loamy catchments, modelled annual nitrate leaching was relatively stable around 50 kg N ha⁻¹ during 2003-2013, after which it decreased by approximately 8 kg N ha⁻¹ in 2014 and 2015 and increased again to the level of 2003-2013 in 2016-2019. For the sandy catchments, the annual leaching of 81 kg N ha⁻¹ in 2003 was relatively low. After this year, the leaching increased to an interval of 81-93 kg N ha⁻¹ in the period 2004-2014. In 2015-2019 the annual leaching decreased to a lower level than in 2003 (77-81 kg N ha⁻¹). The lower leaching in these five years is mainly due to a higher share of catch crops after cereals and maize.

After 2003, the average annual modelled leaching for both loamy and sandy catchments varied between 63 and 67 kg N ha⁻¹. Nitrate leaching was lower in 2014 and 2015 (58 kg N ha⁻¹) due to a large area of catch crops, but increased to 60-62 kg N ha⁻¹ in the years 2016-2019, which is similar to the values for 2013 and 2014. The higher leaching in 2019 is due to a higher proportion of bare soil in the autumn especially in sandy dominated catchments. The extreme drought in the summer and autumn in 2018 caused unfavourable conditions for sowing and plant growth leaving bare soils as a higher proportion of the cropped area. No significant trend in leaching for the individual LOOP catchments was detected after the year 2003, except for a slight decreasing trend in the sandy catchment in Northern Jutland.

The purpose of the root zone modelling is to show the effects of measures introduced to mitigate nutrient losses from agriculture. The modelling is therefore carried out for normalised growth conditions, i.e. averaging the model output for a 20-year period: The model is run for each year in the 20-year period and model outputs are then averaged for the period. The climatic data used cover the period 1990-2010. Actual measurements of nitrate leaching will show higher annual variations than the climatic average of the modelled values as the measurements depend on the actual climate.

Certain forms of soil cultivation and ploughing of grass fields in autumn were prohibited as from autumn 2011. This circumstance is not considered in the leaching model due to lack of actual measurements that could otherwise have been applied in the model development. It is estimated that postponed soil tillage will reduce root zone leaching by 2,400 t N at the national level, corresponding to an average effect of about 1 kg N ha⁻¹ (Børgesen et al., 2013).

4.5 Measurements of nitrate in water leaving the root zone

In five of the six Agricultural Monitoring Catchments, soil water samples are collected regularly at 30 sites. One of the sites is covered by forest and is therefore not included in the data on nitrate concentrations measured in agricultural areas. Measurements were ceased on a sandy site in 2011 as the farmers did not want to participate in the monitoring. Two sites on a loamy catchment are located very close to the edge of the field, and tractor transport in and out of the fields and results in high damage to crops, possible uneven fertilizer application and very high values of measured nitrate leaching in some of the monitored years. Out of the remaining 27 sites on agricultural areas, 14 are located on loamy soils and 13 on sandy soils and the data on these are considered valid for use in the trend analysis of the loamy and sandy catchments. The samples represent the root zone water (approx. 1 m depth - 30 samples per year) and the upper oxic groundwater (1.5-5 m depth – 6 samples per year). To give an annual representative values for the nitrate leaching, the measured nitrate concentration is multiplied with the percolation in the sampling period. Samples are taken weekly in periods with percolation (autumn, winter and spring) and monthly in summer time when percolation is scarce or zero. Percolation values are modelled, as measurements of soil water content and flow in soil, covering soil variability at field level, are difficult to perform (Blicher-Mathiesen et al., 2014). The annual flowweighted nitrate concentration is calculated by dividing the annual percolation into the nitrate leaching.

Since the previous publication of the annual derogation report, inconsistencies in the precipitation time series has been detected. These inconsistencies affect the reported flow-weighted concentrations, as the precipitation time series are used for the calculation of percolation. Specifically, it was found that the relation between precipitation and stream runoff in the monitoring catchments was inconsistent before and after 2010, respectively. The precipitation is measured at several rain gauge stations and distributed to cover 10 x 10 km² grids by the Danish Meteorological Institute (DMI). The type of rain gauge station was changed from 2011 and also the number of stations decreased significantly. This explains some of the inconsistency related to measured discharge. DMI has delivered new precipitation data for the period after 2010, but all inconsistency in the data has not yet been resolved. In order to address the possible bias or inconsistency in the precipitation time series, we included an uncertainty in the precipitation data, which is reflected in the calculated percolation and flow-weighted nitrate concentration. This uncertainty was derived from an analysis of radar detected precipitation in five subplots within ten precipitation grids of 10x10 km². The error bars on the flow-weighted nitrate concentration in figure 0.6 and figure 0.10 represent this uncertainty from variation in precipitation on field level, but tabulated as an average uncertainty from ten precipitation grids (Blicher-Mathiesen et al., 2021).

The flow-weighted nitrate concentrations are shown as annual average values for loamy and sandy soils, respectively, for the period 1990/91-2018/19 (**Figure 4.7**).




Figure 4.7 Annual flow-weighted nitrate concentrations measured in root zone water (1 m below ground level) and annual average nitrate concentrations measured in upper oxic groundwater (1.5-5 m below ground level), the Agricultural Catchment Monitoring Programme 1990/91-2018/19. Error bars indicate variation in percolation as precipitation variated on local scale within a DMI 10 x 10 km² precipitation grid.

Generally, measured data on nitrate leaching from the root zone on only 27 sites cannot be used directly for estimating the effect of a single variable as the input of fertilizer or manure because of the high variability in actual fertilizer and manure practice between the monitoring fields and the measured years.

Instead, the data were used for the development of the nitrate leaching model, N-LES4, which was subsequently used for calculating the leaching from all the fields in the catchments relative to agricultural practises (**Figure 4.6**). The calculation of statistical trends for the monitoring period is done using flow-weighted nitrate concentrations.

General trend for nitrate concentrations in water leaving the root zone There is strong inter-annual variation in the measured nitrate concentrations due to differences in rainfall and temperature. Therefore, a long time series and a large number of measuring points are needed to detect any statistically significant trend. Such data series are available from the Danish Monitoring Programme. A statistical trend analysis – a Mann-Kendall test, incorporating annual variations in the mean annual flow-weighted nitrate concentrations for water leaving the root zone – showed that concentrations decreased significantly by 1.2 and 2.6 mg NO₃ l⁻¹ a⁻¹ for the measured sites on loamy and sandy soils, respectively, and for the whole 26-year monitoring period from 1990/91 to 2015/16. In loamy catchments, the measured nitrate concentrations in root zone water decreased from 61-155 mg NO₃ l⁻¹ in the 5-year period 1990/91-1994/95 to 37-66 mg NO₃ l⁻¹ in the 5-year period 2011/12-2015/16 and increased to 72, 48 and 108 mg NO₃ l⁻¹ in the three years 2016/17, 2017/18 and 2018/19, respectively. The high nitrate concentrations are seen in years with low percolation as observed on loamy soils in 2004/05, 2010/11, in 2016/17 and in 2018/19. In sandy catchments, the nitrate concentration decreased from 73-207 mg NO₃ l⁻¹ in the 5-year period 1990/91-1994/95 to 54-73 mg NO₃ l⁻¹ in the 5-year period 2011/12-2015/16 and increased to 93, 84 and 99 mg NO₃ l⁻¹ the three years 2016/17, 2017/18 and 2018/19, respectively (**Figure 4.6**).

After 2003/04 (Action Plan III + Green Growth), no statistically significant change in measured nitrate concentrations in soil water leaving the root zone has been recorded. However, before 2011/12, high concentrations were temporarily observed for sandy soils. This is most likely due to growth of crops with high leaching potential on these fields, such as turnover of grassland followed by cereals with no catch crops the following years, growing of maize and winter rape etc.

It should be noted that the measurements of nitrate leaching originate from a small number of sampling stations (27 stations). Furthermore, the measurements are affected by high crop yields, in particular in 2009, and effects of crop rotation, especially of grass in rotation. These conditions induce higher inter-annual variations than seen in the modelled nitrate leaching, which covers a larger area including approx. 129 farms (**Figure 4.6**).

In the upper groundwater (1.5-5.0 m below ground level), nitrate concentrations were lower than in the root zone water, indicating nitrate reduction in the aquifer sediment between the bottom of the root zone and the uppermost groundwater (**Figure 4.**7).

In loamy catchments, the measured nitrate concentrations in the upper oxic ground-water decreased from 41-46 mg NO₃ l⁻¹ in the 5-year period 1990/91-1994/95 to 29-37 mg NO₃ l⁻¹ in the 5-year period 2014/15-2018/19. In sandy catchments, the nitrate concentration decreased from 87-110 mg NO₃ l⁻¹ in the 5-year period 1990/91-1994/95 to 59-77 mg NO₃ l⁻¹ in the 5-year period 2014/15-2018/19.

Nitrate concentrations in water leaving the root zone from cattle holdings

Two to three of the monitoring sites belong to cattle holdings that, on average, used between 130 and 170 kg organic manure N ha⁻¹ in the period 2000/01 -2018/19 and four to five sites belong to holdings that, on average, used more than 170 kg organic manure N ha⁻¹ at the monitoring sites. Measurements of nitrate in water leaving the root zone are shown annually for each site for the period 2000/01-2018/19. At one of the sites, station "st 604", the manure input changed from a high annual input (>170 kg N ha⁻¹) until 2008 (data shown in **Figure 4.8**B and D), to a lower input (<170 kg N ha⁻¹) in the following years (data shown in **Figure 4.8**A and C). Suction cups at site "st 203" were re-established in 2012, which entails that no measurements for this site were available for 2012/13 and 2013/14. On this site, the input of manure increased to >170 kg N ha⁻¹ on average for a five year period monitored from 2012 and onwards. The annual manure input at site "st 202" changed to a much lower level from 2014, and the nitrate concentration in the root zone water is therefore included in **Figure 4.9**A and C from 2014 and onwards. For the site "st 206" the average five-year manure input increased from <130 kg N ha⁻¹ to >170 kg N ha⁻¹ in 2013 and is included in **Figure 4.8**B and D.




Figure 4.8 Measured nitrate concentrations in root zone water (1 m depth) with average application of 130-170 (A) and more than 170 kg organic N ha⁻¹ (B) at the sites (average application of organic manure N is shown in brackets). Figures for annual averages for the measured stations, average application of 130-170 (C) and more than 170 kg organic N ha⁻¹ (D). All data from the period 2000/01-2018/19 are shown.

The manure input at site "st 201" increased from 2013 and was included in **Figure 4.8**B and D and **Figure 4.8**B from this year. One site "st 406" stopped livestock production from 2017 and therefore this site is not included in the data from this year and onwards in **Figure 4.7** and **Figure 4.8**.

Annual variations in measured concentrations at the individual monitoring stations were expected, partly due to crop rotation and variations in yield and meteorological conditions.

The sites that annually received an average of 130-170 kg N in manure ha⁻¹ in the period 2000/01-2018/19 had high average nitrate concentrations in the six years 2005/06, 2008/09-2010/11, 2013/14 and 2015/16-2018/19 (**Figure 4.7A**). At some of the sites with an annual average manure application of more than 170 kg N ha⁻¹, nitrate concentrations were very high. For instance, this is seen at "st 604" in five out of six years between 2004/05 and 2009/10. However, other sites receiving high manure input had relatively lower soil water concentrations (**Figure 4.8**B). The average flowweighted nitrate concentrations in root zone water at four specific sites with an average manure application of 170-230 kg N ha⁻¹ varied between 64 and 110 mg NO₃ l⁻¹ for the recent five hydrological years (2014/15-2018/19).

High nitrate concentrations are most likely a result of crop rotation, especially turnover of clover grass in rotation, followed by cereals without catch crops or high N input to maize, and they cannot be linked to the level of manure input alone.

Phosphorus concentrations in the water leaving the root zone are shown in **Figure 4.9**. Generally, the concentrations varied between 0.005 and 0.050 mg PO_4 -P l⁻¹, irrespective of the use of organic manure. However, in one field receiving an average of 148 kg organic N ha⁻¹ ("st 608"), P concentrations were much more variable. The soil texture in this field is coarse sand and it is located in an area with high rainfall.




Figure 4.9 Measured phosphorus concentrations as dissolved orthophosphate (PO4-P) at soil water stations (1 m depth) with average application of 130-170 (A) and more than 170 kg organic N ha⁻¹ (B) at the sites (average application of organic manure N is shown in brackets). All data for the period 2000/01-2018/19 are shown.

4.6 The nitrogen flow to surface water in agricultural catchments

When percolating water leaves the root zone, it is partitioned into a component that discharges to surface water and a component that discharges to groundwater from where it will eventually – often some years later – drain into the streams. The pathways for water and nutrients in agricultural catchments are analysed in the Agricultural Catchment Monitoring Programme. Nitrate concentrations are measured in soil water and in water from tile drains from three loamy catchments and two sandy catchments.

The monitoring programme does not allow a specific evaluation of the effect of derogation farms on the nitrate transport in the streams since measurements at the catchment outlet integrate the effects of all activities in the catchment. However, the monitoring programme will provide an overview of the general trend for surface water, including the effect of any derogation farms in the catchment.

This chapter gives an overview of the nitrogen pathways in the hydrological cycle and describes the trends for nitrate in water for the period 1990-2019. Continued monitoring within the framework of the Agricultural Catchment Programme and the Stream Programme will provide indicators for the future development.

The hydrological pathways

An analysis of the water flow in the streams of the five agricultural catchments has shown that it can be conceptually divided into three components – rapid, intermediate and slow response to precipitation (**Table 4.8**). These components may be regarded as flow from the upper soil layers (including tile drainage), from the upper oxic groundwater and from deep groundwater.

Table 4.8 Partitioning of water flow in streams into three components – rapid, intermediate and slow responding water. The analysis included three loamy catchments and two sandy catchments (1989/90-2002/03).

	Flow response				
	Rapid	Intermediate	Slow		
Loamy catchments	41 %	16 %	43 %		
Sandy catchments	20 %	23 %	57 %		

Nitrate concentrations in the hydrological cycle (2014/15-2018/19)

(The arrows show the dominant pathways of the waterflow)




Figure 4.10 Measured mean of nitrate concentrations in the hydrological cycle in three loamy catchments and two sandy catchments in the Agricultural Catchment Monitoring Programme. Stream values for the two sandy catchments are for the two individual stream outlet. The values are calculated as an annual mean for the period 2014/15-2018/19.

In loamy catchments, the flow path is characterised by relatively rapidly responding water (from upper soil layers), whereas there is a larger proportion of slowly responding water (from deeper groundwater) in sandy catchments.

Figure 4.10 illustrates measurements of nitrate concentrations (mg NO3 l⁻¹) in soil root zone water, upper oxic groundwater (1.5-5 m below ground level) and in streams. When water percolates from the root zone to the upper groundwater, denitrification processes take place. Thus, nitrate concentrations in the upper groundwater are lower than in the root zone water. When the water passes through the deeper aquifers, it will usually reach the redox cline where the remaining nitrate will be removed by biological and geo-chemical reduction processes.

As sandy catchments are dominated by slow groundwater flow, the water discharging to the streams has been exposed to reduction processes. Thus, nitrate concentrations in the stream water are relatively low. In loamy catchments, the discharging water has mainly passed through the upper soil layers and through the drainage system where there is less nitrate reduction. Hence, nitrate concentrations in the streams are higher than in sandy catchments.

In this context, it should be noted that cattle farms, i.e. the derogations farms, are mainly located in the western and northern parts of Jutland characterised by sandy soils and deep groundwater flow, leading to high nitrate removal and low nitrogen concentrations in the streams.

Trends in nitrate concentrations in the hydrological cycle

The development in nitrate concentrations in root zone water, upper oxic groundwater and stream water is shown in **Figure 4.11**. Statistical analyses incorporating the annual variations showed that the nitrate concentration in water leaving the root zone decreased significantly by 1.2 and 2.6 mg NO₃ l⁻¹ a⁻¹ at the measured sites on loamy and sandy soils, respectively, for the 26-year monitoring period from 1990/91 to 2015/16. However, as mentioned before, nitrate concentrations increased to 72, 48 and 108 mg NO₃ l⁻¹ in the three years 2016/17, 2017/18 and 2018/19, respectively, on loamy soils and to 93, 84 and 99 mg NO₃ l⁻¹ in the corresponding three years, respectively, on sandy soils (see section 4.5). In the Stream Monitoring Programme, the development is analysed for a larger number of streams. This programme reported that during the period 1989-2019, in 51 agriculturally dominated catchments representing both loamy and sandy soils, there was an average reduction of 35 % (confidence interval: 27-39 %) of the total nitrogen transport (Thodsen et al., 2021).




Figure 4.11 Nitrate concentrations in root zone water, upper groundwater and in streams for three loamy catchments and two sandy catchments according to the Agricultural Catchment Monitoring Programme, 1990/91-2018/19.

References

Blicher-Mathiesen, G., Petersen, R.J., Holm, H., Houlborg, T., Rolighed, J., Andersen, H.E., Carstensen, M.V., Jensen, P.G., Wienke, J., Hansen, B. & Thorling, L. 2021. Landovervågningsoplande 2019. NOVANA. Aarhus Universitet, DCE – Nationalt Center for Miljø og Energi, 255 s. - Videnskabelig rapport nr. xxx

Blicher-Mathiesen, G., Andersen, H.E. & Larsen, S.E. (2014). Nitrogen field balances and suction cup-measured N leaching in Danish catchments. Agriculture, Ecos. Environm. 196, 69-75.

Kristensen, K., Waagepetersen, J., Børgesen, C.D., Vinther, F.P., Grant, R. & Blicher-Mathiesen, G. (2008). Reestimation and further development in the model N-LES - N-LES₃ to N-LES₄. DJF Plant Science No. 139.

Kronvang, B., Andersen, H.E., Børgesen, C., Dalgaard, T., Larsen, S.E., Bøgestrand, J. & Blicher-Mathiasen, G., (2008). Effects of policy measures implemented in Denmark on nitrogen pollution of the aquatic environment. Environmental Science & Policy 11, 144-152.

Thodsen, H., Tornbjerg, H., Rasmussen, J.J., Bøgestrand, J., Larsen, S.E., Ovesen, N.B., Blicher-Mathiesen, G., Kjeldgaard, A. & Windolf, J. (2019). Vandløb 2018. NOVANA. Aarhus Universitet, DCE – Nationalt Center for Miljø og Energi, 70 s. -Videnskabelig rapport nr. 353 <u>http://dce2.au.dk/pub/SR353.pdf</u>

Thodsen, H., Tornbjerg, H., Bøgestrand, J., Larsen, S.E., Ovesen, N.B., Blicher-Mathiesen, G., Rolighed, J., Holm, H. & Kjeldgaard, A. 2021. Vandløb 2019 - Kemisk vandkvalitet og stoftransport. NOVANA. Aarhus Universitet, DCE – Nationalt Center for Miljø og Energi, xx s. - Videnskabelig rapport nr. 416

SVANA (2016). Vandområdeplaner 2015-2021. Styrelsen for Vand og Naturforvaltning. Miljø- og Fødevareministeriet. https://mst.dk/naturvand/vandmiljoe/vandomraadeplaner/vandomraadeplaner-2015-2021/vandomraadeplaner-2015-2021/

Wiberg-Larsen, P., Windolf, J., Bøgestrand, J., Larsen, S.E., Thodsen, H., Ovesen, N.B., Bjerring, R., Kronvang, B. & Kjeldgaard, A. (2015). Vandløb 2013. NOVANA. Aarhus Universitet, DCE – Nationalt Center for Miljø og Energi, 50 s. - Videnskabelig rapport fra DCE - Nationalt Center for Miljø og Energi nr. 121 http://dce2.au.dk/pub/SR121.pdf

Windolf, J., Larsen, S.E., Thodsen, H., Bøgestrand, J., Ovesen, N. & Kronvang, B. (2011). A distributed modelling system for simulation of monthly runoff and nitrogen sources, loads and sinks for ungauged catchments in Denmark. J. Environmental Monitoring 13, 2645-2658.

5. Reinforced monitoring in areas characterized by sandy soils

This chapter is based on selected data from the National Monitoring Programme of Water and Nature (NOVANA), provided by the Danish Environmental Protection Agency, and data on derogation farm location, provided by the Danish Agricultural Agency.

5.1 Introduction

Prior to 2018, data on water quality in the derogation report was based on data from the national agricultural catchment monitoring programme. This programme combines detailed information on both agricultural practice and crop rotation as well as data on water quality in root zone water, uppermost groundwater and small local streams. Monitoring takes place in five agricultural catchments throughout the country, of which three are located in parts of Denmark characterized by loamy soils and two in the western part, where sandy soils predominate. The latest, relevant results from the programme are reported in chapter 4 of this report.

Due to the limited size of the area monitored within the national agricultural catchment monitoring programme, only very few derogation farms are located in the five catchments. The majority of derogation farms are found in the western part of Denmark, especially in the western part of middle and southern Jutland, as shown on the maps in chapter 2 of this report. This part of Denmark is also characterized by predominantly sandy soils.

The derogation decision from 2017 (2017/847/EU) introduced the requirement that water quality should be reported using data from reinforced monitoring. The reinforced monitoring is carried out on sandy soils and in an area that comprises fields belonging to at least 3% of all derogation farms. The derogation decision from 2018 (2018/1928/EU) and the latest derogation decision from 2020 (2020/1074/EU) specifies in Article 10 (2) that, in addition to the monitoring obligations in prior derogation decisions, "[...] *Reinforced monitoring of water quality shall be carried out in areas with sandy soils. In addition, nitrates concentrations in surface and groundwater shall be monitored in at least 3 % of all holdings covered by an authorisation.*"

5.2 Method

Selection of relevant monitoring stations

Besides the results from the national agricultural catchment monitoring programme (see chapter 4), which previously has formed the basis for annual reporting according to the derogation decision, Danish authorities also collect data through a number of other national monitoring programmes. As part of the "National Monitoring Programme of Water and Nature" (NOVANA), data from approximately 500 water quality stations in streams and rivers are collected on a regular basis. The primary purpose is to determine nutrient loads to sensitive recipients, i.e., coastal waters and lakes. Water samples from more than 1,000 groundwater monitoring stations are also analysed on a regular basis; the sampling frequency varies from several times annually to once during a multi-year period, according to the monitoring and reporting requirements of the Nitrates Directive and the Water Framework Directive. One of the usual parameters that both groundwater and surface water samples are analysed for is nitrate concentration.

Simultaneously, the Danish Agricultural Agency registers which fields belong to derogation farms. The approach is based on the identification of either surface water or groundwater monitoring stations located in close proximity to a field belonging to a derogation farm. More precisely, the GIS-analysis is based on the coordinates of the surface water or groundwater monitoring station as well as the surrounding area within a fixed 15-metre radius. This circle allows for an overlap between the position of the monitoring station and any fields in close proximity.

Only water course and groundwater monitoring stations located within 15 metres of a field registered to a derogation farm are selected. To determine whether this criterion is met, the latest registry data from the Danish Agricultural Agency is used.

If a groundwater monitoring well fulfils the location criterion but contains several monitoring stations at different depth ("multi-filter wells"), only one of these stations is selected; typically, the station that has the largest number of prior nitrate concentration samples.

Groundwater monitoring stations at a depth of 80 metres or more have been excluded from the data set, as data from the national groundwater monitoring ("GRUMO") programme shows that nitrate levels are no longer quantifiable (<1 mg/L) at these depths.

Only surface water monitoring stations that are part of the national programme monitoring "Transport of nutrients in streams" have been considered for the reinforced monitoring. A few mobile stations used for lake monitoring that would have fulfilled the proximity criterion have been excluded from the data set, as their locations typically change every year, making it impossible to create time series. Monitoring stations that have been installed in water courses to monitor the outflow from constructed wetlands have also been excluded.

In all, this selection method has identified a total of 46 monitoring stations. 28 stations of these (61 %) are groundwater monitoring stations, while 18 stations (39 %) are located in water courses (**Figure 5.1**). The distribution between station types is a direct consequence of the higher density of groundwater as opposed to surface water monitoring stations throughout Denmark.




Figure 5.1: Map showing the locations of the 46 monitoring stations selected as the reporting basis for the reinforced monitoring. The squares show the location of in total 28 groundwater monitoring stations at different depths – these may overlap due to the scale of the map. The circles show the location of the 18 water course monitoring stations. Grey shading indicates all fields belonging to Danish derogation farms.

The majority of derogation farms are located in the western part of Denmark, especially the western, northern and southern parts of the peninsula of Jutland, also illustrated in chapter 2 of this report. These parts of the country are characterized by sandy soils, whereas loamier soils dominate the more eastern parts of the country. Consequently, the described approach of linking the locations of monitoring stations to fields belonging to derogation farms results in a considerable enlargement of the data basis for reporting of water quality in sandy areas.

The geological map in **Figure 5.2** below illustrates the soil substrates throughout Denmark.




Figure 5.2 Geological map of Denmark showing the substrates that are the basis for soil development. Modified from a map produced by GEUS. The legend is only available in Danish, but the four main soil substrate types that can be categorized as "sand" have been marked with an "S" in the legend.

Coverage of Danish derogation farms

The locations of the 46 monitoring stations have been linked to 60 fields, which in turn belong to 40 different derogation farms. The number of stations exceeds the number of derogation farms, as some farms own several fields in proximity to a monitoring station, and as there may be more than one monitoring station located very close to a given field. Out of the 40 farms, 20 are subject to the reinforced monitoring due to the proximity of their fields to a water course monitoring station, while 21 farms (one both water course and groundwater) are included owing to proximity to groundwater monitoring stations. The total number of farms encompassed by the reinforced monitoring corresponds to approximately 3.1 % of all holdings that make use of the derogation.

5.3 Characterization of monitoring stations and data analysis

Groundwater

The selected groundwater monitoring stations are located at depths below the surface ranging from 1.75 m to 72 m^9 . The majority of the stations monitor water quality in comparatively shallow groundwater, at an average depth of 21.4 m and a median

⁹ Based on data available up until 2018, the deepest selected monitoring station was located at 62 m below the surface. Data from the deeper groundwater is expected to be included in future reports, when water from these selected stations will be sampled and analysed for nitrate again.

depth of 13.45 m. Of the selected groundwater monitoring stations, 35.7 % the samples are of very shallow groundwater from a depth of less than 10 m. 28.6% are located from 10-20 metre below surface and the rest 35.7 % of the stations are located from 20-80 metres.

All of the groundwater monitoring stations will be sampled at least once per year. Historic data since 2002 – the year Denmark obtained a derogation from the Nitrates Directive for the first time – have been included to the extent that they are available. If groundwater has been sampled more than once per year, the average annual nitrate concentration has been calculated for this station for each respective sampling year.

For the purpose of presenting the data in the results section below, the stations have been grouped into three different categories, stations at a depth of less than 10 m below surface, stations at 10 to almost 20 m depth and stations at 20 m depth or deeper. Annual average nitrate concentrations have been calculated for each depth category for each year since 2002, based on the actual number of stations sampled in the respective year.

Surface water

The monitored water courses vary considerably in size and flow rate. The widths of the water courses at the monitoring station vary from 2 m to 23 m. The average water course width at the monitoring station is 7 metres, while 5 out of the 18 stations are located in small streams of less than 5 metres' width.

Samples from water courses are generally analysed for Nitrite- and Nitrate-Nitrogen (N). Nitrite-N-concentrations are typically negligible, and under this assumption, nitrate concentrations in the water samples could be calculated by multiplying the Nitrate-N concentration by a factor of 4.4268. In this chapter, the surface water concentration is generally given in Nitrite- and Nitrate-Nitrogen. Historic data since 2002 is included to the extent that it is available. Only data from monitoring stations that have been sampled at least 9 times annually in the period before 2017 are displayed in the results. In 2019, each water course monitoring station has been sampled more frequently, from 12 to 18 times annually.

For the purpose of presenting the data in the results section, stations have been grouped based on the approximate width of the water course at the sampling station site into three different categories, as also displayed in figure 5.1: less than 5 m, 5 to 10 m and more than 10 m width. Average nitrate concentrations have been calculated for each category for each year since 2002, based on the actual number of stations sampled in the respective year.

As a consequence of the political agreement on the Food and Agricultural Package from December 2015, the number of water course monitoring stations has been significantly increased. 8 out of the 18 water course stations selected for the reinforced monitoring were established in 2016 as a consequence of the agreement, and now, it is decided to continue the monitoring.


5.4 Results and Discussion


Groundwater


Figure 5.3 shows the nitrate concentration of each groundwater monitoring station selected for reinforced monitoring, as well as the average nitrate concentrations per sampling year for the period 2002 to 2019 for each of the depth categories. The quality limit value of 50 mg nitrate per litre is also shown.

The data generally shows great variability in nitrate concentrations from one year to another in water samples from individual monitoring stations. Especially in the shallowest groundwater (Figure 5.3A), absolute concentration changes of up to more than 80 mg nitrate per litre can be observed from one sampling year to the other.

The average nitrate concentration remains below the quality limit value for each depth category throughout the whole period 2002 until 2019, with the exception of 2010 for the deepest category. However, this value is only based on a single monitoring station, as none of the others was sampled in 2010.









depth categories of groundwater stations: (A) stations at less than 10 m depth; (B) stations at 10-20 m depth and (C) stations at 20 m depth and deeper below the surface. A red dashed line at 50 mg nitrate per litre is inserted in each figure.

No clear trend in the average nitrate concentration can be observed over time for any of the three depth categories. Due to the limited number of stations and samples per year, the annual average values are highly influenced by the variability in nitrate concentration in the water sampled from some individual stations.

Table 5.1 shows the average nitrate concentration of all stations for each year in the period 2002 to 2019, irrespective of their depth and the number of stations sampled (n) in the respective year that form the basis of this calculation. The annual average nitrate concentration varies between 20.9 mg/L, as sampled in 2003 (n=11), and 41.9 mg/L in the groundwater samples from 2009 (n=13).

	l average nitrate concen riod 2002-2019 and num	tration of all stations in reinforced mon- ber of stations sampled
Sampling vear	Average nitrate concentration	Number of sampled stations (n)

Sampling year	Average nitrate concentration [mg/L]	Number of sampled stations (n)		
2002	29.2	11		
2003	20.9	11		
2004	27.0	10		
2005	37.0	18		
2006	38.1	18		
2007	37.8	18		
2008	32.1	16		
2009	41.9	13		
2010	40.7	13		
2011	28.7	18		
2012	28.8	23		
2013	29.2	18		
2014	28.8	23		
2015	35.1	21		
2016	35.2	22		
2017	31.9	26		
2018	30.5	29		
2019	26.9	28		

When calculated across the entire period from 2002 to 2019, the (non-weighted) mean value of the annual average concentrations is 30.0 mg/L. The 2019 average is lower than the mean value for the whole 2002-19 period.

Surface water

Figure 5.4 shows the Nitrite- and Nitrate-Nitrogen concentration of the individual water course monitoring stations selected for reinforced monitoring, as well as the average nitrate concentrations per sampling year for the period 2002 to 2019 for each of the width categories. The quality limit value for groundwater of 50 mg nitrate per litre, which corresponds to approximately 11.3 mg Nitrate-N per litre, is also shown.




Figure 5.4: Nitrite- +Nitrate-Nitrogen (N) concentration of the individual surface water monitoring stations selected for reinforced monitoring, as well as the average nitrate concentrations per sampling year for the period 2002 to 2019 for each of the three width categories (determined at sampling site): (A) less than 5 m wide; (B) 5 to 10 m wide and (C) wider than 10 m. A red dashed line is inserted in each figure at 11.3 mg Nitrate-N/L, corresponding to approx. 50 mg nitrate per litre.

At the level of the individual monitoring station, nitrite- + nitrate-nitrogen concentrations can vary significantly from year to year mainly due to variation in amount and timing of precipitation. The standard deviation in absolute concentration are 1.45 mg/l, corresponding to 6.4 mg nitrate per litre. Nevertheless the year-to-year variations are not as pronounced as those seen in groundwater samples.

For all water course categories it is, however, important to underline that the N transport is not determined by the nitrogen concentration alone, but also by the water flow in the water course, which can significantly vary due to the specific and local weather conditions of a given year. In low flow rate situations, nitrogen levels may be relatively high while total N transport remains unchanged, and vice versa. As smaller water courses typically have a smaller catchment area than rivers, variations in local weather conditions are expected to have a greater impact on the nitrogen concentration in water sampled from small water courses.

For all individual water course monitoring stations, nitrate-N concentrations remain well below the quality limit for groundwater and drinking water throughout the whole period from 2002 to 2019. Absolute concentrations tend to be higher in the smaller water courses than in the larger ones, which is likely to be a result of nitrate being removed through natural processes along the course of the water. Overall, the annual average for each category has been steadily decreasing over the last 10 years of the period shown.

Table 5.2 shows the annual average nitrite- and nitrate-N concentration in water sampled at all water course stations – irrespective of their width – and the number of stations sampled in the respective year (n).

Sam- pling year	Average nitrite- +ni- trate-N concentration [mg/L]	Number of sam- pled stations (n)
2002	4.1	5
2003	3.8	5
2004	4.4	7
2005	4.0	7
2006	4.4	10
2007	4.1	10
2008	3.9	10
2009	3.9	10
2010	3.8	10
2011	3.3	11
2012	3.4	11
2013	3.4	11
2014	3.3	11
2015	3.3	11
2016	3.3	11
2017	3.0	20
2018	3.0	20
2019	2.9	18

Table 5.2: Annual average nitrite- + nitrate-N concentration in water sampled at all stations selected for reinforced monitoring, as well as the number of stations sampled in each year

The annual average nitrite- + nitrate-N concentration has decreased from 4.4 mg/L in the early years of the reported period (e.g. 2006, n=10) down to 3.3 mg/L in 2016 (n=11). Since 2017, 8 additional water course monitoring stations have been established, improving the data basis significantly. Hence, it is now possible to follow the development in water courses, which have been equipped with new monitoring stations as a consequence of the political agreement on the Food and Agricultural Package from December 2015. Despite the significant increase in number of monitoring stations, which are considered in the reinforced monitoring, the average nitrite- + nitrate-nitrogen concentrations for the different water course stations was 2.9 mg/L, i.e. at the same level as in 2018.

General discussion

It is important to highlight that the reinforced monitoring does not provide data that can be used to examine any potential effect on water quality that might be the result of the use of the derogation. A range of other fluctuating factors influence nutrient concentrations in the aquatic environment, and as such, it would not be possible to identify or isolate such an effect.

Because the reinforced monitoring method is based on linking the locations of monitoring stations to fields belonging to derogation farms in a two-dimensional way: The approach does not account for the actual catchment area and subsurface water paths for the respective monitoring stations. Hence, it is only to a very limited degree possible to get a picture of the effects of land use on surface water and groundwater quality. A clearer picture would require a catchment-based approach, which takes into account that water quality in the recipient water is affected by land use in the whole catchment area.

The present method does not include a reference group of monitoring stations that are not located in proximity to fields belonging to derogation farms. However, by including the data from this selected set of the surface water and groundwater monitoring stations, the data basis for water quality in sandy areas has been considerably enlarged from the two sandy catchments within the national agricultural catchment monitoring programme (see chapter 4), which formed the basis for reporting prior to 2018 and still provides comprehensive data on land use at farm-level.

6. Indicator and monitoring system for application of phosphorus in Denmark

Ministry of Environment of Denmark

6.1 Introduction

In consultation with the European Commission, the Ministry of the Environment and Food (since November 2020 the Ministry of Environment) has agreed that Denmark must monitor the use of phosphorus (P) in organic fertilizer and commercial fertilizer, so that it is ensured that the average use does not exceed the national phosphorus ceiling. The monitoring is based on data from the fertilizer accounts, which is available approximately one year after a planning period is completed, when the farmers submit their fertilizer accounts to the Danish Agricultural Agency. The first planning period with limiting phosphorus use by specific ceilings at farm level was 2017/2018. Deadline for fertilizer accounting for this first period was 1 April 2019. Hence, the current report contains data for two planning periods with phosphorus ceilings.

As a supplement to monitoring, it has been agreed that an "indicator system" must be established, where data from the NOVANA monitoring program in Agricultural Catchments (LOOP) in combination with available data on livestock production and sales of fertilizer and other phosphorus sources can provide an updated overview of the average amount of phosphorus used in Danish agriculture.

These results from the P monitoring and indicator system, respectively, should be compared with the phosphorus ceilings. In this connection, it was agreed, that the total amount of phosphorus used should be divided by the total agricultural area in order to calculate the average fertilizer rate per year per ha on agricultural land. No requirement has been set for the first planning period 2017/2018, but in 2018 (planning period 2018/2019) the average use should be below 34.7 kg P/ha. In 2019 (planning period 2019/2020) the average use must be below 34.1 kg P/ha. If the average use exceeds 34.1 kg P/ha in 2019, the phosphorus ceilings must be lowered.

6.2 Results from the P monitoring system

The Danish Agricultural Agency compiled data from the fertilizer accounts with data from the planning period 2018/2019. The compiled data has not been processed or checked thoroughly for exorbitant values and other "noise", e.g. typos. If there are exorbitant values, it is estimated that only extremely high values in a few fertilizer accounts can have an important influence on the overall results, so the results represent a "worst case" scenario of phosphorus use.

	Produced P	Used P
	(tons)	(tons)
Poultry/fur	5,200	
Finishers	10,900	
Sows and piglets	8,600	
Cattle (non-derogation)	10,700	
Cattle (derogation)	7,400	
Manure – Total	42,800	44,400

Table 6.1: Compiled data from fertilizer accounts 2018/2019 (rounded numbers)

Waste and other P	3,600	3,200
Manure + waste		47,600
Chemical fertilizers		14,800
Used P – Total		62,300
		Mio. ha
Agricultural area		2,600
Harmony area		2,500
Average P-ceiling in 2018/2019		34. 7
kg P/ha agricultural area		23.8
kg P/ha harmony area		25.2

6.3 Results from P indicator system

The following table shows the phosphorus inputs as reported in the NOVANA report "Land Surveillance Survival 2019" from July 2021¹⁰. The table shows an increase as expected in the use of phosphorous in 2017 due to the increase in the P-ceiling from 2016 to 2017. In the coming years the P-ceiling will be decreased back to a lower level, so the increase in the use of phosphorous is not expected to continue.

	2012	2013	2014	2015	2016	2017	2018	2019
Use of P (1,000 tons) in different inputs:								
- Chemical fertilizer	11.8	11.3	13.0	13.3	13.3	20.8	14.8	14.6
- Livestock manure	45.8	45.3	46.1	46.1	44.3	43.0	44.3	44.9
- Seed	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1,0
- Sludge	2.4	2.4	2.4	2.4	2.4	2.4		
- Waste from industry	3.1	3.1	3.1	3.1	3.1	3.1		
- Other organic ferti- lizer ¹¹							2.8	3.1
- Deposition	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Total use of P	64.4	63.4	65.9	66.2	64.4	70.5	63.2	63.9
Agricultural area (1,000 ha)12	2,679	2,671	2,661	2,633	2,625	2,610	2,602	2,613
kg P/ha in average	24.0	23. 7	24. 7	25.1	24.4	27.0	24.3	24.4
kg P/ha P-ceiling					[32.2] ¹³	35.2	34. 7	34.1

Table 6.2: The use of P-input in Danish agriculture in 2012-2019¹⁰

¹⁰ Source: Blicher-Mathiesen *et al.* (2021): *Landovervågningsoplande 2019*. Aarhus University. <u>http://dce2.au.dk/pub/SR451.pdf</u>

¹¹ From 2018 onwards, amount of other organic waste, such as sludge and waste from industry, is derived from the fertilizer accounts.

¹² Agricultural area for the years 2016, 2017 and 2018 has been updated after submission of the Derogation Report 2019.

¹³ This figure indicates the average phosphorus protection level in 2016 expressed as a theoretical P-ceiling, before the P-ceilings were introduced, and is included for comparison. In the dialogue with the EU Commission, it was expected that the development in livestock production should be monitored via data from the CHR register, since Denmark previously prepared an annual status on the size of livestock production in various catchments. This annual status is now done instead on the basis of the fertilizer accounts, which is why the best data material on the development in livestock production is the annual status of the livestock population, which is made by Statistics Denmark. Statistics Denmark's information on livestock in 2017-2019 can be seen in **Table 6.3**.

	Number of animals 2017	Number of animals 2018	Number of animals 2019	% change in total number of ani- mals 2017-2019
Number of all kinds of cattle and dairy cows on all farms	1,545,417	1,540,446	1,491,433	-3.50 %
Number of all kinds of pigs on all farms	12,307,667	12,781,247	12,298,993	-0.07 %
Number of all kinds of poultry on all farms	21,483,698	19,973,164	23,059,881	7.89 %
Number of all kinds of mink on all farms	3,429,472	3,379,931	2,489,751	-27.80 %

Table 6.3: The development in the livestock production according to Statistics Denmark in 2017, 2018 and 2019¹⁴

The manure production based on data from the fertilizer accounts shows that 12 % of the total manure production comes from mink and poultry, 46 % from pigs and 42 % from cattle. There are no signs that indicate that a considerably larger amount of live-stock manure will be produced in 2020, and that the average phosphorus application in Denmark will exceed 25-28 kg P/ha, as the phosphorus ceiling from 2018 onwards will be reduced continuously. This level will be well below the average phosphorus ceilings of 35.2 kg P/ha in 2017, 34.7 kg P/ha in 2018, 34.1 kg P/ha in 2019, 33.2 kg P/ha in 2020 and further reductions set for 30-31 kg P/ha in 2025.

¹⁴ Data from Statistics Denmark: for cattle, pigs, poulty and mink: <u>https://www.statistikbanken.dk/10472</u>

7. Targeted catch crops scheme and targeted nitrogen regulation

Kari Lundsager and Marie Dam, The Danish Agricultural Agency, Ministry of Food, Agriculture and Fisheries of Denmark, October 2020

As part of the political agreement on the Food and Agricultural Package of December 2015, the reduction of the nitrogen application standards was removed. It was also agreed to develop a new nitrogen regulation, the "targeted nitrogen regulation", which was to be implemented in 2019.

The Danish government introduced an intermediate initiative, the "targeted catch crops scheme", to reduce N-losses through promoting the establishment of additional catch crops in 2017 and 2018. The scheme was designed to protect both groundwater bodies and coastal waters. The scheme consisted of a voluntary phase, where farmers applied for participation in the scheme, and a subsequent mandatory requirement for catch crops if the voluntary scheme did not reach the predefined targets. The latter requirement was uncompensated whereas the voluntary part was compensated with de minimis support.

In November 2017, a political agreement for targeted nitrogen regulation was reached and would be implemented from 2019. The targeted nitrogen regulation is similar to the targeted catch crops scheme in many ways. The most significant difference is the introduction of the possibility to use alternative measures to catch crops. Conversion factors are used to secure that the alternatives have the same effect as catch crop. Like the targeted catch crops scheme, the targeted nitrogen regulation is divided into a voluntary and a mandatory part. The targeted nitrogen regulation was subsidised by de minimis in 2019 and by RDP funds in 2020.

After the application deadline in the voluntary crop scheme, the farmer is bound by any commitment made, either through catch crops or alternatives, as well as by any additional catch crop requirement imposed through the mandatory round. The farmer will not be able to opt out of any of these requirements without consequences. The voluntary and obligatory targeted catch crops or alternatives must be additional to the national mandatory requirement for catch crops on 10.7 or 14.7% of the farm's crop base area, and they cannot be established on the area used for catch crops to meet the EFA requirement under direct payments.

If the farmer opts out afterwards or non-compliance is detected during control, the nitrogen quota for the farm (calculated on the basis of the composition and distribution of crops and the soil and crop-specific nitrogen standards) is reduced corresponding to the non-compliance with the voluntary and/or mandatory requirement and according to a conversion factor between the nitrogen reduction effect of catch crops and the nitrogen quota reduction for the planning period. This quota reduction will contribute to meeting the objectives of the Nitrates Directive. Furthermore, if the reduced nitrogen quota is exceeded, the farmer will be in breach of the Fertilizer Act and will be sanctioned accordingly cf. Annex III point 1.3 of the Nitrates Directive. This is similar to the current practice for the general catch crop requirements and additional catch crop requirements for holdings using organic manure.

In 2019, the targeted nitrogen regulation contributed to a nitrogen reduction of 1,174 tons in coastal waters, including reductions of nitrogen leaching to the groundwater. Further, in 2019, it was decided by political agreement to increase the effort of the

targeted nitrogen regulation in 2020 for additional contribution to meet the objectives of The Water Framework Directive. In 2020, the targeted nitrogen regulation will thus contribute to a nitrogen reduction of 3,514 tons in coastal waters.

7.1 Results from 2017 to 2020

Prior to 2017 and 2018, respectively, the ministry calculated the need for further nitrates efforts for each of the years, which can be expressed as the amount of additional catch crops required in the individual water catchment areas, in terms of hectares and as a percentage of the crop base area. The calculation is based on the estimated need for reductions in the nitrates contents of groundwater bodies and coastal waters, adjusted by the estimated soil nitrates retention in the water catchment area. In 2019 and 2020, the targeted nitrogen regulation was dimensioned to comply with the Danish implementation of The Water Framework Directive.

In 2017, the need for further nitrogen efforts was calculated to 137,560 ha. By the application deadline, the farmers had applied for a total of 144,220 ha of catch crops. However, the geographical distribution of the catch crops was not optimal in relation to the efforts needed. Calculations revealed that an additional **3,253** ha catch crops were needed in order to reach the target. It was decided politically to postpone the residual effort until 2018.

In 2018, the need for further nitrogen effort was calculated to 114,300 ha catch crops (including the postponed 3,253 ha). By the application deadline, the farmers had applied for a total of 105,000 ha of catch crops. It was furthermore decided to postpone the effort related to aquaculture (fish farming, mariculture, etc.), as extensions of existing aquaculture facilities had not been approved. Calculations revealed that an additional **3,000** ha catch crops were nevertheless needed in order to reach the target. This has been implemented as a mandatory uncompensated requirement in 2018.

In 2019, the need for nitrogen efforts in targeted nitrogen regulation was calculated to 138,200 ha of catch crops. By the application deadline, the farmers had applied for 139,350 ha of catch crops (and alternatives). Calculation revealed that an additional 275 ha were needed to reach the set effort. The reason was the geographical distribution of the catch crops, which was not optimal. It was decided politically to postpone this insignificant residual effort to 2020

In 2020 the need for nitrogen efforts in targeted nitrogen regulation was calculated to 373,000 ha of catch crops and included the residual effort from 2019. By the application deadline, the farmers had applied for 370,000 ha of catch crops (and alternatives). Some applications had to be dismissed, as the set effort for the individual water catchment areas was already reached. Calculations of the geographically specific retention disclosed that an additional 12,493 ha were needed to reach the set national nitrogen reduction effort. Consequently, this was implemented as a mandatory uncompensated requirement in 2020. Excluding a minor residual effort of 350 ha of catch crops, which was decided politically to postpone.

8. Conclusions

8.1 Cattle holdings and controls on farm level

In the planning period 2018/2019, a total of 1,284 cattle holdings made use of the derogation. This corresponds to 3.9 % of the total number of agricultural holdings in Denmark. These holdings produced 39.1 million kg N corresponding to 17.8 % of the total kg N produced. The arable land encompassed by the derogation in year 2018/2019 was 195,804 hectares corresponding to around 8.1 % of the total arable area. Compared to the previous reporting period, in 2018/2019 there has been a decrease in the number of farms and the number of hectares encompassed by the derogation. The average livestock size was 30,475 kg N/holding in 2018/2019.

In January – February 2020, 85 inspections of compliance with the derogation management conditions were carried out. All of these inspections were closed without remarks.

For the year 2017/2018, 84 inspections (0.3 % of all Danish holdings) at the holding were made concerning compliance with the harmony rules (amount of livestock manure applied per hectare). 84 of the inspected farms used the derogation. 60 of these inspections were closed without remarks. Three holdings had minor violations and they received a warning. 21 holdings are still under investigation.

All 33,056 fertilizer accounts submitted in 2017/2018 (100 %) were automatically screened by the IT-system according to normal procedure. Of these, 1,461 (4.4 %) were subject to administrative control. In all, 139 of these holdings used the derogation. Of the inspections of derogation farms, 89 (64.0 %) were closed without remarks, 3 (2.2 %) were closed with remarks and 47 (33.8 %) are still under investigation.

In total, approximately 7.0% of derogation farms were selected for physical inspections. In total, more derogation farms have been subject to controls due to the aforementioned administrative controls. As holdings are automatically selected - based on a previously agreed set of risk criteria - for both physical inspections and administrative control, the Danish Agricultural Agency has no direct influence on the share of holdings using the derogation that are inspected each year. Therefore, the share of derogation farms that in some way has been subject to controls varies from year to year.

8.2 Water quality

General conclusions from the Agricultural Catchment Monitoring Programme Measured nitrate concentrations in root zone soil water (1.0 m below ground level) decreased steadily from 1990/91 to 2015/16, albeit with annual variations. From 2016/17, the concentration generally increased, except on loamy soil in 2017/18. In loamy catchments, the measured nitrate concentrations in the root zone soil water decreased from 61-155 mg NO3 l⁻¹ in the 5-year period 1990/91-1994/95 to 37-66 mg NO3 l⁻¹ in the 5-year period 2011/12-2015/16. However, the concentration changed to 72, 48 and 108 mg NO3 l⁻¹ in the three years 2016/17,2017/18 and 2018/19, respectively. High nitrate concentrations are seen in years with low percolation as observed on loamy soils in 2004/05, 2010/11, 2016/17 and in 2018/19. In sandy catchments, the nitrate concentrations were 73-207 mg NO3 l⁻¹ in the 5-year period 1990/91-1994/95 and decreased to 54-73 mg NO3 l⁻¹ in the 5-year period 2011/12-2015/16, but increased to 93, 84 and 99 mg NO3 l⁻¹ in the three years 2016/17, 2017/18 and 2018/19, respectively. Measured nitrate concentrations in the upper oxic groundwater (1.5-5.0 m below ground level) decreased to a level below the limit of 50 mg NO3 l^{-1} for loamy catchments and to a level between 59-77 mg NO3 l^{-1} for the two sandy catchments in the period 2014/15-2018/19.

Measured average flow-weighted nitrate concentrations in root zone water at three to four specific sites receiving, on average, 170-230 kg organic manure N ha⁻¹ varied between 24-110 mg NO3 l⁻¹ in the hydrological years in the period 2014/15-2018/19.

Modelling of nitrate leaching for three loamy and two sandy catchments:

- For the loamy catchments, modelled annual nitrate leaching was relatively stable around 50 kg N ha⁻¹ during 2003-2013, after which it decreased by app. 8 kg N ha⁻¹ in 2014 and 2015 and increased again to the level of 2003-2013 in 2016-2019.
- For the sandy catchments, the modelled annual leaching of 81 kg N ha⁻¹ in 2003 was relatively low. After this year, the leaching increased to an interval of 81-93 kg N ha⁻¹ in the period 2004-2014, but decreased a level of 77-81 kg N ha⁻¹ in 2015-2019. The lower leaching in these five years is mainly due to a higher share of catch crops after cereals and maize.
- The higher leaching in 2019 is due to a higher proportion of bare soil in the autumn, especially in sandy dominated catchments. The extreme drought in the summer and autumn in 2018 caused unfavourable conditions for sowing and plant growth leaving a higher proportion of the cropped area as bare soils. No significant trend in leaching for the individual LOOP catchments was detected after the year 2003, except for a slight decreasing trend in the sandy catchment in Northern Jutland.

8.3 Targeted catch crops and targeted nitrogen regulation

For the year 2017, a total of app. 144,000 ha voluntary targeted catch crops were established, and a further effort of 3,250 ha were postponed to 2018. In 2018, a total of app. 105,000 ha voluntary catch crops were established, and in addition a mandatory effort of app. 3,000 ha has been applied (uncompensated). In 2019, first year of targeted nitrogen regulation, a total of 139,350 ha voluntary catch crops (or alternatives) were established, a further effort of 275 ha was postponed. In 2020, the targeted nitrogen regulation continued with a total of app. 370,000 ha voluntary catch crops established, and an additional mandatory effort of app. 12,500 ha applied (uncompensated). A further effort of 350 ha was postponed.

8.4 The reinforced monitoring

The reinforced monitoring does not provide data that can be used to examine any potential effect on water quality that might be the result of the use of the derogation. A range of other fluctuating factors than proximity to a derogation farm influence nutrient concentrations in the aquatic environment. However, by including the data from the selected set of the surface water and groundwater monitoring stations, the data basis for water quality in sandy areas is considerably enlarged. The total number of farms encompassed by the reinforced monitoring corresponds to approximately 3.1% of all holdings that make use of the derogation.

8.5 The phosphorus indicator and monitoring system

Neither the phosphorus indicator nor the P monitoring system indicate that the average phosphorus application in Denmark exceeds the average phosphorus ceilings of 34.7 kg P/ha in 2018 and 34.1 kg P/ha in 2019, respectively. There is currently also no risk for exceeding future P-ceilings, which are reduced compared to current level.



Ministry of Environment of Denmark - Departement Slotsholmsgade 12 DK - 1216 Copenhagen K

www.mim.dk